# Assessment of controlled release buccal inserts containing pilocarpine hydrochloride: a multicentre, double-blind, placebo-controlled, randomised, cross-over study in patients diagnosed with Sjögren's Syndrome

Recruitment status	Prospectively registered
No longer recruiting	☐ Protocol
Overall study status	Statistical analysis plan
Completed	Results
Condition category	Individual participant data
Musculoskeletal Diseases	Record updated in last year
	No longer recruiting  Overall study status  Completed  Condition category

# Plain English summary of protocol

Not provided at time of registration

# Contact information

# Type(s)

Scientific

#### Contact name

Mr John Hamburger

#### Contact details

The University of Birmingham School of Dentistry St Chad's Queensway Birmingham United Kingdom B4 6NN +44 (0)121 237 2888 j.hamburger@bham.ac.uk

# Additional identifiers

EudraCT/CTIS number

**IRAS** number

#### ClinicalTrials.gov number

#### Secondary identifying numbers

PILO-BUC-002

# Study information

#### Scientific Title

Assessment of controlled release buccal inserts containing pilocarpine hydrochloride: a multicentre, double-blind, placebo-controlled, randomised, cross-over study in patients diagnosed with Sjögren's Syndrome

#### **Study objectives**

Not provided at time of registration

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Not provided at time of registration

#### Study design

Multi-centre double-blind placebo-controlled randomised cross-over study

#### Primary study design

Interventional

# Secondary study design

Randomised controlled trial

#### Study setting(s)

Not specified

# Study type(s)

Treatment

#### Participant information sheet

## Health condition(s) or problem(s) studied

Sjögren's Syndrome

#### **Interventions**

This trial is a 'cross-over' study with four treatment cycles. For each treatment cycle, patients will be provided with 12 buccal inserts, containing one of the following: 2.5, 5 or 10 mg pilocarpine or placebo. A treatment cycle is four days long, however, there will be a three day washout period between each new dose. The buccal insert is a hydrogel polymer with pilocarpine dispersed throughout the matrix. Each buccal insert should be inserted high in the upper buccal sulcus between the buccal mucosa and gingivae towards the back of the mouth and remain in situ for at least three hours. One buccal insert should be inserted three times daily in the period between meals.

#### Intervention Type

Other

#### Phase

**Not Specified** 

#### Primary outcome measure

Not provided at time of registration

#### Secondary outcome measures

Not provided at time of registration

#### Overall study start date

01/01/2002

#### Completion date

31/12/2002

# Eligibility

#### Key inclusion criteria

Approximately 30 patients with primary or secondary Sjögren's Syndrome, as diagnosed by the modified European Diagnostic Classification, will be enrolled into the study, to accrue 25 evaluable patients

#### Participant type(s)

**Patient** 

#### Age group

Adult

#### Sex

Both

## Target number of participants

30

#### Key exclusion criteria

Not provided at time of registration

#### Date of first enrolment

01/01/2002

#### Date of final enrolment

31/12/2002

# Locations

Countries of recruitment

## England

**United Kingdom** 

Study participating centre
The University of Birmingham
Birmingham
United Kingdom
B4 6NN

# Sponsor information

#### Organisation

Controlled Therapeutics (Scotland) Ltd

## Sponsor details

1 Redwood Place Peel Park Campus East Kilbride United Kingdom G74 5PB

#### Sponsor type

Industry

#### **ROR**

https://ror.org/03e9kb581

# Funder(s)

#### Funder type

Industry

#### **Funder Name**

Controlled Therapeutics (Scotland) Ltd (UK)

# **Results and Publications**

# Publication and dissemination plan

Not provided at time of registration

Intention to publish date
Individual participant data (IPD) sharing plan

**IPD sharing plan summary**Not provided at time of registration