# Effectiveness of Bupivacaine in Pain Management following Breast Augmentation

Submission date	Recruitment status	Prospectively registered
28/09/2007	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
28/09/2007	Completed	Results
Last Edited	Condition category	Individual participant data
30/08/2012	Signs and Symptoms	<ul><li>Record updated in last year</li></ul>

# Plain English summary of protocol

Not provided at time of registration

# Contact information

# Type(s)

Scientific

#### Contact name

Mr S Jayagopal

#### Contact details

Countess of Chester Hospital NHS Foundation Trust Liverpool Road Chester United Kingdom CH2 1UL +44 01244 365000

# Additional identifiers

**EudraCT/CTIS** number

IRAS number

ClinicalTrials.gov number

**Secondary identifying numbers** N0072186040

# Study information

### Scientific Title

# Study objectives

- 1. Is bupivacaine better than placebo in reducing the post-operative pain?
- 2. Does bupivacaine used as supplement reduce the opiate dosage?

# Ethics approval required

Old ethics approval format

# Ethics approval(s)

Not provided at time of registration

# Study design

Randomised double blind controlled trial

# Primary study design

Interventional

# Secondary study design

Randomised controlled trial

# Study setting(s)

Hospital

# Study type(s)

Treatment

### Participant information sheet

# Health condition(s) or problem(s) studied

Signs and Symptoms: Pain

#### **Interventions**

A blinded member of the team will assess the pain at 2,4 and hours after the operation. Length of hospital stay is also recorded.

The data analysis will take place at Countess of Chester Hospital (COCH). The research team at COCH will analyse the data using SPSS software.

# Intervention Type

Drug

#### Phase

**Not Specified** 

# Drug/device/biological/vaccine name(s)

Bupivacaine

### Primary outcome measure

Is bupivacaine superior than placebo in pain control following breast augmentation?

# Secondary outcome measures

- 1. Does bupivacaine reduce opiate requirement?
- 2. Does it reduce hospital stay?
- 3. Does it reduce the incidence of opiate related adverse effects?

# Overall study start date

01/08/2006

# Completion date

01/08/2009

# **Eligibility**

# Key inclusion criteria

- 1. Patients undergoing breast augmentation
- 2. For correction of congenital anomaly or asymmetry. This group is usually young population, so usually not associated with co-morbid conditions which might complicate result.

# Participant type(s)

**Patient** 

### Age group

Adult

### Sex

Female

# Target number of participants

Total 50; 25 in each group.

# Key exclusion criteria

- 1. Patients undergoing augmentation following mastectomy or as part of other reconstruction
- 2. Patients with multiple co-morbidity

### Date of first enrolment

01/08/2006

### Date of final enrolment

01/08/2009

# Locations

### Countries of recruitment

England

**United Kingdom** 

# Study participating centre

# Countess of Chester Hospital NHS Foundation Trust

Chester United Kingdom CH2 1UL

# Sponsor information

# Organisation

Record Provided by the NHSTCT Register - 2007 Update - Department of Health

## Sponsor details

The Department of Health, Richmond House, 79 Whitehall London United Kingdom SW1A 2NL +44 (0)20 7307 2622 dhmail@doh.gsi.org.uk

# Sponsor type

Government

### Website

http://www.dh.gov.uk/Home/fs/en

# Funder(s)

# Funder type

Government

### **Funder Name**

Countess of Chester NHS Foundation Trust

# **Funder Name**

NHS R&D Support Funding

# **Results and Publications**

# Publication and dissemination plan

Not provided at time of registration

Intention to publish date
Individual participant data (IPD) sharing plan

**IPD sharing plan summary**Not provided at time of registration