Adjuvant cyclophosphamide, methotrexate and 5-Fluorouracil (CMF) versus 5-Fluorouracil, epirubicin and cyclosphosphamide (FEC) in node negative poor-risk primary breast cancer

Submission date	Recruitment status	[X] Prospectively registered
19/08/2002	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
19/08/2002	Completed	Results
Last Edited	Condition category	Individual participant data
06/11/2012	Cancer	Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr - -

Contact details

UKCCCR Register Co-ordinator MRC Clinical Trials Unit 222 Euston Road London United Kingdom NW1 2DA

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

Study information

Scientific Title

Study objectives

Not provided at time of registration

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Not Specified

Participant information sheet

Health condition(s) or problem(s) studied

Breast cancer

Interventions

Patients randomised to receive chemotherapy will receive CMF or FEC no sooner than 2 weeks and no later than 4 weeks following primary surgery.

Patients will receive either:

- 1. CMF Regimen: Chemotherapy, CMF (cyclophosphamide, methotrexate, 5-fluorouracil) repeated every 4 weeks for six cycles.
- 2. FEC Regimen: Chemotherapy, FEC (5-fluouracil, 4-epidoxorubicin, cyclophosphamide) repeated every 4 weeks for six cycles.

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Not provided at time of registration

Secondary outcome measures

Not provided at time of registration

Overall study start date

01/01/2003

Completion date

01/01/2004

Eligibility

Key inclusion criteria

- 1. Aged less than 65 years
- 2. Histologically proven invasive adenocarcinoma axillary nodes must be histologically negative
- 3. Tumour is T1c to T3
- 4. Tumour shows one of the following features of poor prognosis:
- 4.1. Oestrogen receptor less than or equal to 10 fmol/mg cytosol protein
- 4.2. Grade III
- 4.3. Vessel invasive
- 4.4. High labelling index of S phase
- 4.5. Aneuploidy
- 4.6. Tumour size at least 2 cm
- 5. Adequate renal, hepatic and haematological function
- 6. No evidence of metastatic disease
- 7. No evidence of inflammatory carcinoma
- 8. Patients with bilateral malignancy or with a mass in the opposite breast, unless there is biopsy proof that it is not a malignancy, are excluded
- 9. Patients with findings that relate them to a category of more advanced disease are not eligible
- 10. Patients with clinically positive nodes in the axilla opposite the affected breast, or with palpable supraclavicular or infraclavicular nodes are considered ineligible unless there is a biopsy evidence that these are not involved with the tumour
- 11. No previous or concomitant malignancy, except squamous or basal cell carcinoma which has been effectively treated and carcinoma in situ of the cervix which has been treated operatively
- 12. No prior therapy for the present breast cancer including radiotherapy, chemotherapy, immunotherapy and/or hormonal therapy
- 13. No non-malignant systemic disease which would preclude their being subjected to any of the treatment options or prevent prolonged follow up
- 14. No active or previous cardiac disease that would preclude the use of 4-epidoxorubicin
- 12. No psychiatric or addictive disorders which should preclude obtaining informed consent

Participant type(s)

Patient

Age group

Adult

Sex

Female

Target number of participants

Not provided at time of registration

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/01/2003

Date of final enrolment

01/01/2004

Locations

Countries of recruitment

England

United Kingdom

Study participating centre UKCCCR Register Co-ordinator

London United Kingdom NW1 2DA

Sponsor information

Organisation

Pharmacia Ltd & Upjohn (UK)

Sponsor details

Davy Avenue Milton Keynes United Kingdom MK5 8PH +44 (0)1908 661101 info@adreco.co.uk

Sponsor type

Industry

Website

http://www.pharmacia.com

ROR

https://ror.org/04x4v8p40

Funder(s)

Funder type

Industry

Funder Name

Pharmacia and Upjohn (UK)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration