

Effects of intra-articular administration of Hymovis on the expression and activity of matrix metalloproteinases in synovial fluids of patients affected by osteoarthritis of the knee

Submission date 01/08/2014	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 15/09/2014	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 15/10/2020	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims:

Osteoarthritis (OA) is a degenerative disease of the joint. Over time, the cartilage covering the end of the bones at the joints become rough and thin, and in some cases wearing out completely, so that the bones in the joint grind against each other. Symptoms include painful and stiff joints and a limited range of movement. It can have a serious impact on the life of the individual with the condition as well as on public health services. Matrix metalloproteinase 3 (MMP3) is an enzyme known to be involved in the breakdown of cartilage and it can be found in the joint fluid (synovial fluid) of patients suffering from OA of the knee. Here, we want to see if injecting the knee with a synovial fluid replacement, Hymovis (a hyaluronic acid), can reduce the amount of MMP3 present and relieve pain. Hymovis has already been shown to work in the laboratory. We also hope to see whether Hymovis can also have an effect on other proteins that may be involved in OA.

Who can participate

Adults aged at least 46 with OA of the knee.

What does the study involve?

Participants are randomly allocated into one of two groups. Those in group 1 are treated with Hymovis. Those in group 2 are treated with another hyaluronic acid called Synvisc. Both groups are given 3 injections in all. The synovial fluid in the knee is collected and analysed before the first and second injection to check for MMPs and other molecules that suggest that there is inflammation. All patients are followed up for 6 months. At the end of the study, we assess the performance of the two treatments.

What are the possible benefits and risks of participating?

There are no immediate direct benefit for those taking part in this study, but there should be benefits for future patients. It is possible that the treatment will relieve knee OA pain. The main risks are pain, swelling or infections.

Where is the study run from?

This study has been set up by Fidia farmaceutici S.p.A. and it will be carried out by the Medicine Department of the University of Perugia (Italy)

When is the study starting and how long it is expected to run for?

September 2014 to September 2015

Who is funding the study?

Fidia farmaceutici S.p.A. (Italy)

Who is the main contact?

Cinzia Santin

csantin@fidiapharma.it

Contact information

Type(s)

Scientific

Contact name

Prof Roberto Gerli

Contact details

Università degli Studi di Perugia

Dep. Medicina

Via Enrico Dal Pozzo

Perugia

Italy

06122

-

roberto.gerli@unipg.it

Additional identifiers

Protocol serial number

EQC5-14-01

Study information

Scientific Title

A post-marketing, pilot study to assess the role of intra-articular administration on the expression and activation of metalloproteinases in synovial fluid.

Study objectives

It is hypothesized that Hymovis has the capacity to decrease the expression and activation of MMPs in synovial fluid, the purpose of the study is to verify this activity comparing it to that of another product for the same indication

Ethics approval required

Old ethics approval format

Ethics approval(s)

Umbria Ethic Committee (CEAP), 21/05/2014, ref. CEAS N 2333/14

Study design

Pilot study, randomized open label post marketing

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Knee Osteoarthritis

Interventions

Patients are randomly allocated into one of two groups. One group is treated with Hymovis hyaluronic acid (HA) and the other group, Synvisc HA.

All patients will undergo one cycle of intra-articular injections with HA, synovial fluid will be collected at baseline and before the second injection. 6 visits are scheduled:

Screening visit: enrolment

Visit 1: Start of therapy (first injection)

Visit 2: During study treatment (second injection)

Visit 3: During study treatment (third injection)

Visit 4: First FU (3 months after baseline)

Visit 5: Second FU (6 months after baseline)

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

1. Synovial fluid analysis will be carried out at baseline and at visit 1
2. MMP3 activity will be evaluated via ELISA test

Key secondary outcome(s)

The following variables will be evaluated at baseline and visit 1:

1. Macroscopic analysis of SF
2. Microscopic analysis of SF
3. Expression of MMP2 and MMP13 (ELISA test)
4. Expression of inflammatory cytokines (Flowcytomix analysis)

At baseline and FU visits the following will be considered:

1. VAS
2. WOMAC
3. Physicians global assessment

For all the duration of the study safety and tolerability will be assessed by recording all AE

Completion date

03/09/2015

Eligibility

Key inclusion criteria

1. Ambulant patient > 45 years of age in good health conditions
2. Patient of any race, both outpatients or hospitalized, affected by OA diagnosed both clinically and radiologically according to the American College of Rheumatology criteria:
 - 2.1. Pain
 - 2.2. Presence of radiographic osteophytes
 - 2.3. At least one of the following items:
 - 2.3.1. Age > 50
 - 2.3.2. Morning stiffness < 30 minutes of duration
 - 2.3.3. Crepitus on motion
3. Kellgren-Lawrence grade II-IV confirmed by X-ray (performed within 6 months preceding the inclusion in the study)
4. Patient diagnosed with idiopathic OA of the knee since at least one year
5. Patient presenting synovial effusion of the knee joint echographically confirmed
6. Patient who discontinued previous oral/topical NSAIDs therapy since at least one week before the first intra-articular injection
7. Male or fertile, non pregnant or breast feeding female who consents to use a reliable birth control method for the whole treatment duration
8. Patient without conditions or psychological impairment who could interfere with compliance to the protocol and the study conclusion in the investigator's opinion
9. Patient able to understand study obligations and who provided a written informed consent in compliance with the Helsinki declaration (version October 2013)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Concomitant chronic inflammatory diseases (e.g. rheumatoid arthritis or psoriasis)
2. Painful knee condition other than OA
3. Secondary knee OA or relevant knee injuries in the target knee within 6 months before the inclusion
4. Known or suspected allergic reactions to hyaluronic preparations or to other components of the devices under study
5. Presence of infections and/or skin diseases in the area of injection site; e.g. psoriasis
6. Known coagulopathy
7. Therapy with one of the devices under study within 6 months before the inclusion

8. Systemic anti-inflammatory therapy at a fixed dose within 1 month before the inclusion
9. In therapy with i.a. HA or i.a. corticosteroids within 6 months before the inclusion
10. In therapy with biphosphonates, and/or oral pharmaceutical products containing glucosamine and/or chondroitin sulphate and/or diacerein within 1 month before inclusion
11. Participation in any other study involving investigational or marketed products simultaneously or within one month prior to study entry
12. Previous surgery in the target knee within 9 months prior to inclusion, or planned surgery throughout the duration of the study
13. Presence of pathologies or conditions that might interfere with subject compliance /cooperation during the study, in the judgment of the investigator
14. History of alcoholism, treatment abuse, psychological or other emotional problems that, in the judgment of the investigator, could invalidate informed consent or limit the subjects compliance with protocol requirements
15. Females who are pregnant, lactating or not in menopause and are not using recognized contraceptive measures as indicated by investigators for the duration of the study

Date of first enrolment

03/09/2014

Date of final enrolment

03/09/2015

Locations

Countries of recruitment

Italy

Study participating centre

Università degli Studi di Perugia

Perugia

Italy

06122

Sponsor information

Organisation

Fidia Pharmaceuticals S.p.a (Fidia Farmaceutici S.p.a) (Italy)

ROR

<https://ror.org/00stv9r10>

Funder(s)

Funder type

University/education

Funder Name

University of Perugia (Italy)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration