

Haemodynamic stability during anaesthesia induction with propofol the impact of phenylephrine

Submission date 07/06/2014	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 11/07/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 29/05/2020	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Propofol is a drug that is widely used to anaesthetize a person before they have an operation. One of the effects of propofol is a drop in blood pressure. Other drugs, called vasopressors, have to then be given to the patient in order to raise the blood pressure to a more normal level. In this study, we will assess how well a continually administered dose of a vasopressor (phenylephrine), given when the patient is being put under a general anaesthetic with propofol, is at maintaining a more stable blood pressure.

Who can participate?

Adult patients with abdominal cancers about to have high-risk abdominal surgery.

What does the study involve?

Patients are randomly allocated into one of two groups. One group are given a continuous dose of phenylephrine while being put under a general anaesthetic with propofol. The other group are given a saline solution instead. The blood pressure of both groups of patients are measured during the start (induction) of anaesthesia and compared.

What are the possible benefits and risks of participating?

All patients are closely monitored, including continually measuring both their blood pressure and heart rate, throughout the procedure. Any possible increase or decrease in blood pressure compared to the norm will be immediately identified and treated. Therefore patients are not exposed to any additional risk.

Where is the study run from?

The Maribor University Clinical Centre (Slovenia)

When is study starting and how long is it expected to run for?

November 2013 to January 2015

Who is funding the study?
The Maribor University Clinical Centre (Slovenia)

Who is the main contact?
Professor Mirt Kamenik
mirt.kamenik@gmail.com

Contact information

Type(s)
Scientific

Contact name
Prof Mirt Kamenik

Contact details
V zatiju 20
Bresternica
Slovenia
2354

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
IRP-2013/02-06

Study information

Scientific Title
The impact of phenylephrine infusion on haemodynamic stability during bispectral index (BIS) guided anaesthesia induction with propofol a double-blind randomised controlled trial

Acronym
BIS PROPphen

Study objectives
We will compare the induction of general anaesthesia with propofol and propofol combined with phenylephrine both titrated to the same bispectral index value. We expect less hypotension and cardiac output decrease during and after the induction of anaesthesia with the use of the combination of propofol and phenylephrine. We are also interested which impact does phenylephrine have on other haemodynamic parameters during and after induction.

Ethics approval required
Old ethics approval format

Ethics approval(s)

National Medical Ethics Committee;10/10/2013; ref. 229/09/13

Study design

Single-centre double-blinded prospective randomized controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Other

Study type(s)

Other

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Cardiovascular disease and abdominal cancer surgery

Interventions

We will study the haemodynamic effects of intravenous anaesthesia induction with propofol titrated to the appropriate anaesthesia depth monitored with spectral index monitor. During induction the patients will receive a continuous infusion of either phenylephrine or saline in a double blind randomized fashion. Hypotension (<55 mmHg) will be managed with additional phenylephrine. Hypertension (>100 mmHg) will be managed with the stop of infusion, additional, fentanyl and nitroglycerine. Tachycardia (>90/min) will be managed with fentanyl and esmolol. Bradycardia (<40/min) will be managed with atropine.

Intervention Type

Procedure/Surgery

Phase

Not Applicable

Primary outcome measure

1. Mean arterial pressure
2. Cardiac output
3. Systemic vascular resistance

All outcomes will be measured by the LIDCCORapid monitor for measuring cardiac output. The new generation of LIDCCORapid monitor enables also the measurement of the depth of anaesthesia with the BIS monitor. The measurements will be done at baseline and for 20 minutes during the induction of anaesthesia. The LIDCCORapid monitor enables automatic online collection of the data which will be averaged over one minute intervals for the period of

induction. The averaged data for both groups of the results will be presented for baseline values, the values immediately before laryngoscopy and intubation, 2 minutes after intubation and then in 3 minute intervals till the end of measurements.

Secondary outcome measures

1. Heart rate
2. Stroke volume
3. Bispectral index
4. Oxygen saturation
5. Dose of propofol
6. Time from start of anaesthesia induction to laryngoscopy and intubation
7. Dose of rescue drugs needed

The other secondary outcome measures: the dose of propofol during induction of anaesthesia, the time to laryngoscopy and intubation, the frequency of rescue medication used and the dose of rescue medication given is registered during the study for each study subject and the data are collected on our protocol sheet.

Overall study start date

01/11/2013

Completion date

01/01/2015

Eligibility

Key inclusion criteria

ASA II-III patients scheduled for major abdominal surgery (large bowel resection, stomach resection, liver resection, Whipple resection) and major urologic surgery (bladder resection, prostatic cancer).

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

50

Total final enrolment

40

Key exclusion criteria

1. Alcohol-abuse
2. Drug abuse
3. Chronic use of benzodiazepines, opioids or other psychotropic substances
4. Body mass index > 30

- 5. Anticipated difficult intubation (Mallampati 3 and 4)
- 6. Kidney disease (serum creatinine > 120 mmol/l)
- 7. Manifest liver disease
- 8. Alzheimer disease
- 9. Epilepsy

Date of first enrolment

01/11/2013

Date of final enrolment

01/01/2015

Locations

Countries of recruitment

Slovenia

Study participating centre

V zatiju 20

Bresternica

Slovenia

2354

Sponsor information

Organisation

Maribor University Clinical Centre (Slovenia)

Sponsor details

Ljubljanska 5

Maribor

Slovenia

2000

Sponsor type

Hospital/treatment centre

ROR

<https://ror.org/02rjj7s91>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Maribor University Clinical Centre (Slovenia)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/03/2018	29/05/2020	Yes	No