Impact of graded nursing interventions based on

Submission date 18/04/2025	Recruitment status No longer recruiting	Prospectively registeredProtocol
Registration date 25/04/2025	Overall study status Completed	Statistical analysis plan
		Results
Last Edited 23/04/2025	Condition category Cancer	Individual participant data
		[X] Record updated in last year
Plain English summary of protocol Background and study aims Liver cancer is one of the malignancies with high incidence and mortality rates worldwide. This study aims to explore the impact of quantitative graded nursing interventions on psychological stress and postoperative recovery in patients undergoing primary liver cancer resection by constructing a grading assessment system centered on Child-Pugh scores and HAMA/HAMD scales, combined with dynamic nursing resource allocation. Who can participate? Patients preparing for liver cancer surgery. What does the study involve? Health education and nursing before and after liver cancer surgery.		
What are the possible benefits and risks of participating? Helps liver cancer patients recover after surgery. No potential risks.		
Where is the study run from? The study was conducted at the First Affiliated Hospital of Ningbo University (China)		
When is the study starting and how long is it expected to run for? January 2023 to February 2024		
Who is funding the study? Provincial Medical Project (No.2025KY249) (China)		

Contact information

Who is the main contact?

Type(s)

Public, Scientific, Principal investigator

Dr Qinhong Xu, xuqinhong_xqh03@126.com

Contact name

Dr Qinhong Xu

ORCID ID

https://orcid.org/0009-0008-8726-2439

Contact details

59 Liu Ting Street, Haishu District Ningbo China 315000 +86 0574 87085014 xuqinhong_xqh03@126.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

The impact of graded nursing interventions based on quantitative risk assessment on psychologica

Study objectives

Graded nursing interventions based on quantitative assessment may reduce postoperative psychological stress levels and negative emotions in patients undergoing primary liver cancer surgery, promoting postoperative recovery

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 13/07/2023, Ethics Committee of The First Affiliated Hospital of Ningbo University (59 Liu Ting Street, Haishu District, Ningbo, 315000, China; +86 0574 87085233; sychenjy@163.com), ref: 2023Y-095RS

Study design

Single-center interventional double-blind randomized controlled trial

Primary study design

Interventional

Study type(s)

Other

Health condition(s) or problem(s) studied

Nursing care for patients undergoing resection of primary hepatic cancer

Interventions

Random number sequences are generated using SPSS 26.0, stratified by Child-Pugh score (A/B grade), with a block size of 4 for each stratum. Sealed opaque envelopes are used to store allocation results, managed by an independent statistician. After enrollment, research nurses assign patients to groups according to the envelope instructions, dividing them into an observation group and a control group, each with 40 patients.

The control group receive routine nursing care, which included:

Preoperative health education: Nursing staff provided patients with knowledge about liver cancer through oral communication; Postoperative pain management (VAS score ≤ 3): Close monitoring of patients' pain, bowel movements, and drainage tube conditions after surgery; Early ambulation (within 24 hours postoperatively): Guidance on early ambulation to actively prevent complications. A targeted assessment of the condition of enrolled liver cancer patients is conducted. Treatment risk was categorized based on the comprehensive score: low risk medium risk and high risk.

Based on the quantitative assessment results, clinical nursing staff from various departments were allocated appropriately, and stratified nursing measures were implemented. Low-risk patients: Assigned N0–N1 nursing staff to provide routine ERAS care + psychological support (10 minutes of mindfulness training daily). Medium-risk patients: Assigned N2 or higher nursing staff to implement low-risk measures + personalized exercise plans (bed activities starting 6 hours postoperatively). High-risk patients: Assigned N3 or higher nursing staff, team leaders, or specialist nurses to implement medium-risk measures + multidisciplinary team interventions (joint daily rounds by psychiatric nurses and nutritionists).

Intervention Type

Behavioural

Primary outcome(s)

- 1. Psychological stress response is measured using Psychological Stress Response Questionnaire (SRQ) at one day before surgery, three days after surgery, and at discharge.
- 2. Depressive state is measured using Hamilton Depression Scale (HAMD) at one day before surgery, three days after surgery, and at discharge.
- 3. Anxiety level is measured using Hamilton Anxiety Scale (HAMA) at one day before surgery, three days after surgery, and at discharge.

Key secondary outcome(s))

Measured using patient records:

- 1. Recorded the time to tolerate semi-liquid diet (time from surgery to first intake of ≥200 mL semi-liquid food)
- 2. Time to first out-of-bed activity (time from surgery to independent standing and walking ≥5 meters)
- 3. Time to first postoperative anal exhaust, and length of hospital stay

Completion date

06/02/2024

Eligibility

Key inclusion criteria

- 1. Meeting the surgical indications outlined in the "Guidelines for the Diagnosis and Treatment of Primary Liver Cancer (2022 Edition)" (single tumor diameter ≤ 5 cm, no portal vein trunk tumor thrombus, Child-Pugh A/B grade)
- 2. Age 18-75 years
- 3. Signed informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

75 years

Sex

Αll

Total final enrolment

80

Key exclusion criteria

- 1. Severe cardiovascular disease (NYHA III-IV grade)
- 2. Immune system disease or coagulation dysfunction (INR > 1.5, PLT < $50 \times 10^9/L$)
- 3. Malignant tumors in other locations
- 4. History of cognitive impairment or mental illness.

Date of first enrolment

06/01/2023

Date of final enrolment

31/12/2023

Locations

Countries of recruitment

China

Study participating centre

The First Affiliated Hospital of Ningbo University

59 Liu Ting Street, Haishu District Ningbo China 315000

Sponsor information

Organisation

The First Affiliated Hospital of Ningbo University

Funder(s)

Funder type

Government

Funder Name

Provincial Medical Project

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available due to privacy and ethical restrictions.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output typeDetailsDate createdDate addedPeer reviewed?Patient-facing?Participant information sheetParticipant information sheet11/11/202511/11/2025NoYes