

# Molecular basis of depression disorder and influence of omega-3 fatty acid on clinical symptoms and biomarkers

<b>Submission date</b> 10/10/2016	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 19/10/2016	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 28/02/2024	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Mood disorders in children and adolescents are a serious problem worldwide and are becoming more common in younger children. In recent years, research has been focused on the adjuvant therapy of depression (extra treatment to enhance the effects of the main treatment) to reduce the consumption of antidepressants, prolong remission (disease-free periods) and improve the outlook for these patients. The use of Omega-3 fatty acids as an adjuvant therapy is promising, with beneficial effects in the prevention and treatment of depression. This study aims to assess the effectiveness of omega-3 fatty acids-rich fish oil in the treatment of depression symptoms in children and adolescents treated for depression as well as to find out what effects this has on the body by looking at biochemical markers (substances).

### Who can participate?

Children and adolescents aged 10 -18 years suffering from depression who are registered at the Department of Child Psychiatry of Comenius University and Child University Hospital in Bratislava.

### What does the study involve?

Patients are randomly allocated to one of two groups. Those in the first group take 20 mL of omega-3 fish oil emulsion every day for 12 weeks in addition to their usual treatment, followed by 4 weeks of not taking the oil. Those in the second group take 20 mL of omega-6 sunflower oil emulsion every day for 12 weeks in addition to their usual treatment, followed by 4 weeks of not taking the oil. At the start of the study, every two weeks during the 12-week treatment period, and then at 16 weeks, participants have their symptoms of depression assessed and have blood and saliva samples taken to measure for chemical indicators of the effects the supplements are having on the body.

### What are the possible benefits and risks of participating?

Participants who receive the omega-3 oil may benefit from a reduction of their depressive symptoms. There are no known risks involved with participating.

Where is the study run from?

1. Comenius University (Slovakia)
2. Comenius University and Child University Hospital (Slovakia)

Who is funding the study?

Slovak Research and Development Agency (Slovakia)

When is the study starting and how long is it expected to run for?

March 2013 to June 2020

Who is the main contact?

Professor Zdenka Durackova

zdenka.durackova@fmed.uniba.sk

## Contact information

**Type(s)**

Scientific

**Contact name**

Prof Zdeňka Ďuračková

**ORCID ID**

<http://orcid.org/0000-0001-6956-7038>

**Contact details**

Institute of Medical Chemistry

Biochemistry and Clinical Biochemistry

Faculty of Medicine

Comenius University

Bratislava

Slovakia

813 72

+421 1 59357 564

zdenka.durackova@fmed.uniba.sk

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

APVV-15-0063

## Study information

**Scientific Title**

Molecular bases of depressive disorders in children and adolescents, effect of omega-3 fatty acids, inflammation and oxidative stress

## **Acronym**

Depoxin

## **Study objectives**

Current study hypothesis as of 05/12/2023:

The current study aims to compare the efficacy of 12 weeks of administration of omega-3 fatty acids present in fish oil emulsion with a control oil emulsion (omega-6 fatty acids rich sunflower oil emulsion) alongside standard antidepressant treatment on depressive symptoms in children and adolescents (10 to 18 years old).

The study will investigate:

1. Clinical parameters: CDI score
2. Biochemical parameters: Lipid profile, fatty acid composition in serum LDL- and HDL-subfractions of lipoproteins, markers of oxidative stress and inflammation in blood, steroidal hormones in saliva, metabolites of kynurenine and serotonin pathway
3. Biophysical parameters: Fluidity of erythrocyte membranes
4. The length of telomeres in the buccal cells

Previous study hypothesis:

The aim of the current study is to compare the efficacy of 12 weeks administration of omega-3 fatty acids present in fish oil emulsion with a control oil emulsion (omega-6 fatty acids rich sunflower oil emulsion) alongside standard antidepressant treatment on depressive symptoms in children and adolescents (8 to 18 years old).

The study will investigate:

1. Clinical parameters: CDI score
2. Biochemical parameters: Lipid profile, fatty acid composition in serum LDL- and HDL-subfractions of lipoproteins, markers of oxidative stress and inflammation in blood, steroidal hormones in saliva
3. Biophysical parameters: Fluidity of erythrocyte membranes
4. The length of telomeres in the buccal cells

## **Ethics approval required**

Ethics approval required

## **Ethics approval(s)**

Approved 20/03/2013, Ethics Committee of the National Institute of Children's Diseases and the Faculty of Medicine, Comenius University Bratislava (Limbova 1, Bratislava, 833 40, Slovakia; +421 2 59371209, +421 2 59371927; detska.klinika@nudch.eu), ref: None available

## **Study design**

Single-centre randomized double-blind comparator controlled interventional study

## **Primary study design**

Interventional

## **Secondary study design**

Randomised controlled trial

## Study setting(s)

Home

## Study type(s)

Treatment

## Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet in Slovakian

## Health condition(s) or problem(s) studied

Depressive disorder (F32) or mixed anxiety and depressive disorder (F41.2)

## Interventions

Patients are allocated in a 1:1 ratio to the two arms (Om3 and Om6) according to a computer-generated random sequence using block randomisation with a block-size of four. The randomisation was performed by the independent statistician. Patients are enrolled and assigned sequentially to adjuvant interventions by the physician. The allocation sequence is not available to any member of the research team until the databases had been completed and locked.

Intervention group: Participants receive 20ml of an omega-3 FA rich fish oil emulsion (Om3), which provides 2400 mg of total omega-3 FA; 1000 mg EPA and 750 mg DHA, EPA:DHA ratio = 1.33:1, for 12 weeks in addition to their standard antidepressant therapy, followed by a 4 week wash out period.

Control group: Participants receive an identically looking comparator omega-6 sunflower oil emulsion containing min 2467 mg of omega-6 linoleic acid provided by Cultech Ltd, Port Talbot (UK) for 12 weeks in addition to their standard antidepressant therapy, followed by a 4 week wash out period.

Clinical examinations take place at the beginning of the project (week 0), and every 2 weeks for 3 months (weeks 2, 4, 6, 8, 10, 12) and after following 4 week wash-out period, at week 16. Ratings are made using the self rated scale Children's Depression Inventory (CDI). Biological samples (blood, urine, saliva) are taken at the week 0, 6, 12 and 16.

## Intervention Type

Supplement

## Primary outcome measure

Clinical symptoms of depression are determined using Children's Depression Inventory score (CDI) at baseline, 2, 4, 6, 8, 10, 12 and 16 weeks.

## Secondary outcome measures

1. Basic biochemical marker levels (glucose, uric acid, lipid profile (total cholesterol, LDL-CH, HDL-CH, TAG), creatinine, CRP, liver enzymes ALT, AST) are measured by testing blood samples taken at baseline, at the week 6 and 12 of intervention and after wash-out period, at the week 16
2. Markers of oxidative stress (antioxidant enzymes SOD, GPx, CAT in Er, PON1 in serum, markers of lipid damage - lipoperoxides, 8-IsoP, protein damage - AOPP, serum protein carbonyls, nitrotyrosine, DNA damage in lymphocytes by comet assay, total antioxidant status-TEAC) are measured by testing blood samples taken at baseline, at the week 6 and 12 of intervention and

after wash-out period, at the week 16

3. Markers of inflammation (CRP, calprotectin, cytokines and eicosanoids, thromboxanes) are measured by testing blood samples taken at baseline, at the week 6 and 12 of intervention and after wash-out period, at the week 16

4. Hormones in saliva (cortisol, aldosterone) are measured by testing saliva samples taken at baseline, at the week 6 and 12 of intervention and after wash-out period, at the week 16

5. Erythrocyte (Er) membrane fluidity is measured by spectrofluorimeter at baseline, at the week 6 and 12 of intervention and after wash-out period, at the week 16

6. Composition and content of serum fatty acids (FA) are measured by gas chromatography at baseline, at the week 6, 12 of intervention and after wash-out period, at the week 16

8. Length of DNA telomers in buccal cells is measured by qPCR at baseline and at 12 week intervention

**Overall study start date**

01/03/2013

**Completion date**

30/06/2020

## **Eligibility**

**Key inclusion criteria**

1. Children and adolescents who met diagnostic criteria for depressive disorder or mixed anxiety and depressive disorder according to ICD 10
2. No signs of chronic somatic disease
3. Normal eating habits
4. Parents/guardians are willing to provide a signed informed consent
5. Children and adolescents are willing to provide blood, urine and saliva samples

**Participant type(s)**

Patient

**Age group**

Child

**Lower age limit**

10 Years

**Upper age limit**

18 Years

**Sex**

Both

**Target number of participants**

60 = 30 in omega-3 group and 30 in omega-6 group

**Total final enrolment**

58

**Key exclusion criteria**

1. Chronic somatic diseases (endocrine, metabolic, autoimmune)
2. Dietary restrictions (vegetarians, lactose intolerance, celiac disease)
3. Psychotic disorders
4. Eating disorders
5. Addiction to psychoactive compounds
6. Personality disorders
7. Organic mental disorders
8. Pervasive developmental disorders

**Date of first enrolment**

01/03/2014

**Date of final enrolment**

30/06/2019

**Locations****Countries of recruitment**

Slovakia

**Study participating centre****Comenius University**

Department of Paediatric Psychiatry, Faculty of Medicine, The National Institute of Children's Diseases  
Limbová 1  
Bratislava  
Slovakia  
833 40

**Study participating centre****Comenius University**

Institute of Medical Chemistry, Biochemistry and Clinical Biochemistry  
Faculty of Medicine  
Sasinkova 2  
Bratislava  
Slovakia  
813 72

**Sponsor information****Organisation**

Comenius University

**Sponsor details**

Faculty of Medicine  
Špitálska 24  
Bratislava  
Slovakia  
813 72  
+421 2 593 57 466  
dekan.sekretariat@fmed.uniba.sk

**Sponsor type**

University/education

**Website**

<https://www.fmed.uniba.sk/>

**ROR**

<https://ror.org/0587ef340>

**Funder(s)****Funder type**

Research organisation

**Funder Name**

Slovak Research and Development Agency (Agentúra na Podporu Výskumu a Vývoja)

**Results and Publications****Publication and dissemination plan**

Planned publication in a high-impact peer reviewed journals of papers discussing:

1. Clinical symptoms – results of pilot study - 2017
2. Markers of oxidative stress – pilot results -2017
3. Markers of inflammation – pilot results 2018
4. Fluidity of membranes and fatty acid composition - 2019
5. Telomere length – 2019
6. Mutual relation between clinical symptoms and markers of oxidative stress and inflammation - 2020

**Intention to publish date**

30/06/2021

**Individual participant data (IPD) sharing plan**

The datasets generated during and/or analysed during the current study will be available upon request from [zdenka.durackova@fmed.uniba.sk](mailto:zdenka.durackova@fmed.uniba.sk) and [jana.trebaticka@fmed.uniba.sk](mailto:jana.trebaticka@fmed.uniba.sk)

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	Pilot study results	05/07/2017		Yes	No
<a href="#">Results article</a>	Association of depression symptoms with levels of omega-3 and omega-6 fatty acids in serum	01/05/2020	18/03/2020	Yes	No
<a href="#">Results article</a>	Lipid profile results	08/10/2020	22/10/2020	Yes	No
<a href="#">Results article</a>	results for levels of thromboxane, brain-derived neurotrophic factor, homocysteine and vitamin D	27/03/2021	06/04/2021	Yes	No
<a href="#">Results article</a>	Changes in stress hormones and markers of oxidative stress	10/08/2022	30/08/2022	Yes	No
<a href="#">Results article</a>	The kynurenine and serotonin pathway, neopterin and biopterin in depressed children and adolescents: an impact of omega-3 fatty acids, and association with markers related to depressive disorder. A randomized, blinded, prospective study	13/02/2024	28/02/2024	Yes	No