

A Phase II, open-label study to evaluate the safety, tolerability, and pharmacokinetic profile of Proxinium™ in patients with recurrent squamous cell carcinoma of the head and neck who have received at least one prior anti-cancer treatment regimen for recurrent disease

Submission date 06/04/2006	Recruitment status Stopped	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 17/08/2006	Overall study status Stopped	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 28/01/2019	Condition category Cancer	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Jerald Kahan

Contact details
4009 Banister Lane
Austin
United States of America
78704
+1 512 685 5779
jerald.kahan@austin.ppd.com

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

NCT00272181

Secondary identifying numbers

VB4-845-01-IIA

Study information

Scientific Title

A Phase II, open-label study to evaluate the safety, tolerability, and pharmacokinetic profile of Proxinium™ in patients with recurrent squamous cell carcinoma of the head and neck who have received at least one prior anti-cancer treatment regimen for recurrent disease

Study objectives

To evaluate the safety, tolerability and pharmacokinetic profile of Proxinium™ in patients with recurrent Squamous Cell Carcinoma of the Head and Neck (SCCHN).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved by Institutional Review Boards at various Universities, Hospitals and Clinics in North America, also approved by the Food and Drug Administration on 17/11/2005 (reference number: 12610).

Study design

Multicenter open-label safety study

Primary study design

Interventional

Secondary study design

Multi-centre

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet**Health condition(s) or problem(s) studied**

Squamous Cell Carcinoma of the Head and Neck

Interventions

Proxinium™ injected intratumorally weekly.

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

Proxinium₂

Primary outcome measure

Determine safety, tolerability and pharmacokinetic profile of Proxinium™ in SCCHN patients.

Secondary outcome measures

1. Tumour response rates
2. Time to progression
3. Overall Survival
4. Progression free survival associated with intratumoral injection of Proxinium™

Overall study start date

16/12/2005

Completion date

30/11/2006

Reason abandoned (if study stopped)

Participant recruitment issue

Eligibility

Key inclusion criteria

1. The patient must have histologically confirmed SCCHN
2. The patient must have immunohistochemically confirmed Epithelial Cell Adhesion Molecule (Ep-CAM) positive SCCHN
3. The patient must have received therapy for their primary disease (i.e., SCCHN) consisting of radiotherapy with or without surgery and with or without chemotherapy
4. Patients must have progressed on or after receiving at least one prior anti-cancer treatment regimen containing one or more anti-cancer agents (e.g., chemotherapy, biologic therapy, or photodynamic therapy) for their recurrent disease
5. The patient must have fully recovered or reached a stable state of symptomatology from any previous treatment-related toxicity
6. The patient must have at least one accessible target tumor without direct carotid artery involvement (ie, a distance of less than 5 mm between a tumor and carotid) and must be likely to retain study drug

Participant type(s)

Patient

Age group

Not Specified

Sex

Both

Target number of participants

18

Key exclusion criteria

1. The patient has known brain tumor or brain metastases
2. The patient has nasopharyngeal SCCHN
3. The patient has concurrent or documented history of any one of the following:
 - a. Human Immunodeficiency Virus (HIV)
 - b. Hepatitis C virus
 - c. Hepatitis B surface antigen
4. The patient has uncontrolled bleeding from any target tumor(s) that are being considered for Proxinium™ treatment
5. The patient has a history of tumor hemorrhage that has required medical intervention (other than direct compression)
6. The patient is a candidate for surgical tumor resection of their target tumor(s)
7. The patient is pregnant or lactating
8. The patient has clinically significant renal or hepatic disease
9. The patient requires regular use of aspirin, full-dose warfarin, or heparin. Use of low-dose agents to maintain patency of vascular catheters is allowed

Date of first enrolment

16/12/2005

Date of final enrolment

30/11/2006

Locations

Countries of recruitment

Canada

United States of America

Study participating centre

4009 Banister Lane

Austin

United States of America

78704

Sponsor information

Organisation

Viventia Biotech Inc (Canada)

Sponsor details

5060 Spectrum Way
Suite 405
Mississauga
Canada
L4W 5N5

Sponsor type

Industry

ROR

<https://ror.org/0440s3562>

Funder(s)**Funder type**

Industry

Funder Name

Viventia Biotech Inc (Canada)

Results and Publications**Publication and dissemination plan**

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration