

Evaluation of gum and bone changes after immediate dental placement with two different surgical techniques

Submission date 22/07/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 23/07/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 23/07/2020	Condition category Oral Health	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

The introduction of dental implants as an option to replace missing teeth was considered one of the most revolutionary developments in modern dentistry. However, one of the main drawbacks with the first dental implants was the time required for the treatment to be completed. One of the stages that delayed the overall treatment considerably was the waiting time for the gum and the bone to heal once a tooth had been extracted (3-6 months) before a dental implant could be placed. Quite recently, a new approach was described in which a dental implant was placed immediately in the socket left after a tooth was removed. This proved to be a very valid modality of treatment that reduced considerably the time needed to place an implant. As surgical techniques evolved, it was noticed that the less trauma caused during the immediate implant placement, the better the healing was. This led to the consideration of placing immediate dental implants without lifting the gums due to the potential advantages this may have for the healing of the gum and the bone afterwards. At the present, there are not many studies looking into this surgical aspect of immediate dental implants, so the aim of this study is to compare the gum and bone healing in immediate dental implants placed with and without gum lifting/flap.

Who can participate?

Patients aged over 18 requiring single tooth extraction in the upper anterior and premolar region of the mouth

What does the study involve?

Participants are randomly allocated to receive immediate dental implants placed with and without gum lifting/flap. Both groups will receive hard and soft tissue grafting as part of the procedure. Dental implant survival and success are measured after 1 year.

What are the possible risks and benefits of participating?

The main benefit of this study is providing the participants with a dental implant to replace a missing tooth instead of another treatment modality that would not suit their needs or may have a negative impact in neighbouring teeth, such as fixed dental prosthesis. The potential risks

associated with this treatment modality are mainly associated with the minor surgical procedure needed to place a dental implant and the unlikely scenario in which a dental implant failed. The main items are listed below:

Swelling of the lips, chin, cheeks, and other tissues of the face, mouth, and neck with possible bruising and discoloration. Damage to and possible loss of other teeth, fillings, or other dental work, which will require root canal treatment, tooth extraction, or new fillings or restorations. Infection which will require antibiotic treatment, possibly further surgery, and may lead to loss of the implant. Discomfort which may require the use of pain medication for several days or longer, and which will also require at-home recuperation for the same period of time or longer. Bleeding which may be prolonged or heavy, and may require additional surgery or other treatment. Sinus or nasal problems, including but not limited to nasal stuffiness, bleeding, infection, or creation of an opening between the nose or sinus and mouth, which may require surgery. Poor healing of the gum tissue which may result in exposure of the implant and its loss. Loss or resorption of the bone which may result in failure of the implant. Injury of the nerves near the treatment site which cause pain, numbness, or tingling of the lips or areas of the face. This effect is usually temporary but may be permanent. Stretching or damage to the corners of the mouth with cracking, bleeding, and bruising.

Where is the study run from?

Universidad de Murcia (Spain)

When is the study starting and how long is it expected to run for?

October 2016 to May 2020

Who is funding the study?

1. Universidad de Murcia (Spain)
2. Avivent (Spain)
3. Osteogenoss (Spain)

Who are the main contacts?

1. Dr Ruben Garcia
ruben.garciasanchez@gstt.nhs.uk
2. Dr Guillermo Pardo Zamora
gparza@um.es

Contact information

Type(s)

Scientific

Contact name

Dr Ruben Garcia Sanchez

ORCID ID

<http://orcid.org/0000-0003-2051-0668>

Contact details

Periodontology Unit
Floor 25
Guy's Hospital
Great Maze Pond

London
United Kingdom
SE1 9RT
+44 (0)7864929396
ruben.garciasanchez@gstt.nhs.uk

Type(s)
Scientific

Contact name
Dr Guillermo Pardo Zamora

Contact details
Facultad de Odontología - Universidad de Murcia Hospital Morales Meseguer
Avda. Marqués de los Vélez s/n
Murcia
Spain
30008
+34 (0)606064848
gparza@um.es

Additional identifiers

EudraCT/CTIS number
Nil known

IRAS number

ClinicalTrials.gov number
Nil known

Secondary identifying numbers
1738/2017

Study information

Scientific Title
Evaluation of soft and hard tissue remodelling in post-extraction immediate dental implants placed with a flap and flapless approach in the aesthetic region. Clinical and radiographic prospective evaluation at one year.

Study objectives
The aim of this study is to test the main null hypothesis that there are no differences in survival /success rate in a group of patients requiring single immediate DI in the anterior maxillary region treated with a flapless vs. flap approach against the alternative hypothesis of a difference. For the secondary outcomes, the study tests the null hypotheses that there are no differences in "Pink aesthetic score (PES)", cortical plate resorption and "patients' reported outcomes (PROMS)" in a group of patients requiring single immediate DI in the anterior maxillary region treated with a flapless vs. flap approach against the alternative hypothesis of a difference.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 01/03/2018, ethical committee of Murcia University (Comité Ético de Investigación Clínica (CEIC); Calle Santo Cristo, 1, Murcia, Spain, 30001; +34 (0)868883614; comision.etica.investigacion@um.es), ref: 1738/2017

Study design

Single-center randomized controlled parallel-arm blind study

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Soft and hard tissue remodelling in post-extraction immediate dental implants

Interventions

Patients requiring single tooth extraction in the anterior and premolar areas were recruited for this study. Self-tapping, screw-shaped Biomimetic OCEAN, Avinent® implants of different lengths and diameters were immediately placed after dental extraction. In the control group, implant placement was performed with the elevation of a mucoperiosteal flap whereas in the test group this was performed flawless. In all study cases, test and control, discrepancies between the extraction socket and the implant surface were filled with a mix of bone replacement graft (Gen-Os, Osteógenos s.r.l.) and autologous bone, covered with a collagen membrane (Evolution, Osteógenos, s.r.l.). In addition, soft tissues were augmented by the use of autogenous connective tissue graft.

Randomization was done by balanced block randomization using a computer-generated table with random numbers. Treatment assignment was concealed to the treating surgeon by opaque envelopes that were opened only after completion of tooth extraction and final assessment of the feasibility of immediate implant placement. Clinical/radiographic measures, aesthetic evaluations and statistical analyses were performed blind with respect to treatment assignment.

Intervention Type

Procedure/Surgery

Primary outcome measure

1. Dental implant survival, defined as percentage of dental implants that presented successful osseointegration, lack of complications and bone loss less than 0.2 mm/year after loading at 1-year post-loading
2. Dental implant success, defined as percentage of implants that survived and the pink aesthetic score was above 7 and the white aesthetic score was above 5 at 1-year post-loading

Secondary outcome measures

1. Aesthetic of gums measured using the pink aesthetic index at 1-year post-loading
2. Aesthetic of false tooth measured using the white aesthetic index at 1-year post-loading
3. Buccal plate resorption measured using cone-beam CT at 1-year post-loading
4. Patients' reported outcomes measured using questionnaire and visual analogue scale at 1-year post-loading

Overall study start date

01/10/2016

Completion date

22/05/2020

Eligibility

Key inclusion criteria

1. Healthy individuals above 18 years who underwent immediate implant placement following single tooth extraction and were intended to receive implants
2. Non-smokers
3. Enough quantity of native bone to achieve primary stability
3. Patient has adequate mesiodistal space for implant placement
4. Patient available for follow-up according to protocol for 12 months post-loading to be able to assess the results of the proposed investigation

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

30

Total final enrolment

28

Key exclusion criteria

1. Medically compromised patients
2. Smokers
3. Presence of active periodontal disease
4. Symptomatic apical pathology before extraction

Date of first enrolment

20/04/2018

Date of final enrolment

29/05/2019

Locations

Countries of recruitment

Spain

Study participating centre**Universidad de Murcia**

Universidad de Murcia Hospital Morales Meseguer

Avda. Marqués de los Vélez s/n

Murcia

Spain

30008

Sponsor information

Organisation

Hospital General Universitario Morales Meseguer

Sponsor details

Facultad de Odontología - Universidad de Murcia Hospital Morales Meseguer

Avda. Marqués de los Vélez s/n

Murcia

Spain

30008

+34 (0)868888584

gparza@um.es

Sponsor type

Hospital/treatment centre

Website

<http://www.murciasalud.es/seccion.php?idsec=367>

ROR

<https://ror.org/00cfm3y81>

Funder(s)

Funder type

University/education

Funder Name

Universidad de Murcia

Alternative Name(s)

University of Murcia

Funding Body Type

Government organisation

Funding Body Subtype

Local government

Location

Spain

Funder Name

Avinent

Funder Name

Osteógenos s.r.l.

Results and Publications

Publication and dissemination plan

Planned publication in a high-impact peer-reviewed journal. Study protocol, PIS, the results of any statistical analysis, statistical analysis plan are provided upon request and under the conditions stated.

Intention to publish date

01/11/2020

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Dr. Guillermo Pardo Zamora (gparza@um.es). This data will become available once the study results are published. The data available will be all of the individual participant data collected during the trial, after deidentification. This data will be available 3 months after publication and ending 5 years following article publication. This data will be shared with investigators whose proposed use of the data has been approved by an independent review committee identified for this purpose. The data will be shared to achieve the aims in the approved proposal. Proposals should be directed to Dr Guillermo Pardo Zamora (gparza@um.es). To gain access, data requestors will need to sign a data access agreement.

IPD sharing plan summary

Available on request