

# Evaluation of the effectiveness of the ePATH Guided Re-entry Catheter Kit in bypassing a blockage of a leg artery

<b>Submission date</b> 11/03/2020	<b>Recruitment status</b> Stopped	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 02/07/2020	<b>Overall study status</b> Stopped	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 17/01/2022	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Patients who have no blood flow in their leg arteries because they are blocked can be treated via special tubes called catheters that can be inserted into the artery. First an attempt is made to pierce the blockage with a sharp wire that goes inside the catheter. If this is not possible then a wire is passed around the blockage by going inside the wall of the artery. Once past the blockage, the wire needs then to be redirected back into the downstream artery in order to reconnect to the main blood stream, a channel for blood flow that bypasses around the blockage can be made by implanting special tubes called grafts using the wire as a guide. Then blood will flow around the blockage back into the artery to reach the rest of the leg, and solve the problems caused by the blockage. One step in this process, the re-entry of the initial wire back into the vessel past the blockage, can sometimes be extremely challenging. The existing devices to achieve this re-entry can be difficult to use and can miss the artery. The aim of this study is to evaluate a new device, the ePATH Catheter Kit, which incorporates a new method of electronic guidance to assist the doctor to do this more efficiently and accurately.

### Who can participate?

Patients aged 18 to 90 who have an artery that needs unblocking (chronic total occlusion of the femoral, popliteal, external iliac or common iliac suitable for endovascular intervention [PTA or peripheral stenting])

### What does the study involve?

The treatment will follow the usual course, except the new ePATH Catheter Kit will be used to create a channel around the blockage instead of the existing devices. Once this is done the device will be removed and the treatment will continue unchanged, and the same graft will be implanted as before. The study will be carried out on 10 patients at Northwick Park Hospital in the second part of 2020 by the same doctor who normally carries out this treatment, the patient will not see any change to the care given or the number of hospital visits. The patient will attend the usual 6-week check-up after the treatment and leave the study at that point.

What are the possible benefits and risks of participating?

The potential benefit of this new approach is that the positioning of the wire around the blockage will be quicker, and less likely to miss the artery requiring re-positioning. There is a risk that the new ePATH Catheter Kit will not be any better at positioning the wire, if that happens then an existing device will be used. The new catheter has been thoroughly tested in the factory, but this is the first time it is used in patients which is why it is being used in a carefully observed study.

Where is the study run from?

Northwick Park Hospital (UK)

When is the study starting and how long is it expected to run for?

January 2020 to December 2020

Who is funding the study?

National Institute of Health Research (NIHR) (UK)

Who is the main contact?

Dr Robert Dickinson

robert.dickinson@imperial.ac.uk

## Contact information

### Type(s)

Public

### Contact name

Dr Robert Dickinson

### ORCID ID

<https://orcid.org/0000-0002-5999-7015>

### Contact details

Dept Of Bioengineering

Imperial College

Exhibition Road

London

United Kingdom

SW7 2AZ

+44 (0)20 7594 5662

robert.dickinson@imperial.ac.uk

### Type(s)

Scientific

### Contact name

Dr Robert Dickinson

### ORCID ID

<https://orcid.org/0000-0002-5999-7015>

## Contact details

Dept Of Bioengineering  
Imperial College  
Exhibition Road  
London  
United Kingdom  
SW7 2AZ  
+44 (0)20 7594 5662  
robert.dickinson@imperial.ac.uk

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

### Integrated Research Application System (IRAS)

274700

### ClinicalTrials.gov (NCT)

Nil known

### Protocol serial number

19IC5677, IRAS 274700

## Study information

### Scientific Title

Safety and performance evaluation of the ePATH guided percutaneous re-entry device for guiding the delivery of a guidewire around a chronic total occlusion of lower limb artery

### Acronym

eFACTOR

### Study objectives

To investigate whether the ePATH Catheter Kit and Display can be used to safely deliver a guidewire that bypasses a CTO.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Approved 21/05/2020, London - Dulwich Research Ethics Committee (Health Research Authority, Skipton House, 80 London Road. London SE1 6LH, UK; +44 (0)207 104 8241; dulwich.rec@hra.nhs.uk), REC ref: 20/LO/0330

### Study design

Prospective non-randomised single-arm single-centre observational study

### Primary study design

Observational

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Vascular disease

**Interventions**

Percutaneous bypass of Chronic Total Occlusions in arteries of the leg. This involves placing a guidewire around a chronic total occlusion via the sub-intimal space and back into the artery under the electronic guidance of the ePATH system. Subsequently, a graft is delivered over the guidewire and deployed to create a channel for blood perfusion to the distal vasculature. Follow up for 6 weeks.

**Intervention Type**

Device

**Phase**

Not Applicable

**Primary outcome(s)**

Measured immediately after the procedure:

1. Incidence of peri-procedural Major Adverse Events
2. Success rate of target lesion crossing defined by crossing with a 0.014 inches guide wire from true lumen to true lumen

**Key secondary outcome(s)**

Measured immediately after the procedure with the exception of late adverse events:

1. Number of deployment attempts until the guidewire enters true lumen
2. Distance between the distal cap and the re-entry point (using X-ray image post procedure and target catheter tip length as reference)
3. Time from first steps taken to insert the ePATH target catheter into vessel until successful lesion crossing, documented using fluoroscopic images/video timestamps
4. Incidence of late adverse events reported at 6-week follow-up

**Completion date**

31/12/2020

**Reason abandoned (if study stopped)**

Participant recruitment suspended and the study was ultimately closed due to public health guidance causing restrictions to study activities

**Eligibility****Key inclusion criteria**

1. Patient has a chronic total occlusion of the femoral, popliteal, external iliac or common iliac requiring treatment
2. Male or female patient, aged  $\geq 18$  and  $\leq 90$  years at time of enrolment
3. Suitable for endovascular intervention (PTA or peripheral stenting)
4. Able and willing to give written informed consent

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

**Key exclusion criteria**

1. Any of the following blood disorders, confirmed by the pre-assessment laboratory tests taken as standard of care, within the month before the procedure:

1.1. Anaemia with haemoglobin < 80g/L

1.2. Thrombopaenia with platelets  $\leq$  50 000/mL

1.3. Coagulation disorder with INR  $\geq$  1.5

2. If female, patient is pregnant or lactating at the time of enrolment or planning to become pregnant during the study time period

3. Patient has participated in another clinical study involving an investigational drug or investigational device within 30 days prior to enrolment or is scheduled to participate in another clinical study involving an investigational drug or investigational device during the course of this study. Patients enrolled in observational registries not involving investigational drug or investigational device may still be eligible

4. Patient is in custody or an institution

5. Patient has close affiliation with the study site or sponsor (e.g. employee, close relative of an employee)

**Date of first enrolment**

11/03/2020

**Date of final enrolment**

01/12/2020

**Locations****Countries of recruitment**

United Kingdom

England

**Study participating centre**

**Northwick Park Hospital**

LNWH NHS Trust Watford Road

Harrow

London  
United Kingdom  
HA1 3UJ

## Sponsor information

### Organisation

Imperial College London

### ROR

<https://ror.org/041kmwe10>

## Funder(s)

### Funder type

Government

### Funder Name

National Institute for Health Research

### Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

### Funding Body Type

Government organisation

### Funding Body Subtype

National government

### Location

United Kingdom

## Results and Publications

### Individual participant data (IPD) sharing plan

Patient identifiable data will be restricted to the study site, Northwick Park Hospital, and will be in hard copy form. Patients will be asked to consent to their data being used for this study. Requests for access to the datasets should be made to Simon Lewis, Research Governance Manager LNWH NHS Trust, R&D Office, Northwick Park Hospital, Watford Road, Harrow HA1 3UJ, Tel: +44 (0)20 8869 5173, email: [simon.lewis4@nhs.net](mailto:simon.lewis4@nhs.net). Anonymised data and summarised data will be made available to the Chief Investigator Dr R J Dickinson, Imperial College.

## IPD sharing plan summary

Available on request

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">HRA research summary</a>			28/06/2023	No	No