

Minimizing needle poke pain in newborn infants with a pain relieving cream and sugar water

Submission date 22/08/2007	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 23/08/2007	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 18/04/2008	Condition category Signs and Symptoms	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Anna Taddio

Contact details
Department of Pharmacy and Child Health Evaluative Sciences
The Hospital for Sick Children
555 University Avenue
Toronto, Ontario
Canada
M5G 1X8
+1 416 783 5263
anna.taddio@sickkids.ca

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
MCT-82947

Study information

Scientific Title

Evaluation of liposomal lidocaine and oral sucrose for treatment of pain in newborn infants undergoing venipuncture: a randomised controlled trial

Study objectives

Public title: Minimising needle poke pain in newborn infants with a pain relieving cream and sugar water

Hypothesis 1: Sucrose plus liposomal lidocaine will be superior to either agent alone in reducing pain during venipuncture.

Hypothesis 2: Plasma levels of lidocaine will be below toxicologically significant levels (1 mcg /ml), providing objective evidence of safety.

Hypothesis 3: Plasma levels of endomorphins-1,-2 will be higher in infants pre-medicated with sucrose versus lidocaine, confirming that sucrose exerts its analgesic effects via an opioid-mediated mechanism.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval was gained from the Research Ethics Board of Mount Sinai Hospital Toronto, Ontario, Canada on August 03, 2007 (ref: 07-0099-A).

Study design

Randomised, controlled, double-dummy, single-centre, three arm trial with study participant and investigator, caregiver, outcome assessor, and data analyst blinded.

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Newborns undergoing painful procedure

Interventions

Experimental arm one: 1 g of liposomal lidocaine to the dorsum of hand for 30 - 40 minutes prior to venipuncture and 2 ml of placebo water by mouth using a syringe over 1 - 2 minutes prior to venipuncture.

Experimental arm two: 2 ml of 24% sucrose administered by mouth using a syringe over 1 - 2 minutes prior to venipuncture and 1 g of placebo liposomal lidocaine to the dorsum of hand for 30 - 40 minutes prior to venipuncture.

Experimental arm three: both 1 g of liposomal lidocaine to the dorsum of hand for 30 - 40 minutes prior to venipuncture and 2 ml of 24% sucrose administered by mouth using a syringe over 1 - 2 minutes prior to venipuncture.

Due to double dummy, all infants receive oral sucrose or water (placebo) and all infants receive liposomal lidocaine or placebo. Each baby gets one intervention (experimental arm 1, 2, or 3).

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Liposomal lidocaine, oral sucrose

Primary outcome measure

Infant pain during venipuncture as assessed by facial grimacing response - the score will incorporate three facial actions: brow bulge, eye squeezed shut, naso-labial furrow, during venipuncture.

Secondary outcome measures

1. Visual Analog Scale (VAS), during venipuncture
2. Cry duration, during venipuncture
3. Heart rate, during venipuncture
4. Number of attempts until procedure completion, from first needle poke to completion
5. Endomorphins -1, -2, levels, before and 10 minutes after sucrose/sucrose placebo administration
6. Procedure duration, from first needle poke to completion
7. Lidocaine levels, 5 - 15 minutes after the liposomal lidocaine cream/placebo cream is removed

Overall study start date

01/08/2007

Completion date

28/02/2010

Eligibility

Key inclusion criteria

All healthy full-term (greater than 37 weeks gestational age) newborn infants (either sex)

Participant type(s)

Patient

Age group

Neonate

Sex

Both

Target number of participants

330

Key exclusion criteria

1. Neonatal Intensive Care Unit (NICU) admission
2. Asphyxia, seizures
3. Major birth defects (heart, brain, genetic syndrome)
4. Circumcised during study
5. Receiving analgesia/sedatives

Date of first enrolment

01/08/2007

Date of final enrolment

28/02/2010

Locations

Countries of recruitment

Canada

Study participating centre

Department of Pharmacy and Child Health Evaluative Sciences

Toronto, Ontario

Canada

M5G 1X8

Sponsor information

Organisation

The Hospital for Sick Children (Canada)

Sponsor details

c/o Ms. Julie Gibson, Manager

Clinical Research Office

555 University Avenue

Toronto, Ontario

Canada

M5G 1X8
+1 416 813 8481
julie.gibson@sickkids.ca

Sponsor type

Hospital/treatment centre

Website

<http://www.sickkids.ca/>

ROR

<https://ror.org/057q4rt57>

Funder(s)

Funder type

Research organisation

Funder Name

Canadian Institutes of Health Research (CIHR) (Canada) - <http://www.cihr-irsc.gc.ca> (ref: MCT-82947)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration