Comparision of bone-drilling techniques in the management of long bone infection following fracture repair by metal rod insertion

Submission date	Recruitment status No longer recruiting	Prospectively registered		
23/07/2019		☐ Protocol		
Registration date 29/07/2019	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited 10/01/2020	Condition category Musculoskeletal Diseases	Individual participant data		

Plain English summary of protocol

Background and study aims

Treatment for fractures of long bones in the leg (femur and tibia) usually involves the insertion fo a metal rod into the bone cavity to provide additional strength and support (intermedullary nailing). In one to two percent of cases the bone cavity can become infected. Treating the infected area involves removal of the metal rod and drilling out (reaming) the infected bone. The aim of this study is to compare the postoperative effects of two methods of intermedullary reaming.

Who can participate?

Patients aged over 18 years diagnosed with an infection of the bone cavity previously treated by insertion of a metal rod into the bone.

What does the study involve?

Patients undergo surgical removal of the infected metal rod followed by either conventional reaming or reaming using the Reamer / Irrigator / Aspirator (RIA) system (Synthes, Inc. West Chester, Philadelphia, PA, USA). Patients then attended routine follow up visits to the hospital at 1, 3, 6, 12 and 24 months after the operation.

What are the possible benefits and risks of participating? During the research, there is no direct benefit to the participant

Where is the study run from? Federal University of São Paulo - Hospital São Paulo (UNIFESP)

When is the study starting and how long is it expected to run for? August 2013 to July 2015

Who is funding the study? Universidade Federal de São Paulo Who is the main contact? Mauro Salles mauro.salles@unifesp.br

Contact information

Type(s)

Scientific

Contact name

Mr Mauro Salles

ORCID ID

https://orcid.org/0000-0002-5443-6024

Contact details

Rua Visconde de Ouro Preto 77, apto193 São Paulo Brazil 01303-060 +5511985360055 mauro.salles@unifesp.br

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

354.934

Study information

Scientific Title

The intramedullary reaming modality for the management of postoperative long bone infection: a prospective randomized controlled trial

Study objectives

To compare the efficacy of reamer/irrigator/aspirator (RIA) with conventional reaming followed by insertion of antibiotic-loaded cement, for the treatment of intramedullary nail infection of long bones

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 16/08/2013, Research and Ethics Committee of the Federal University of São Paulo (UNIFESP) (Rua Botucatu, 572 1° Andar Conj. 14, Vila Clementino, Sao Paulo, Brazil; +55 115539-7162; cepunifesp@unifesp.br), ref: 354.934

Study design

Prospective randomized controlled single-blinded clinical trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Intramedullary nail infection

Interventions

A noninferiority, randomized clinical trial was carried out involving patients of whom a locked IMN implant of the femur and/or tibia was retrieved and who all met the clinical and radiological criteria for IMN-associated osteomyelitis. Patients were randomized into two groups: RIA alone versus conventional reaming followed by antibiotic-loaded cement insertion. Both groups also underwent six-weeks of antibiotic treatment according to the results of the antibiogram. Patients were evaluated after 1, 3, 6, 12 and 24 months for radiological and clinical follow-up.

Consecutive patients with clinically suspected intramedullary infection who underwent nailing explantation and surgical debridement were randomised to receive either conventional reaming or reamer/irrigator/aspirator (RIA) of the presumed infected medullary canal.

Patients presenting intramedullary infection were assessed at the trauma unit and the outpatient setting by the infectious disease specialist and recruited after signed consent. Allocation of patients to the treatment groups took place at the time of anesthesia within the operative room. Then, randomization to conventional reaming or RIA was set up in a simple randomization procedure (1:1 ratio) with sealed opaque envelopes, sequentially numbered. The CONSORT guidelines were followed throughout the study.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Intramedullary infection two-year remission rate, performed using all randomized patients that completed at least one-year of follow-up in the per-protocol (PP) analysis. Patients were considered in remission when there was an absence of clinical, laboratory and radiological signs of infection, assessed in the last medical consultation (at least 12 months follow-up). Cases that did not require re-operation or further antibiotic administration for the same site of infection were also considered to be in remission.

Key secondary outcome(s))

Preoperative and postoperative pain measured by the Visual Analog Scale (EVA) scale at baseline and 1, 3, 6, 12 and 24 months postoperatively

Completion date

Eligibility

Key inclusion criteria

- 1. 18 years and over
- 2. Previous tibia or femur intramedullary locking nail fixation who met the definitive clinical and radiological diagnosis of osseous infection

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

44

Key exclusion criteria

- 1. Diaphyseal diameter < 10 mm
- 2. Previous infection of the affected bone and patients with HIV or chronic renal failure were excluded from the study

Date of first enrolment

30/08/2013

Date of final enrolment

20/07/2015

Locations

Countries of recruitment

Brazil

Study participating centre

Federal University of São Paulo - Hospital São Paulo (UNIFESP)

Rua Napoleão de Barros 715 Vila Clementino São Paulo

Sponsor information

Organisation

Federal University Of São Paulo (UNIFESP)

ROR

https://ror.org/02k5swt12

Funder(s)

Funder type

University/education

Funder Name

Universidade Federal de São Paulo

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request. Carlos Augusto Finelli (email: c.finelli@cecot.com. br). The type of data is a complete Excel datasheet. The data is already available upon request and will become available for a long as necessary. Data will be shared upon reasonable request. The criteria access would be a simple email contact with the corresponding author, explaining the reasons for accessing the data. Additional analysis may also be requested upon reasonable rational. No further participants consent will be required including additional comments on data anonymization, as it has already been gathered and declared.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	02/12/2019	10/01/2020	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes