

Study comparing a regional leg block vs. local anaesthetic infiltration for leg donor sites in burns patients - pain scores and analgesic requirements

| | | |
|--|---|--|
| Submission date 22/03/2016 | Recruitment status No longer recruiting | <input type="checkbox"/> Prospectively registered |
| | | <input type="checkbox"/> Protocol |
| Registration date 04/05/2016 | Overall study status Completed | <input type="checkbox"/> Statistical analysis plan |
| | | <input type="checkbox"/> Results |
| Last Edited 17/08/2018 | Condition category Injury, Occupational Diseases, Poisoning | <input type="checkbox"/> Individual participant data |
| | | <input type="checkbox"/> Record updated in last year |

Plain English summary of protocol

Background and study aims

In the UK about 250,000 people sustain burn injuries each year. Of these, 175,000 attend accident and emergency departments and 13,000 need to be admitted to hospital. During treatment patients might need to be taken to theatre for the burn wound to be cleaned thoroughly and also to apply a skin graft to the wound. This will usually be done under a general anaesthetic. The skin graft is a piece of skin taken from the patient's unburnt skin in order to cover the burn wound and promote healing. This piece of skin is usually taken from the thigh. The amount of skin that is taken will depend on the size of the burn that needs to be covered. The site that the skin graft is taken from can be a cause of more pain than the original burn. In order to prevent this pain the surgeons sometimes use a local anaesthetic to make the skin numb. Sometimes the anaesthetists might give a local anaesthetic block called the fascia iliaca compartment block to prevent pain. This involves injecting local anaesthetic drugs into the front of the thigh near the groin crease. It is a single injection and the exact place of injection near the groin crease is guided by using ultrasound. This injection of local anaesthetic blocks the nerves that detect sensation from the thigh and thereby give pain relief. We wish to find out whether this block (fascia iliaca compartment block) is better at providing pain relief to the donor site (where the skin graft was taken) than local anaesthetic infiltration. We also want to find out if this block might provide a longer period of pain relief.

Who can participate?

Adult patients with burns undergoing skin grafts where the donor site is the thigh.

What does the study involve?

Participants are randomly allocated to be treated with either fascia iliaca compartment block or local anaesthetic infiltration. Participants are offered additional pain relief medications after the surgery, if required. All other treatment is the same as standard practice. Participants' pain and use of painkillers is assessed.

What are the possible benefits and risks of participating?

The fascia iliaca block is a very safe local anaesthetic block, which has been used for many years to provide pain relief to problems in the thigh region such as femoral fractures and hip fractures. It has also been used in burn centres to provide pain relief to the burn donor site. The risk is very low. Those patients who receive a block will find that their leg is numb and weak for some time after the block, making walking difficult for a few hours. Most of the time the patient would be kept on bed rest anyway for the grafts to heal. There is also a very small risk of infection and nerve damage but these risks are further minimised because we do this block under ultrasound guidance. The risk of nerve damage is small. The block when performed correctly is not in the path of any major nerve. Local anaesthetic infiltration around the donor site is also a long-standing technique that is used and side effects are rare. The patient will be carefully monitored in theatre and the ward regardless of the type of pain relief that they receive.

Where is the study run from?

Mid Essex Hospital Services NHS Trust (UK)

When is the study starting and how long is it expected to run for?

June 2014 to May 2017

Who is funding the study?

Mid Essex Hospital Services NHS Trust (UK)

Who is the main contact?

1. Dr Malka Sandunmalee Liyanage
2. Dr Peter Berry

Contact information

Type(s)

Public

Contact name

Dr Malka Sandunmalee Liyanage

Contact details

Broomfield Hospital
Court Road
Chelmsford
United Kingdom
CM1 8ET

Type(s)

Public

Contact name

Dr Peter Berry

Contact details

Broomfield Hospital
Court Road
Chelmsford

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

R&D 1012

Study information

Scientific Title

Study of fascia iliaca blocks vs. conventional local anaesthetic infiltrations for leg donor sites in burns patients – pain scores and analgesic requirements

Acronym

FICB

Study objectives

Patients with burns often require skin grafts to the site of the burn. A common place to take the skin graft from is the side of the thigh. It is often seen that the pain from the site where the skin graft is taken is more intense than that of the burn wound itself.

Fascia Iliaca Compartment Block (FICB) is a safe and simple method of giving local anaesthetic agents to the groin to provide pain relief for surgery to the thigh and hip. The local anaesthetic agents cause numbness by blocking the nerves supplying the hip and the thigh. It has been used to provide pain relief for patients with fractures of the neck of femur in emergency rooms for a long period of time and has a demonstrated safety profile.

Currently the most common method for providing pain relief after surgery to the skin graft is by infiltration of local anaesthetic to the tissues via multiple injections.

The purpose of the study is to find out if there is a difference in the pain relief obtained by the patients depending on whether they have a FICB or the direct local anaesthetic infiltration. This would be measured by the amount of oral pain relief medications they need postoperatively and also their level of pain as determined by a simple pain score.

Ethics approval required

Old ethics approval format

Ethics approval(s)

East of England - Cambridgeshire and Hertfordshire Research Ethics Committee, 09/02/2016, 15/EE/0066

Study design

Prospective partially blinded randomised controlled study

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet.

Health condition(s) or problem(s) studied

Pain in the donor sites of burns patients

Interventions

Patients coming to the burns or plastics theatres for skin grafting and where the graft is to be taken from the thigh would be included after applying the exclusion criteria. Burn size would be less than 7% and donor site >0.5% TBSA in order to meet the inclusion criteria. The anaesthetist in charge of the list on that day would have the discretion to either include or exclude the patient in the study.

If all criteria are satisfied, the patient would be consented for both the block and local infiltration at the same time as consenting to take part in the study.

Randomisation would be by random number tables (generated by Random.org) and sealed numbered envelopes to be picked up by the anaesthetist. These would state the arm of the study to which the patient would belong.

In those patients who are randomised to the block arm, the fascia iliaca block would be performed under ultrasound guidance. The block would be performed using 0.8 ml/kg 0.25% levobupivacaine as the local anaesthetic dosing regimen, with a maximum local anaesthetic volume of 40 ml.

For those patients randomised to local infiltration, the surgeons would directly infiltrate the same local anaesthetic regimen to the donor site, after the skin graft had been taken. The levobupivacaine may contain 1:200,000 adrenaline according to surgical preference, in order to limit blood loss.

Intraoperatively, the patient would have a target controlled infusion of Remifentanyl 3-5 ng/ml, titrated according to the responsible anaesthetists' judgement. Paracetamol 1g and Diclofenac 75 mg would be given intravenously if not contraindicated. Morphine 0.1 mg/kg-0.3 mg/kg will be given towards the end of the procedure. All other anaesthetic factors would be determined by the responsible anaesthetist. Should further analgesia be felt necessary, the responsible anaesthetist would document the rationale in the study proforma.

A standardised postoperative regimen of regular Paracetamol plus Ibuprofen if not contraindicated would be prescribed. Breakthrough analgesia in the form of Oromorph 0.2 mg/kg (rounded to the nearest 10 mg) m, max 1 hourly would also be available. Any pre-existing analgesia for the burn pain, for example Gabapentin or Clonidine, would be continued. Naloxone 100mcg prn would be prescribed in case of opiate toxicity.

Intervention Type

Procedure/Surgery

Primary outcome measure

The use of opiate painkillers at 24 and 48 hours. Our chosen threshold for a clinically significant difference will be a 20% difference in opiate usage at 24 and 48 hours between patients receiving fascia iliaca blocks, compared to those receiving the same volume and concentration of local anaesthetic via direct infiltration into the skin.

Secondary outcome measures

1. Maximum pain score 6 hours after surgery, first post-operative day and second post-operative day
2. Time to first dose of opiate medication after discharge from recovery
3. Patient satisfaction with post-operative analgesia, assessed at 24 and 48 hours
4. The density and duration of muscle weakness associated with fascia iliaca blocks by the use of bromage scores in the first and second post-operative day (does not include patients who are not allowed to move the grafted leg)

Overall study start date

27/06/2014

Completion date

01/12/2018

Eligibility

Key inclusion criteria

Adult patients with burns admitted to the burns service at St. Andrews with <7% burns and undergoing split skin grafts where the donor site will be the anterolateral thigh only and expected to be at least 0.5% total body surface area (TBSA).

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

60

Key exclusion criteria

1. Patients <16 years old
2. Patients unable to comment on the pain scores effectively e.g., unable to speak English
3. Patients unable to give informed consent e.g., dementia
4. Patients refusing consent for the fascia iliaca block
5. Patients requesting a particular form of analgesia
6. Patients previously on treatment for chronic pain conditions
7. Patients from whom multiple donor sites are anticipated
8. Patients with renal impairment sufficient to alter the planned opiate dosage
9. Patients with a current or previous history of opiate abuse

Date of first enrolment

01/03/2016

Date of final enrolment

01/03/2017

Locations

Countries of recruitment

England

United Kingdom

Study participating centre

Mid Essex Hospital Services NHS Trust

Broomfield Hospital

Court Road

Chelmsford

United Kingdom

CM1 7ET

Sponsor information

Organisation

Mid Essex Hospital Services NHS Trust (UK)

Sponsor details

Research and Development Department

Broomfield Hospital

Court Road

Broomfield

England

United Kingdom

CM1 7ET

Sponsor type

Hospital/treatment centre

ROR

<https://ror.org/0280r7h11>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Mid Essex Hospital Services NHS Trust (UK)

Results and Publications

Publication and dissemination plan

We are aiming to publish this study within the scientific community. All patient data will be anonymous.

Intention to publish date

01/12/2017

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Available on request

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|--------------------------------------|---------|--------------|------------|----------------|-----------------|
| HRA research summary | | | 28/06/2023 | No | No |