

# Clinical value and cost-effectiveness of intra-dialytic exercise for the improvement of health-related quality of life in people with stage 5 chronic kidney disease undergoing maintenance haemodialysis

<b>Submission date</b> 11/12/2014	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 17/12/2014	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 24/06/2021	<b>Condition category</b> Urological and Genital Diseases	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Hemodialysis is a form of treatment that replicates many of the kidney's functions, filtering the blood to rid the body of harmful waste, extra salt, and water. The aim of this study is to test whether exercise carried out during hemodialysis benefits patients in terms of their general wellbeing and quality of life and also if it benefits physical function and ability to participate in daily activities, like walking and rising from a chair. The effects of participation in the exercise programme on other aspects that may be important to the health of hemodialysis patients (for example, blood pressure) will also be examined. In addition, we will also examine whether this type of additional treatment option is cost effective within the health service setting.

### Who can participate?

Adult patients from dialysis units throughout the UK will participate in this study.

### What does the study involve?

Once patients have agreed to take part they will complete a series of assessments to measure their quality of life and functional limitations. After this, patients will be allocated to either a usual care haemodialysis plus exercise group or a usual care haemodialysis group. The exercise group will receive the opportunity to exercise during their dialysis for a total duration of 6 months. All patients will be invited to come back for reassessment of quality of life and functional limitations after 6, 9 and 15 months.

### What are the possible benefits and risks of participating?

The benefits of undertaking exercise training are that the patients are likely to feel better and healthier, have a potentially improved cardiovascular health, have improved physical, mental and emotional wellbeing, and an improved dialysis efficiency. All participants will have access to additional monitoring not available as part of routine care. For example, fitness and the health

of blood vessels will be assessed. After the study patients will be able to use this information to guide their lifestyle choices, potentially impacting positively upon their health. For patients in the exercise group, there is a very small chance of getting a muscle strain or some joint pain following exercise. Also, blood sugar levels and blood pressure may drop during the exercise, which may make the patient feel sick or dizzy. Rarely, exercise can cause a change to heart function. To reduce the risks of exercise the patients' medical history will be checked to ensure that health risks do not outweigh any benefits.

Where is the study run from?

King's College Hospital, Salford Royal Hospital, Glasgow Western Infirmary, Royal Derby Hospital and Heartlands Hospital (UK).

When is the study starting and how long is it expected to run for?

January 2015 to February 2020

Who is funding the study?

NIHR Health Technology Assessment Programme - HTA (UK).

Who is the main contact?

Mrs Claire White (clairewhite4@nhs.net)

Professor Iain Macdougall (iain.macdougall@nhs.net)

## Contact information

### Type(s)

Public

### Contact name

Mrs Claire White

### Contact details

King's College Hospital

Denmark Hill

London

United Kingdom

SE5 9RS

+44 (0)2032999000

clairewhite4@nhs.net

### Type(s)

Scientific

### Contact name

Prof Iain Macdougall

### Contact details

King's College Hospital

Denmark Hill

London

United Kingdom

SE5 9RS

+44 (0)2032999000  
iain.macdougall@nhs.net

## Additional identifiers

ClinicalTrials.gov (NCT)  
NCT02222402

Protocol serial number  
HTA 12/23/09

## Study information

### Scientific Title

PrEscription of intra-Dialytic exercise to improve quAlity of Life in patients with chronic kidney disease (PEDAL)

### Acronym

PEDAL

### Study objectives

To determine, in comparison to usual care, whether usual care augmented by intra-dialytic exercise training improves health-related quality of life in stage 5 CKD patients receiving maintenance haemodialysis renal replacement therapy.

More details can be found at <http://www.nets.nihr.ac.uk/projects/hta/122309>

Protocol can be found at [http://www.nets.nihr.ac.uk/\\_\\_data/assets/pdf\\_file/0006/130659/PRO-12-23-09.pdf](http://www.nets.nihr.ac.uk/__data/assets/pdf_file/0006/130659/PRO-12-23-09.pdf)

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

London – Fulham, 27/11/2014, ref: 14/LO/1851

### Study design

Multi-centre pragmatic single-blind randomised controlled trial

### Primary study design

Interventional

### Study type(s)

Treatment

### Health condition(s) or problem(s) studied

Chronic kidney disease

### Interventions

Once patients have agreed to take part they will complete a series of assessments to measure their self-reported QOL and functional limitations. After this, patients will be allocated to either

a usual care haemodialysis plus intradialytic exercise group or a usual care haemodialysis group. The exercise group will receive the opportunity to exercise during their dialysis for a total duration of 6 months. All patients will be invited to come back for reassessment of QOL and functional limitations after 6, 9 and 15 months.

## **Intervention Type**

Mixed

## **Primary outcome(s)**

A change in Kidney Disease QOL questionnaire Physical Composite Score (KDQOL-PCS) between baseline and 6 months. The KDQOL is a disease-specific quality of life measure.

## **Key secondary outcome(s)**

Assessed at 6, 9 and 15 months. Change between baseline and follow-up for:

1. KDQOL-PCS
2. KDQOL-MCS
3. KDQOL-Vitality subscale
4. KDQOL- symptom burden subscale
5. IPAQ
6. Height (at screening visit only), weight and anthropometric (BMI, waist to hip ratio)
7. Peak aerobic capacity
8. Duke Activity Status Index (self-report 12-item activity of daily living questionnaire)
9. 10 m Timed-up-and-go (composite test of leg strength, balance, coordination and gait speed)
10. Sit-to-stand 60 (proxy measure of lower extremity muscular endurance)
11. Tinetti Falls Efficacy Scale
12. Habitual physical activity (a self-report assessment via International Physical Activity Questionnaire)
13. Habitual physical activity (GT3X tri-axial accelerometer)
14. Arterial stiffness (via pulse wave velocity and augmentation index)
15. Blood pressure
16. Blood lipid levels – LDL, HDL, triglyceride levels
17. Hb
18. ESAs
19. HBA1c
20. CRP
21. Bicarbonate
22. Phosphate
23. Parathyroid hormone (PTH)
24. Dialysis efficiency
25. Change in medication use (vascular and diabetic medications)
26. EQ-5D, a generic multi-attribute health-related QOL questionnaire for use in cost-utility analysis
27. Health Economic questionnaire regarding resource use
28. All hospitalisations
29. All-cause mortality
30. Cardiovascular mortality
31. Safety/harms (AE, SAE)

## **Completion date**

29/02/2020

# Eligibility

## Key inclusion criteria

1. Prevalent Stage 5 CKD patients (GFR <15 mL/min) receiving maintenance haemodialysis therapy for more than 3 months
2. Male or female
3. Aged >18 years
4. Able to provide written informed consent

## Participant type(s)

Patient

## Healthy volunteers allowed

No

## Age group

Adult

## Lower age limit

18 years

## Sex

All

## Total final enrolment

380

## Key exclusion criteria

1. Patients unlikely to be on HD for > 6 months (this includes cachectic patients, those with severe heart failure, patients in whom dialysis withdrawal is being considered, and patients likely to receive a live-donor transplant or transfer to PD in that period of time)
2. Less than 3 months after the initiation of haemodialysis (patients in this time-frame are generally less clinically stable, many having vascular access procedures performed, and rates of inter-current events, including death and hospitalisation, are very much higher in the first 3 months after commencement of chronic haemodialysis)
3. Deemed to be clinically unstable by treating physician
4. Bilateral lower limb amputees
5. Dementia or severe cognitive impairment (as will be unable to give consent and/or complete questionnaire assessments)
6. Severe psychiatric disorders – except treated stable

## Date of first enrolment

01/01/2015

## Date of final enrolment

30/09/2015

# Locations

## Countries of recruitment

United Kingdom

England

Scotland

**Study participating centre**

**King's College Hospital**

Denmark Hill

London

United Kingdom

SE5 9RS

**Study participating centre**

**Salford Royal Hospital**

Stott Lane

Salford

United Kingdom

M6 8HD

**Study participating centre**

**Glasgow Western Infirmary**

Dumbarton Road

Glasgow

United Kingdom

G11 6NT

**Study participating centre**

**Royal Derby Hospital**

Uttoxeter New Road

Derby

United Kingdom

DE22 3NE

**Study participating centre**

**Heartlands Hospital**

Bordesley Green East

Birmingham

United Kingdom

B9 5SS

# Sponsor information

## Organisation

King's College Hospital NHS Trust

## ROR

<https://ror.org/01n0k5m85>

# Funder(s)

## Funder type

Government

## Funder Name

National Institute for Health Research

## Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

## Funding Body Type

Government organisation

## Funding Body Subtype

National government

## Location

United Kingdom

# Results and Publications

## Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		10/09/2020	10/05/2021	Yes	No
<a href="#">Funder report results</a>		01/06/2021	24/06/2021	No	No
<a href="#">HRA research summary</a>			28/06/2023	No	No