Depression: a trial of antenatal guided self help for women

Submission date 08/08/2014	Recruitment status No longer recruiting	[X] Prospectively registered		
		[X] Protocol		
Registration date 08/08/2014	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited 07/08/2020	Condition category Pregnancy and Childbirth	Individual participant data		

Plain English summary of protocol

Background and study aims

Pregnancy does not appear to be protective against the existing or development of new mental disorders. Depression is the most common antenatal mental disorder and can have a considerable impact upon a woman and her family. To date, there has been no study of mild psychological interventions in the antenatal setting which may be more effective and cost-effective for mild and moderate depression during pregnancy. We therefore propose to modify and evaluate Guided Self Help materials for antenatal depression in women using NHS maternity services. This study aims to get initial evidence on the how effective is the Guided Self Help intervention in improving depressive symptoms and other outcomes (e.g. psychological symptoms after giving birth and quality of life) for women with antenatal depression.

Who can participate?

Adult pregnant women who are up to 26 weeks pregnant and having depressive symptoms can take part.

What does the study involve?

Pregnant women experiencing depression will be randomly allocated to either antenatal Guided Self Help (given by an NHS Psychological Wellbeing Practitioner) plus usual care, or to usual care alone. The antenatal Guided Self Help intervention has been modified from existing Guided Self Help materials used in psychological services for the treatment of depression. Modifications include addressing pregnancy-specific worries, and including sections on health issues in pregnancy and planning for parenthood. Women allocated to the this group will be seen for up to eight sessions by a Psychological Wellbeing Practitioner (including an initial assessment session); there will also be a appointment at 12 weeks after delivery. Research measures will be taken from all women before random allocation, after 14 weeks of allocation and at 12 weeks after delivery. Data will also be collected to find out if this method is cost-effective.

What are the possible benefits and risks of participating?

The opportunity to talk about symptoms with a researcher can be beneficial and, where appropriate, participants will be encouraged to discuss concerns with their midwife and given information on sources of help and support, which may be helpful in managing their situation. Some participants may find that being asked questions about their mental health and wellbeing

is sensitive, and the potential for the interview to bring up these feelings is described on the Participant Information Sheet. At the beginning of the interview, participants will be asked if there is anyone they would like the researcher to contact for support if they become distressed. Participants will be closely monitored during interviews for signs for distress and appropriate action will be taken. If there are concerns about the health and wellbeing of participants or their children at the end of the interview, participants will be asked if they would like any of the information they disclosed to be conveyed to their clinic midwife.

Where is the study run from?
The study is being run from Kings Health Partners (UK)

When is the study starting and how long is it expected to run for? October 2014 to June 2017

Who is funding the study? National Institute for Health Research (NIHR) (UK)

Who is the main contact? Ms Kylee Trevillion kylee.trevillion@kcl.ac.uk

Contact information

Type(s)

Scientific

Contact name

Ms Kylee Trevillion

ORCID ID

https://orcid.org/0000-0001-5865-4726

Contact details

Section of Women's Mental Health PO 31, 16 De Crespigny Park London United Kingdom SE5 8AF

kylee.trevillion@kcl.ac.uk

Additional identifiers

Protocol serial number 17048

Study information

Scientific Title

Depression: an exploratory parallel-group randomised controlled trial of Antenatal guided self help for WomeN (DAWN)

Acronym

DAWN Trial

Study objectives

Women with mild or moderate antenatal depression treated with Guided Self Help will have significantly lower Edinburgh Postnatal Depression Scale (EPDS) depressive symptoms at 14 weeks post-randomisation compared with women with mild or moderate antenatal depression receiving usual care.

Ethics approval required

Old ethics approval format

Ethics approval(s)

NRES Committee London-Camberwell St Giles, 11/06/2014, ref: 14/LO/0597

Study design

Randomised; Interventional; Design type: Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: Mental Health, Reproductive health and childbirth; Subtopic: Depression, Reproductive Health and Childbirth (all Subtopics); Disease: Depression, Reproductive Health & Childbirth

Interventions

Women will be randomly assigned to:

- 1. Guided Self Help (delivered by Psychological Wellbeing Practitioners) plus usual care
- 2. Usual care alone (i.e., treatment as usual)

Guided Self Help, Details: Existing Guided Self Help materials used in psychological services for the treatment of depression are being modified for pregnancy. The Guided Self Help (with usual care) intervention for antenatal depression is being delivered by NHS Psychological Wellbeing Practitioners (PWPs). The intervention comprises an initial session at the beginning of therapy, followed by up to eight 30-minute sessions. An additional session will also be conducted, at six to eight weeks post-delivery (i.e. before the post-delivery research outcomes are collected)

Intervention Type

Other

Phase

Phase II

Primary outcome(s)

Depression, measured using the Edinburgh Postnatal Depression Scale - depressive symptoms at baseline and 14 weeks

Key secondary outcome(s))

- 1. Anxiety, measured using the Generalized Anxiety Disorder Scale (GAD-7) at baseline and 14 weeks
- 2. Depression, measured using the Patient Health Questionnaire-9 (PHQ-9) at baseline and 14 weeks and the Edinburgh Postnatal Depression Scale and Whooley items at 3 months post-delivery
- 3. Health-related quality of life, measured using the Short Form Health Survey (SF-6D) at baseline, 14 weeks and 3 months post-delivery
- 4. Service contacts, measured using the Adult Service Use Schedule at 14 weeks and 3 months post-delivery
- 5. Parenting stress, measured using the Postpartum Bonding Questionnaire at 3 months postdelivery

Completion date

30/06/2017

Eligibility

Key inclusion criteria

- 1. Pregnant women, up to 26 weeks gestation
- 2. Aged 16 years or older
- 3. Meeting the criteria for DSM--IV depression

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Total final enrolment

53

Key exclusion criteria

- 1. Pregnant women who are unable to complete the trial questionnaires/workbook in English or provide informed consent
- 2. Have psychosis, borderline personality disorder, a current eating disorder, current post-traumatic stress disorder, or suicidality and/or are receiving secondary mental health care, psychological therapies or antidepressants

Date of first enrolment

30/10/2014

Date of final enrolment

30/06/2016

Locations

Countries of recruitment

United Kingdom

England

Study participating centre Croydon Health Services

530 London Road Croydon London United Kingdom CR7 7YE

Study participating centre St Thomas' Hospital

10th floor North Wing Westminster Bridge Road London United Kingdom SE1 7EH

Study participating centre University Hospital Lewisham

Lewisham High Street London London United Kingdom SE13 6LH

Study participating centre South London and Maudsley NHS Foundation Trust

Monks Orchard Road Beckenham United Kingdom BR3 3BX

Study participating centre King's College Hospital

Denmark Hill London United Kingdom SE5 9RS

Sponsor information

Organisation

King's College London (UK)

ROR

https://ror.org/0220mzb33

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/01/2020	07/08/2020	Yes	No
<u>Protocol article</u>	protocol	18/10/2016		Yes	No
HRA research summary			28/06/2023		No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes