Non-surgical treatment of benign paroxysmal positional vertigo (BPPV) - a randomised controlled trial of procedures and practices

Submission date 12/09/2003	Recruitment status Stopped	Prospectively registered
		☐ Protocol
Registration date	Overall study status	Statistical analysis plan
12/09/2003	Stopped	Results
Last Edited	Condition category	Individual participant data
19/10/2011	Signs and Symptoms	Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr R Palaniappan

Contact details

Audiological Medicine Department, RNTNE Royal Free Hampstead NHS Trust 330 Grays Inn Road Kings Cross London United Kingdom WC1X 8DA +44 020 7915 1300 ext 4259 rudipal@hotmail.com

Additional identifiers

Protocol serial number

N0256111276

Study information

Scientific Title

Study objectives

How efficient are the different non-surgical treatments that are in current clinical practice? Is any one method better than the other?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Signs and Symptoms: Benign paroxysmal positional vertigo (BPPV)

Interventions

- 1. Particle repositioning manoeuvre alone
- 2. Particle repositioning manoeuvre plus soft collar & sleeping instructions
- 3. Particle repositioning manoeuvre with bone vibrator
- 4. Particle repositioning manoeuvre with bone vibrator plus soft collar & sleeping instructions
- 5. Control group treated with rehabilitation exercises

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

All the above described methods are recognized standard treatment procedures for BPPV in current clinical practice. Patients will be reviewed in the clinic 8 weeks post-treatment, when they will be required to answer the Dizziness Handicap Inventory (DHI). Questionnaire. A negative Hallpike's test during this visit would be considered as a positive outcome.

Key secondary outcome(s))

Not provided at time of registration

Completion date

31/07/2003

Reason abandoned (if study stopped)

Poor recruitment

Eligibility

Key inclusion criteria

Outpatients who have a positive Dix-Hallpike's positional test diagnostic of BPPV Five groups, approx 20 patients per group. 5 Patients recruited 2001- 2002.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Not Specified

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/02/2002

Date of final enrolment

31/07/2003

Locations

Countries of recruitment

United Kingdom

England

Study participating centre Audiological Medicine Department, RNTNE

London United Kingdom WC1X 8DA

Sponsor information

Organisation

Department of Health (UK)

Funder(s)

Funder type

Government

Funder Name

The Royal Free Hampstead NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration