# Screening for health problems and emotional distress in GP clinics

Submission date	Recruitment status	<ul><li>Prospectively registered</li></ul>		
28/09/2007	No longer recruiting	☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
28/09/2007	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
13/05/2015	Mental and Behavioural Disorders			

#### Plain English summary of protocol

Background and study aims

GPs are encouraged to screen people for health and emotional problems. However, we know little of how people feel about being screened and the impact of screening on their health and well-being.

In order to find this out, we would like to conduct two short interviews with people over the telephone.

#### Who can participate?

Patients aged 18 and over at the participating 30 GP practices.

#### What does the study involve?

All participants will be randomly allocated to one of two groups. The participants in one group will be asked screening questions related to health and emotional problems whilst those in the other group will be asked similar questions at the follow-up interview in 7-10 days. The order in which people will be asked questions will be determined in advance at random (by chance). Selecting the order in which people are asked questions at random provides the best possible way for us to find out what, if any, effect this has.

#### What are the possible benefits and risks of participating?

There is no intended benefit for those people who agree to take part in the study. The main disadvantage of taking part would be the inconvenience of two telephone interviews. These interviews would last no more than 15 minutes and would be held at a time of your choosing. We do not have any evidence that the screening questions you will be asked would cause you any discomfort or distress. However, if the questions cause distress then let us know. A support mechanism is in place.

Where is the study run from? Imperial College London (UK).

When is the study starting and how long is it expected to run for? From July 2006 to February 2009.

Who is funding the study? Brent Primary Care Trust (UK).

Who is the main contact? Dr Priya Bajaj priya.bajaj@nhs.net

# Contact information

#### Type(s)

Scientific

#### Contact name

Dr Priya Bajaj

#### Contact details

Department of Psychological Medicine Claybrook Centre 37 Claybrook Road London United Kingdom W6 8LN +44 07971767269 priya.bajaj@nhs.net

## Additional identifiers

#### Protocol serial number

N0179192264

# Study information

#### Scientific Title

Screening for suicidal ideation: a randomised trial

#### Study objectives

What are the views of GPs, other primary care workers and patients to screening for suicidal ideation and behaviour?

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Not provided at time of registration

#### Study design

Randomised controlled trial

#### Primary study design

#### Interventional

#### Study type(s)

Screening

#### Health condition(s) or problem(s) studied

Mental and Behavioural Disorders: Distress

#### Interventions

The study involves a multi-method evaluation incorporating three linked components:

A. A survey of 100 GP and other primary care workers working in North and West London

B. A randomised trial examining the impact of screening for suicidal ideation on the subsequent incidence of suicidal ideation

C. A survey of the first 100 patients who participate in the trial

#### A. Survey of GPs and Primary Care workers

The lead GP in all GP surgeries in Brent and Westminster will be contacted and invited to help with the study. All GPs and primary care nurses working in selected GP surgeries will be invited to complete a short questionnaire in which they are asked their views about screening for suicidal ideation and behaviour among their patients. Arrangements will then be made to recruit up to 15 patients attending each of these surgeries

#### B. Randomised controlled trial

Patients attending the surgery with non-urgent medical problems will be approached by the GP during the consultation and be given written and verbal information about the study. Patients who agree to be contacted by a member of the research team will be provided with a copy of the Patient Information Sheet, and asked to sign a consent form and a short screening questionnaire to identify whether they have signs of depression (using the screening tool devised by Arroll et al 2003) and provide the researcher with details of how and when they can be contacted. They will also be given a sealed envelope in which they can place the completed questionnaire and the consent form. The sealed envelopes will be collected from the surgeries. Among them, all patients who describe feeling down, depressed or hopeless during the last four weeks or state that they have been bothered by little interest or pleasure in doing things, will be contacted by telephone within one week and asked to consider participating in the study.

Patients who answer no to both screening questions will be contacted to thank them for their help with the study and told that we do not need to ask them any further questions because they do not appear to have symptoms of mental distress. Patients who answered yes to one or both screening questions will be reminded about the aims of the study and asked to confirm their willingness to participate in the study.

Those willing to participate will be asked to complete a short interview over the telephone. The interview will comprise a measure of mental health - the twelve-item General Health Questionnaire (GHQ-12, Goldberg, 1992) followed by either six questions screening either suicidal ideation and behavior (early screening group), or diet and lifestyle (delayed screening group).

#### Randomisation

Each time a patient agrees to have a baseline interview the researcher will open a pre-prepared sealed opaque envelope that will contain either:

1. Early screening: Six questions on suicidal ideation and behaviour which have been widely used

in previous studies of suicidal behaviour among the general public (Meltzer, et al, 2002; Department of Health, 2002; Jenkins et al, 2003). The questions asked are:

- 1.1 In the last few weeks, have you felt that life was not worth living?
- 1.2 Have you wished that you were dead for instance, that you would go to sleep and not wake up?
- 1.3 Have you thought of taking your life even if you would not really do it?
- 1.4 Have you seriously considered taking your life?
- 1.5 Have you attempted to take your life?
- 1.6 Has any member of your family or someone else you are close to attempted to harm themselves or committed suicide in the past?
- 2. Delayed screening: Six questions regarding diet and lifestyle (taken from the Preventive Nutrition Project, 2004).

Allocation to questions on suicidal ideation and behaviour at baseline (early screening) and diet and lifestyle at baseline (delayed screening) will be made by simple randomisation using randomisation lists derived from random number tables which are stored on a secure computer. The randomisation ratio for early and delayed screening arms of the trial will be 1 to 1.

#### Follow-up procedures

Seven to 10 days after completion of the baseline interviews all patients will be contacted by telephone and asked to complete a short (15 minute) interview. The interview will comprise:

- 1. Screening questions for depression (Arroll et al 2003)
- 2. Screening questions for suicidal ideation and behaviour
- 3. An 8-item screening tool to assess personality (SAP-AS, Moran et al, 2003)
- 4. Questions on level of social support (Stansfeld & Marmot, 1992)
- C. Survey of the first 100 patients who participate in the trial

The first 100 patients to be included in the study will also be asked a short series of open-ended questions regarding their feelings and attitudes to being screened for suicidal ideation and behaviour.

#### References

- 1. Arroll B, Khin N, Kerse N (2003) Screening for depression in primary care with two verbally asked questions: cross-sectional study. British Medical Journal, 327, 1144-1146
- 2. Department of Health (2002) Ethnic Minority Psychiatric Illness Rates in the Community (EMPIRIC). (Eds.). London: The Stationery Office
- 3. Goldberg D (1992) General Health Questionnaire (GHQ-12). Windsor, UK, NFER-Nelson 1992
- 4. Jenkins R, Bebbington P, Brugha T, Farrell M, Gill B, Lewis G, Meltzer H & Petticrew M (2003) The National Psychiatric Morbidity Surveys of Great Britain-strategy and methods. International Review of Psychiatry, 15 (1-2), February-May 2003, 5-13
- 5. Meltzer, H., Lader, D., Corbin, T., Singleton, N., Jenkins, R., & Brugha, T. (2002) Non-fatal suicidal behaviour among adults aged 16 to 74 in Great Britain, 1985-86. The Stationery Office: London,
- 6. Moran P, Leese M, Lee T, Walters P, Thornicroft G & Mann A (2003) Standardised Assessment of Personality- Abbreviated Scale (SAPAS): preliminary validation of a brief screen for personality disorder. British Journal of Psychiatry, 183, 228-232
- 7. Preventive Nutrition Project (2004) Department of Family and Preventive Medicine, University of Arkansas For Medical Sciences, 4301 W. Markham St., Little Rock, AR 72205
- 8. Stansfeld, S.A. & Marmot, M.G. (1992) Deriving a survey measure of social support: the reliability and validity of the Close Persons Questionnaire. Social Science and Medicine 35, 1027-1035.

#### Intervention Type

Other

#### **Phase**

**Not Specified** 

#### Primary outcome(s)

Level of thinking life is not worth living

#### Key secondary outcome(s))

- 1. Attitudes of GPs and other primary care workers to screening for suicidal ideation and factors that promote and hinder screening in primary care
- 2. Patients' attitudes about screening for suicidal ideation and behaviour
- 3. Impact of screening for depression and suicidal ideation on the incidence of feelings that life is not worth living

#### Completion date

01/02/2009

# **Eligibility**

#### Key inclusion criteria

A. Inclusion criteria (for the randomised trial and patient survey)

- 1. Aged 18 years and above
- 2. Willingness to provide written informed consent to participate in the study
- 3. Have a contact telephone number where they can be contacted for a baseline and follow-up interview 7–10 days after the baseline interview

B. Inclusion Criteria (for the survey among GP and primary care workers): all GP and primary care workers in primary care clinics in north and west London.

## Participant type(s)

Patient

## Healthy volunteers allowed

No

### Age group

Adult

#### Lower age limit

18 years

#### Sex

All

#### Key exclusion criteria

Exclusion criteria (for the randomised trial and patient survey):

1. Those who decline to provide consent to participate in the study

- 2. All those presenting to the surgery for emergency medical problems
- 3. All those who have insufficient command of English to complete the baseline interview. As one of the researchers is fluent in Hindi, Urdu, Punjabi and Marathi, all patients who can only speak these languages will also be included.
- B. Exclusion criteria (for the survey among GP and primary care workers): all those who decline to participate

**Date of first enrolment** 01/07/2006

Date of final enrolment 01/02/2009

## Locations

**Countries of recruitment** United Kingdom

England

Study participating centre Claybrook Centre London United Kingdom W6 8LN

# Sponsor information

#### Organisation

Record Provided by the NHSTCT Register - 2007 Update - Department of Health

# Funder(s)

## Funder type

Government

#### Funder Name

Brent Primary Care Trust (UK)

# **Results and Publications**

# Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Not provided at time of registration

# **Study outputs**

Output type	Details	Date created Date added	d Peer reviewed?	Patient-facing?
Results article	results	01/05/2011	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	5 No	Yes