ISRCTN85118989 https://doi.org/10.1186/ISRCTN85118989

To evaluate whether a three days course of high doses of amoxicillin in the treatment of pneumonia in children is better compared to standard treatment with co-trimoxazole for five days

Submission date 19/05/2010	Recruitment status No longer recruiting	 Prospectively registered Protocol
Registration date 03/06/2010	Overall study status Completed	Statistical analysis planResults
Last Edited 03/06/2010	Condition category Respiratory	Individual participant dataRecord updated in last year

Plain English summary of protocol Not provided at time of registration

Contact information

Type(s) Scientific

Contact name Dr Syed Zaman

Contact details Health Protection Agency Centre for Infections 61 Colindale Avenue London United Kingdom NW9 5EQ

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Study information

Scientific Title

Efficacy of short course high-dose amoxicillin in the treatment of non-severe community acquired-pneumonia in children: A double-blind, randomised controlled trial

Acronym

HAT

Study objectives

We hypothesised that the treatment failure in community acquired non-severe pneumonia (defined by WHO using respiratory rates) or in community acquired non-severe radiological pneumonia (defined by WHO Radiology Working Group) will be lower in children randomised to twice daily three-day oral amoxicillin 90 mg/kg-per-day (high-dose amoxicillin) compared to five-day standard dose co-trimoxazole (standard therapy).

We also hypothesised that carriage of non-susceptible pneumococci to co-trimoxazole will be lower in children treated with high-dose amoxicillin compared to standard therapy.

Ethics approval required

Old ethics approval format

Ethics approval(s)

The Gambia Government / MRC Laboratories Joint Ethics Committee approved on the 23rd of June 2003

Study design Two arm randomized double blind single centre clinical trial

Primary study design

Interventional

Secondary study design Randomised controlled trial

Study setting(s) Hospital

Study type(s) Treatment

Participant information sheet

Not available in web format, please use contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Community acquired pneumonia in children

Interventions

Children randomised to high-dose amoxicillin received amoxicillin in 45 mg/kg/dose twice daily (maximum daily dose 2000 mg/day) for three days, followed by placebo twice daily for two days. Children randomised to co-trimoxazole received trimethoprim in 4 mg/kg/dose plus sulphamethoxazole in 20 mg/kg/dose twice daily for 5 days.

Intervention Type

Other

Phase Not Specified

Primary outcome measure

1. Treatment failure:

1.1. 3 days after enrolment there was no improvement

ог

1.2. within 5 days after enrolment, the study drug was changed to another antibiotic, severe pneumonia or very severe disease develops, or death occurs.

All children were assessed by nurses on days 3 and 5 post-enrolment. Treatment failures were confirmed by study pediatricians.

Secondary outcome measures

1. Relapse:

Reappearance of signs of non-severe pneumonia or appearance of signs of severe pneumonia or very severe disease by day-14 post-enrolment after being declared as cured on day-5 post-enrollment. All children were assessed by nurses on days 5, 14 and 28 post-enrolment for evaluation of clinical outcomes.

2. Compliance:

Proportion of children given full prescribed dosage (this was assessed by measuring left-over trial antimicrobials in the bottles on days 3 and 5).

3. Carriage rate of co-trimoxazole non-susceptible pneumococci on day-28 post-enrollment. Nasopharyngeal swab (NPS) for culture and sensitivity to antimicrobials was collected on days 0 and 28 of enrolment.

Overall study start date

05/03/2004

Completion date 02/06/2006

Eligibility

Key inclusion criteria

1. Aged 2 to 59 months

2. Either sex

3. Nutritional status: Weight-for-height > 70% of National Center for Health Statistics (NCHS) reference without oedema

4. Non-severe pneumonia according to WHO definition: if the child has fast breathing with cough or difficult breathing and there is no chest indrawing or other danger signs

Participant type(s)

Patient

Age group

Neonate

Sex

Both

Target number of participants

For community acquired pneumonia, the target number of participants was 850 children in each arm. For radiological pneumonia, the target number of participants was 55 children in each arm.

Key exclusion criteria

- 1. Having severe pneumonia or very sever disease or if needs oxygen
- 2. Needed antibiotic, steroid, theophylline or digitalis for treatment of any other condition
- 3. Had been enrolled in the trial for an earlier episode of pneumonia
- 4. Was admitted in a hospital in the previous month
- 5. History of hypersensitivity or intolerance to amoxicillin or co-trimoxazole

6. History of receiving any antibiotic within last 48 hours, this was be confirmed from health cards or village health workers

- 7. A history of three or more episodes of wheeze, acute bronchial asthma
- 8. Evidence of underlying haematologic, renal, hepatic or cardiovascular disease
- 9. Chronic steroid use or concomitant treatment with theophylline or digitalis glycosides
- 10. Living outside the study area

Date of first enrolment

05/03/2004

Date of final enrolment

02/06/2006

Locations

Countries of recruitment England

Gambia

United Kingdom

Study participating centre Health Protection Agency Centre for Infections London United Kingdom NW9 5EQ

Sponsor information

Organisation

Medical Research Council Laboratories (Gambia)

Sponsor details

PO Box 273 Banjul Atlantic Road Fajara Banjul Gambia 273

Sponsor type Research council

ROR https://ror.org/025wfj672

Funder(s)

Funder type Research council

Funder Name Medical Research Council Laboratories (Gambia)

Results and Publications

Publication and dissemination plan Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary Not provided at time of registration