

Blood loss and knee function after unicompartmental knee replacement or total knee replacement surgery

Submission date 06/04/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 06/04/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 06/04/2020	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Knee replacement, also known as knee arthroplasty, is a surgical procedure to replace the weight-bearing surfaces of the knee joint to relieve pain and disability. It is most commonly performed for osteoarthritis, and also for other knee diseases such as rheumatoid arthritis and psoriatic arthritis.

Unicompartmental knee arthroplasty (UKA) is a surgical procedure used to relieve arthritis in one of the knee compartments in which the damaged parts of the knee are replaced. UKA surgery may reduce post-operative pain and have a shorter recovery period than a total knee arthroplasty (TKA) procedure.

Utilization of tourniquet is considered to compromise the outcome of knee arthroplasty. This study aims to evaluate the hidden blood loss and function restoration of UKA without tourniquet by comparing with total knee arthroplasty (TKA).

Who can participate?

Adults over 18 years, scheduled to undergo UKA or TKA

What does the study involve?

This study is a retrospective analysis of patient records looking at patients who underwent UKA or TKA without tourniquet.

What are the possible benefits and risks of participating?

None.

Where is the study run from?

Department of Orthopedic Surgery, The First People's Hospital of Huzhou, Huzhou (China)

When is the study starting and how long is it expected to run for?

August 2017 to December 2019

Who is funding the study?
Department of Orthopedic Surgery, the Second Affiliated Hospital, School of Medicine, Zhejiang University (China)

Who is the main contact?
Dr Zhanfeng Zhang
335980557@qq.com

Contact information

Type(s)
Public

Contact name
Dr Zhanfeng Zhang

ORCID ID
<https://orcid.org/0000-0002-1469-4475>

Contact details
Department of Orthopedic Surgery
The First People's Hospital of Huzhou
158 Guangchang Back Rd
Wuxing District
Huzhou
China
313000
+86 15905720901
335980557@163.com

Additional identifiers

Clinical Trials Information System (CTIS)
Nil known

ClinicalTrials.gov (NCT)
Nil known

Protocol serial number
2019035

Study information

Scientific Title
Blood loss and knee function after unicompartmental knee arthroplasty (UKA) without tourniquet

Acronym
BLAKFAUKAWT

Study objectives

1. UKA or TKA without tourniquet results in lower blood loss compared to with tourniquet.
2. UKA without tourniquet results in less HBL compared with TKA
3. UKA without tourniquet enjoys a better outcome than TKA

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 06/06/2019, Medical Ethics Committee of The First People's Hospital of Huzhou (The First People's Hospital of Huzhou, 158 Guangchang Back Rd, Wuxing District, Huzhou, 313000, China; +86 (0)572 2508930; hzyllwyh@163.com), ref: 2019035

Study design

Single-center retrospective controlled study

Primary study design

Observational

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Osteoarthritis

Interventions

In this retrospective study, patients were included from August 2017 to October 2018. Both the UKA group and the TKA group underwent procedure without the utilization of tourniquet during the whole process.

The gender, age, body mass index, American Society of Anesthesiologists score, Kellgren-Lawrence grade, perioperative Hb, and volume of hidden blood loss (HBL) were recorded and analysed.

Knee function was assessed at 3 months and 12 months after the procedure by using HSS score.

Intervention Type

Procedure/Surgery

Primary outcome(s)

1. Mean volume of HBL during the procedure (ml)
2. Knee function assessed at 3 months and 12 months after the procedure by HSS score

Key secondary outcome(s)

Hb level measured using blood test on the 2nd day, 4th day, 6th day, 8th day postoperatively

Completion date

30/12/2019

Eligibility

Key inclusion criteria

1. Primary UKA or TKA
2. Kellgren-Lawrence (KL) grade of medial knee osteoarthritis grade

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

112

Key exclusion criteria

1. Preoperative abnormality in coagulation function, hematomosis history
2. Postoperative poor general situation
3. More than 2000ml per day in fluid infusion
4. Simultaneous bilateral UKA or TKA, and UKA or TKA secondary to a failed arthroplasty procedure

Date of first enrolment

01/08/2017

Date of final enrolment

30/10/2018

Locations**Countries of recruitment**

China

Study participating centre**Department of Orthopedic Surgery**

The First People's Hospital of Huzhou

158 Guangchang Back Rd

Wuxing District

Huzhou

China

313000

Sponsor information

Organisation

The First People's Hospital of Huzhou

Funder(s)**Funder type**

Hospital/treatment centre

Funder Name

Zhejiang University School of Medicine, Second affiliated hospital

Results and Publications**Individual participant data (IPD) sharing plan**

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

IPD sharing plan summary

Available on request