

Patient satisfaction in the treatment of anal fissure

Submission date 27/01/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 27/01/2006	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 14/08/2009	Condition category Digestive System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr W.F. Tets

Contact details
Sint Lucas Andreas Ziekenhuis
Department Surgery
P.O. Box 9243
Amsterdam
Netherlands
1006 AE
w.vantets@slaz.nl

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
NTR441; 2004/149

Study information

Scientific Title

Study objectives

Treatment of anal fissure with isosorbide dinitrate (ISDN) or Botulinum toxin A gives approximately comparative outcomes. Because of intensive treatment with isosorbide dinitrate cream, patients treated with Botulinum toxin A will be more satisfied. After six weeks treatment with Botulinum toxin A, the patient satisfaction will be significant more than patients treated with isosorbide dinitrate.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Received from local medical ethics committee

Study design

Multicentre randomised single blind active controlled parallel trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Anal fissure

Interventions

ISDN 1% cream versus Botuline Toxine A for a duration of 12 weeks.

ISDN cream application every 4 hours for 12 weeks.

Botulinum Toxin A injection at week 0 and if necessary again at week 6.

Botulinum Toxin A Dysport® is made by Ipsen.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Botulinum toxin A (Dysport®), isosorbide dinitrate

Primary outcome measure

1. Patient satisfaction after 6 weeks
2. Visual Analogue Score on weeks 0, 6 and 12

Secondary outcome measures

1. Patient satisfaction after 12 weeks
2. Pain after defecation at night
3. Incontinence after 6 and 12 weeks
4. Healing of fissure after 6 and 12 weeks

Overall study start date

08/11/2004

Completion date

01/07/2007

Eligibility**Key inclusion criteria**

1. Diagnosed anal fissure with complaints
2. Complaints longer than 2 months
3. Age 21-60 years
4. Dutch speaking
5. Will-competent
6. Informed consent

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

50

Key exclusion criteria

1. Pregnancy, lactation
2. Muscle-sicknesses such as myasthenia gravis
3. Simultaneous use of medication interacting with neuromuscular transmission
4. Fistulas
5. Coagulation disorders or the use of anticoagulants
6. Anal surgery in the past
7. Haemorrhoids or inflammatory bowel sicknesses as a cause of anal fissure
8. Major secondary changes because of the anal fissure

Date of first enrolment

08/11/2004

Date of final enrolment

01/07/2007

Locations

Countries of recruitment

Netherlands

Study participating centre

Sint Lucas Andreas Ziekenhuis

Amsterdam

Netherlands

1006 AE

Sponsor information

Organisation

Ipsen Farmaceutica B.V. (Netherlands)

Sponsor details

Hoofdweg Oostzijde 620

Hoofddorp

Netherlands

2132 MJ

Sponsor type

Not defined

Website

<http://www.ipsen.com>

ROR

<https://ror.org/026jmga48>

Funder(s)

Funder type

Industry

Funder Name

Ipsen Farmaceutica B.V. (Netherlands)

Results and Publications**Publication and dissemination plan**

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration