

Should the wound be closed with absorbable or non-absorbable sutures after surgery for Dupuytren's Contracture

Submission date 30/09/2005	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 30/09/2005	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 25/08/2009	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Ms Kim Howard

Contact details
Fracture and Orthopedic Clinic
Wirral Hospital NHS Trust
Arrowe Park Road
Upton
Wirral
United Kingdom
CH49 5PE

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N0280141212

Study information

Scientific Title

Study objectives

Would surgical treatment of Dupuytren's Contracture involves less discomfort for patients if absorbable (dissolving) sutures were used instead of non-absorbable sutures? Would absorbable sutures significantly reduce the amount of time nursing staff need to allocate to the care of patients having surgery for Dupuytren's Contracture?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Other

Participant information sheet

Health condition(s) or problem(s) studied

Musculoskeletal Diseases: Dupuytren's disease

Interventions

Randomised controlled trial:

1. absorbable-vicryl rapide 5/0 sutures
2. non-absorbable-prolene 5/0 sutures

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Patients discomfort/pain will be measured using visual analogue and non-parametric Mann Whitney test. The difference in the mean time the nurses spend with the patients in the two groups will be investigated using an independent t-test. Healing complications will probably take form of grouped data and comparisons of the two groups will be done by chi-squared test.

Secondary outcome measures

Not provided at time of registration

Overall study start date

01/05/2004

Completion date

31/05/2005

Eligibility

Key inclusion criteria

All patients that are listed for fasciectomy for Dupuytren's Contracture.

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

Minimum of 30 in each group.

Key exclusion criteria

1. Fasciectomy with skin grafting
2. Fasciectomy for recurrent diseases at the site of previous surgery

Date of first enrolment

01/05/2004

Date of final enrolment

31/05/2005

Locations

Countries of recruitment

England

United Kingdom

Study participating centre
Fracture and Orthopedic Clinic
Wirral
United Kingdom
CH49 5PE

Sponsor information

Organisation
Department of Health

Sponsor details
Richmond House
79 Whitehall
London
United Kingdom
SW1A 2NL
+44 (0)20 7307 2622
dhmail@doh.gsi.org.uk

Sponsor type
Government

Website
<http://www.dh.gov.uk/Home/fs/en>

Funder(s)

Funder type
Government

Funder Name
Wirral Hospitals NHS Trust (UK)

Funder Name
NHS R&D Support Funding

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2009		Yes	No