

Effect of postoperative nasal ventilation in patients undergoing coronary artery bypass grafting (CABG)

Submission date 06/04/2011	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 05/07/2011	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 11/11/2013	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Coronary artery bypass graft (CABG) surgery is well known to affect the lungs in a number of ways, mainly as a result of lung basal atelectasis (alveoli collapsing at the bottom of the lung), which is usually seen in the first 48 hours after an operation. Patients can end up staying in hospital longer as a result of this additional problem. We have carried out a study using a non-invasive technique (non-invasive ventilation [NIV] with bilevel positive airway pressure [BiPAP]) in patients undergoing CABG with cardiopulmonary bypass, in order to compare the results of using NIV with BiPAP with using routine standard care.

Who can participate?

Participants were patients referred for "on-pump" first time CABG surgery with mild hypothermia (body temperature between 32 and 35°C).

What does the study involve?

Patients were placed at random into one of two groups: either NIV using BiPAP or routine standard care following the operation. Lung performance was measured by levels of carbon dioxide gas (PCO₂) and a test of lung volume (FEV₁).

What are the possible benefits and risks of participating?

Participants in the NIV with BiPAP group may have a shorter time in hospital with fewer complications. We do not anticipate additional benefits for the routine care group than normally expected. We do not anticipate any side effects from the NIV with BiPAP treatment.

Where is the study run from?

Hammersmith Hospital (London, UK).

When is study starting and how long is it expected to run for?

The study will run from February 2008 to February 2011.

Who is funding the study?
Imperial College Healthcare NHS trust and Respiroics (now Phillips).

Who is the main contact?
Dr Philip Ind
p.ind@imperial.ac.uk

Contact information

Type(s)
Scientific

Contact name
Dr Philip Ind

Contact details
Respiratory Medicine
Imperial College NHS Trust
Hammersmith House
Du Cane Road
London
United Kingdom
W12 0HS
+44 (0)20 8383 3269
p.ind@imperial.ac.uk

Additional identifiers

Protocol serial number
INDP1073

Study information

Scientific Title
Effect on speed of recovery of postoperative nasal ventilation compared to conventional management in patients undergoing coronary artery bypass grafting (CABG): a randomised controlled trial

Study objectives

1. That following CABG surgery post-operative noninvasive ventilation (NIV) support with bilevel positive airway pressure (BiPAP) in all patients may significantly improve respiratory physiology.
2. That this will reduce complications, intensive care unit (ICU) admissions and duration, and length of hospital admissions when compared to conventional management.

Ethics approval required
Old ethics approval format

Ethics approval(s)

Hammersmith and Queen Charlotte's & Chelsea Research Ethics Committee approved on 04 September 2007, reference number: 07/Q0406/79

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Screening

Health condition(s) or problem(s) studied

Post-operative care of coronary artery bypass graft (CABG) patients

Interventions

Nasal bilevel positive airway pressure (BiPAP) support added to routine post-operative care compared to routine post-operative care

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Hospital length of stay from day of operation to day of fitness-to-discharge-until day when judged fit-for-discharge on the basis of clinical criteria

Key secondary outcome(s)

1. Lung function and blood gases-Lung function:

1.1. Pre-op lung function is done within 4 weeks of the operation (bed-side spirometry, and then formal departmental spirometry). In patients with a significant obstructive defect {defined as an FEV1 < 70% or [forced expiratory volume in one second (FEV1)/forced vital capacity (FVC)] of < 70%}, spirometry is then done before and after bronchodilator administration as well as body plethysmography for lung volumes and gas transfer.

1.2. Respiratory muscle strength, measured as mouth pressures [maximum inspiratory pressure (MIP) and maximum expiratory pressure (MEP)] have been measured on majority of patients.

1.3. Post-operative lung function is measured on day 1, 2 and 3 by bed-side spirometry with documentation of oxygen (O₂) saturation and fraction of inspired oxygen (FiO₂) given to patient.

1.4. Arterial Blood gases:

Are measured preoperatively as a baseline, just before the operation. ABG are measured after the operation at the time of ex-tubation, 1 hour later, 12 hours later and 24 hours from the time of ex-tubation. The Fio₂ is documented for each measurement

2. Intensive treatment unit (ITU) length of stay

3. Occurrence of complications

Completion date

31/03/2011

Eligibility

Key inclusion criteria

Any patient accepted for elective CABG surgery

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Pre-existing significant renal (creatinine > 200 umol/L), hepatic or haematological disease
2. Poor left ventricular function
3. Chronic infection
4. Emergency surgery

Date of first enrolment

01/11/2007

Date of final enrolment

31/03/2011

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Respiratory Medicine

London

United Kingdom

W12 0HS

Sponsor information

Organisation

R&D Hammersmith Hospital (UK)

ROR

<https://ror.org/05jg8yp15>

Funder(s)

Funder type

Industry

Funder Name

Imperial College Healthcare NHS trust (UK)- internal funding

Funder Name

Respironics (now Phillips) (UK)- supplied nasal ventilators (BiPAP), consumables and funded staff training

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2013		Yes	No