

# A comparison study of open surgery, minimal invasive surgery and medical management for complex aortic aneurysms

<b>Submission date</b> 20/12/2016	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 27/02/2017	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 24/06/2021	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

An abdominal aortic aneurysm (AAA) is a common condition where the aorta (the main artery that begins at the heart and travels down the torso) begins to bulge and expand at a portion just below the chest, above the level of the navel. In the usual course of events, the bulge expands slowly over many years and can eventually burst (rupture). This results in major internal blood loss often resulting in death and when an emergency lifesaving operation is possible, they have high failure rate. Although the risk of aneurysm rupture is generally low, it starts to increase if the aneurysm exceeds a certain size. A planned operation can be performed to repair the bulging section of aorta to prevent a burst aneurysm. However, not all abdominal aortic aneurysms are equal and technical aspects of aneurysm repair depend upon their location in relation to important branch arteries of the aorta. Over a quarter of aneurysms are classed as juxtarenal aneurysms because they are too close to the arterial branches supplying blood to the kidneys. These aneurysms require more extensive and more complex operations. There are different ways of managing such complex aneurysms and doctors are unsure which method is better. If a patient undergoes surgery, it is likely to be either open repair (where surgery takes place through an opening in the abdomen) or using endovascular techniques (key hole surgery where access to the aorta is gained through another artery, usually in the leg). Endovascular repair usually takes the form of fenestrated endovascular aneurysm repair (FEVAR), the usual method used or standard endovascular repair (Offlabel EVAR), a less complex technique that is originally intended for use in less complex aneurysms. Endovascular techniques have the advantage of a quicker recovery time and a better chance of survival. The aim of this study is to examine how these different treatment methods compare in terms of clinical benefit and utilisation of valuable NHS resources.

### Who can participate?

All patients undergoing juxtarenal aneurysm treatment in England

### What does the study involve?

Patients who are undergoing open repair, FEVAR, Offlabel EVAR undergo their surgery as planned. They are then monitored by the research team to look at how many have survived after

30 days. Those patients who have an AAA which is not very big are treated without surgery and are also followed up, with scans to see if their aneurysm has grown every six months. Some patients are also asked to complete questionnaires about their quality of life four times over the course of a year.

What are the possible benefits and risks of participating?  
There are no notable risks involved with participating.

Where is the study run from?  
Royal Liverpool University Hospital (UK)

When is the study starting and how long is it expected to run for?  
November 2015 to June 2024

Who is funding the study?  
National Institute for Health Research, Health Technology Assessment Programme (UK)

Who is the main contact?  
Miss Charlotte Rawcliffe  
C.Rawcliffe@liverpool.ac.uk

## Contact information

**Type(s)**  
Public

**Contact name**  
Miss Charlotte Rawcliffe

**Contact details**  
Cancer Research UK Liverpool Cancer Trials Unit  
University of Liverpool  
1st floor Block C, Waterhouse Building  
3 Brownlow Street  
Liverpool  
United Kingdom  
L69 3GL  
+44 1517 948167  
C.Rawcliffe@liverpool.ac.uk

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**  
LCTU202

# Study information

## Scientific Title

A Risk-adjusted and Anatomically Stratified Cohort Comparison Study of Open Surgery, Endovascular Techniques and Medical Management for Juxtarenal Aortic Aneurysms - The UK COMplex Aneurysm Study

## Acronym

UK-COMPASS

## Study objectives

Study Objectives:

1. To compare different treatment strategies for their perioperative mortality and morbidity, corrected for confounding physiological and anatomical characteristics, in order to account for baseline risk and indication biases
2. To identify whether particular physiological and/ or anatomical baseline characteristics are associated with better clinical outcomes or better health economic efficiency using one or other treatment strategy
3. To compare different treatment strategies in terms of overall survival and in terms of treatment failure in the long-term follow-up (stent-graft related complications, secondary interventions, aneurysm-related mortality)
4. To perform cost effectiveness analyses from a broad societal perspective (NHS as well as non-NHS costs) to establish incremental cost effectiveness ratios, comparing different treatment strategies in terms of life years and quality adjusted life years gained
5. To establish the clinical and cost utility of fEVAR and of off-label standard EVAR, in patients who are considered physiologically unfit for OSR, and to compare these against medical management

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Not provided at time of registration

## Study design

Observational cohort study

## Primary study design

Observational

## Secondary study design

Cohort study

## Study setting(s)

Hospital

## Study type(s)

Other

## Participant information sheet

No participant information sheet available

## **Health condition(s) or problem(s) studied**

Juxtarenal aortic aneurysms

## **Interventions**

This is an observational study of patients being treated within the NHS for juxtarenal aortic aneurysms. No changes have been proposed in the care of patients for the purposes of this study and with the correct approvals and safeguards pseudo- anonymised data and copies of scans will be collected on all patients treated in the NHS for juxtarenal aortic aneurysms.

Surgeons will advise their patients regarding what is considered the best treatment for them, forming cohorts of different treatments. Care pathways in each of the cohorts of this study:

**Open Surgical Repair (OSR):** Major surgery involving surgically replacing aneurysm with a synthetic graft, through laparotomy. Patients are reviewed to ensure sound healing of surgical wounds and are prepared for return to normal activities, but there is no clinical need for longer term follow-up

**Fenestrated Endovascular Aneurysm Repair (FEVAR):** Specially designed stent-grafts are placed in the aorta bridging the aorta above the level of the renal arteries to normal arterial segments below the aneurysm. The device comprises openings (fenestrations) to allow blood flow into visceral and renal branches of aorta. This technique avoids laparotomy, leading to quicker recovery and improved perioperative safety.

**Off-label EVAR:** Judicious use of a less advanced stent-graft designed to repair simpler infrarenal aneurysms (outside the conditions of manufacturers' Instructions for Use). This is termed 'off-label standard EVAR'. Patients undergoing FEVAR and off-label EVAR will require lifelong follow-up by means of a structured surveillance based on scans (CT / ultrasound) and plan X-rays. Clinical practice varies slightly, but usually imaging is done at 1 month and annually thereafter

**Medical management 'Operation-deferred':** In some patients with comorbidity and aneurysm larger than 55 mm, the risk of death from aneurysm may be considered to be smaller than the risk of death from an elective repair. In such situations, the surgical option may be deferred with a view to proceeding with surgical intervention if the perceived risk of rupture increases through further aneurysm enlargement. Such patients are medically managed with the aim of optimising fitness for surgery. Patients in this cohort undergo periodic ultrasound scans (typically every 6 m) to monitor the aneurysm size as well as best management of comorbidity. Please indicate the differences between the current standard care.

For the quality of life sub-study patients for the selected centres will be approached to take part in the Quality of Life Study. After consent patient will be asked to complete a series of 4 QoL questionnaires (EQ-5D 5SL, A-DQoL, ASRQ, A-TSQ) at 4 time points during their first year after their procedure.

## **Intervention Type**

Procedure/Surgery

## **Primary outcome measure**

Preoperative death, defined as a patient death by any cause occurring within 30 days of surgery, is measured by obtaining data on patients' deaths from the office of national statistics (ONS) and relating any death date to the date of surgery to detainee if death occurred within 30 days.

### **Secondary outcome measures**

1. Perioperative complications (morbidity) is measured using routinely collected data obtained through HES which will be collected annually throughout the course of the study
2. Long-term survival is measured using data on patient deaths obtained from the office of national statistics.
3. Secondary intervention in late follow-up is measured using obtained through HES which will be collected annually throughout the course of the study
4. Intensive care usage is measured using obtained through HES which will be collected annually throughout the course of the study
5. Length of hospital stay is measured using obtained through HES which will be collected annually throughout the course of the study
6. Treatment costs and health economic evaluations are completed using probabilistic modelling to generate estimates of the ICERs and compare the effect of the different structures of treatment provision on different sub-groups of patients
7. Quality of Life is measured using prospectively collected EQ5D forms which will be measured at registration and every 3 months following in the subgroup of X patients who are registered into this cohort of the study

### **Overall study start date**

03/11/2015

### **Completion date**

30/06/2024

## **Eligibility**

### **Key inclusion criteria**

1. Patients undergoing elective Juxtarenal AAA repair in England; juxtarenal AAA defined and stratified into 4 strata of anatomical complexity as specified in 'Anatomical and Physiological Stratification'
2. Patients with a Juxtarenal AAA 55 mm or larger in size and placed on Medical Management 'Operation-deferred'

### **Participant type(s)**

Patient

### **Age group**

Adult

### **Sex**

Both

### **Target number of participants**

2000 routine, 1100 site

## **Key exclusion criteria**

1. Surgeon-modified stent-grafts and devices
2. Emergency operations (elective operations are distinguished from non-elective ones on NVR)
3. Thoracic or thoracoabdominal aneurysms (this is a different extent/location of aneurysm disease, requiring different health technology. These treatments are currently being investigated by ETAA study-HTA 11/147/03. Patients will be included in ETAA),
4. Aneurysm neck anatomy suitable for standard infrarenal EVAR within IFU of any CE marked device
5. 'Operation Declined Medical Management' patients, due to co-morbidity or patient choice

## **Date of first enrolment**

01/07/2017

## **Date of final enrolment**

31/12/2022

## **Locations**

### **Countries of recruitment**

England

United Kingdom

### **Study participating centre**

**Royal Liverpool University Hospital**

Regional Vascular Unit, Link 8C

Prescot Street

United Kingdom

Liverpool

United Kingdom

L7 8XP

## **Sponsor information**

### **Organisation**

The Royal Liverpool and Broadgreen University Hospitals NHS Trust

### **Sponsor details**

RD&I

4th Floor Linda McCartney

Royal Liverpool Hospital

Prescot Street

Liverpool

England  
United Kingdom  
L7 8XP

**Sponsor type**

Hospital/treatment centre

**ROR**

<https://ror.org/009sa0g06>

## **Funder(s)**

**Funder type**

Government

**Funder Name**

Health Technology Assessment Programme

**Alternative Name(s)**

NIHR Health Technology Assessment Programme, HTA

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

United Kingdom

## **Results and Publications**

**Publication and dissemination plan**

1. Planned publication in peer-reviewed journals of high impact factor, presentations and lectures in scientific meetings, as well as releases to non-peer reviewed medical press
2. Planned presentation of findings at The Vascular Society's annual conference during special session in addition to dissemination through the British Society of Interventional Radiology, The British Society for Endovascular Therapy and The Society for Vascular Nursing, all representing professionals with an interest in the welfare of patients with juxtarenal AAA
3. Exploring potential opportunities for dissemination with The Circulation Foundation and the British Heart Foundation
4. Specialised services Clinical Reference Groups will be directly informed of the conclusions of the study after finalisation of reports

**Intention to publish date**

30/06/2023

### **Individual participant data (IPD) sharing plan**

The current data sharing plans for the current study are unknown and will be made available at a later date.

### **IPD sharing plan summary**

Data sharing statement to be made available at a later date