

# To compare the safety and efficacy of 'low dose' vaginal misoprostol and dinoprostone vaginal gel for induction of labour at term

<b>Submission date</b> 12/09/2003	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 12/09/2003	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 11/09/2013	<b>Condition category</b> Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Ms Sarah Gregson

### Contact details

Maternity Unit  
Queen Mary's Sidcup NHS Trust  
Frogna Avenue  
Sidcup, Kent  
United Kingdom  
DA14 6LT

## Additional identifiers

### Protocol serial number

N0533091848

## Study information

### Scientific Title

A single-blind randomised controlled trial comparing low dose vaginal misoprostol and dinoprostone vaginal gel for inducing labour at term

**Study objectives**

To determine if vaginal misoprostol is a better method of inducing labour at term than dinoprostone vaginal gel.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Not provided at time of registration

**Study design**

Randomised controlled trial

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Pregnancy and Childbirth: Induction of labour

**Interventions**

1. Misoprostol
2. Dinoprostone

**Intervention Type**

Other

**Phase**

Not Specified

**Primary outcome(s)**

1. Uterine tachysystole
2. Hyperstimulation
3. Presence of meconium in the amniotic fluid
4. Apgar Scores at 5 min. Umbilical arterial pH and base deficit
5. Neonatal Unit admission
6. Induction - delivery interval
7. Method of delivery
8. Bishop score at onset of labour
9. Oxytocin requirements in labour
10. Mode of delivery
11. Analgesia requirements in labour

**Key secondary outcome(s)**

Not provided at time of registration

**Completion date**

31/01/2004

# Eligibility

## Key inclusion criteria

1. Women at term (37-42 completed weeks of pregnancy)
2. Single fetus, cephalic presentation
3. Membranes may be intact or ruptured
4. Reactive fetal heart tracing

## Participant type(s)

Patient

## Healthy volunteers allowed

No

## Age group

Adult

## Sex

Female

## Key exclusion criteria

Does not match inclusion criteria

## Date of first enrolment

01/07/2000

## Date of final enrolment

31/01/2004

# Locations

## Countries of recruitment

United Kingdom

England

## Study participating centre

### Maternity Unit

Sidcup, Kent

United Kingdom

DA14 6LT

# Sponsor information

## Organisation

Department of Health (UK)

## Funder(s)

### Funder type

Hospital/treatment centre

### Funder Name

Queen Mary's Sidcup NHS Trust (UK)

## Results and Publications

### Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/04/2005		Yes	No