

# Dose-finding of oxantel pamoate in school-aged children infected with *Trichuris trichiura* on Pemba, United Republic of Tanzania

<b>Submission date</b> 08/10/2014	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 12/11/2014	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 22/09/2015	<b>Condition category</b> Infections and Infestations	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

*Trichuris trichiura* (whipworm) is a helminthic parasite, transmitted through soil. Whipworm infection is a neglected tropical disease and causes a huge burden worldwide. The available drugs have limitations. This study will compare the effectiveness of six different doses of a drug, oxantel pamoate, compared to a placebo (dummy) drug in school-aged children infected with *T. trichiura*, in order to find out the right dose of the drug.

### Who can participate?

Children studying in the participating primary schools.

### What does the study involve?

To examine whether children are infected, two stool samples will be collected and analysed. All children who are positive for *T. trichiura* will be randomly allocated to the seven different treatment groups and receive either one of six different doses of the drug or the placebo treatment. Children are interviewed to find out if they had any side effects. After three weeks, children will provide another two stool samples to see whether they got cured and to calculate egg reduction rates.

### What are the possible benefits and risks of participating?

At the end of the study, children remaining positive for the infection will be treated according to national guidelines. All the working procedures and examinations during this study do not bear any particular additional risks. Side effects are mild and include stomach ache and dizziness.

### Where is the study run from?

Two primary schools on Pemba Island, United Republic of Tanzania.

### When is the study starting and how long is it expected to run for?

October 2014 to December 2014.

Who is funding the study?  
Swiss National Science Foundation (Switzerland).

Who is the main contact?  
Prof Jennifer Keiser  
jennifer.keiser@unibas.ch

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Jennifer Keiser

**Contact details**  
Swiss Tropical and Public Health Institute  
Socinstr. 57  
Basel  
Switzerland  
4051

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**  
No

## Study information

**Scientific Title**  
Dose-finding of oxantel pamoate in school-aged children infected with *Trichuris trichiura* on Pemba, United Republic of Tanzania: a randomized controlled phase 2a trial

**Acronym**  
Oxpatri

**Study objectives**  
Different doses of oxantel pamoate have not yet been sufficiently studied in infected school-aged children. Dosing regimens in humans are empirical.

**Ethics approval required**  
Old ethics approval format

**Ethics approval(s)**

**Study design**

Randomized controlled phase 2a dose finding trial

**Primary study design**

Interventional

**Secondary study design**

Randomised controlled trial

**Study setting(s)**

Other

**Study type(s)**

Treatment

**Participant information sheet**

Not available in web format, please use the contact details below to request a patient information sheet

**Health condition(s) or problem(s) studied**

Trichuriasis

**Interventions**

Two stool samples will be collected on two consecutive days. The medical history of the participating schoolchildren will be assessed with a standardized and previously used questionnaire, in addition to a clinical examination carried out by the study physician on the treatment day. All children positive for *T. trichiura* will be randomly assigned (computer-generated stratified block randomization code) to one of the seven treatment arms: 5, 10, 15, 20, 25, 30 mg/kg oxantel pamoate and placebo. School-aged children will also be interviewed before treatment, 2 and 24 hours after treatment about the occurrence of adverse events. The efficacy of the treatment will be determined 21-26 days post-treatment by collecting another two stool samples. All stool samples will be examined with duplicated Kato-Katz thick smears.

1. 400 mg placebo
2. Oxantel pamoate 5 mg/kg
3. Oxantel pamoate 10 mg/kg
4. Oxantel pamoate 15 mg/kg
5. Oxantel pamoate 20 mg/kg
6. Oxantel pamoate 25 mg/kg
7. Oxantel pamoate 30 mg/kg

**Intervention Type**

Drug

**Phase**

Phase II

**Drug/device/biological/vaccine name(s)**

Oxantel pamoate

**Primary outcome measure**

1. Cure rate (CR) and egg reduction rate (ERR) against *T. trichiura*: 14-21 days post-treatment
2. Safety/adverse events: 3 hours and 24 hours post-treatment

**Secondary outcome measures**

CR and ERR against *A. lumbricoides* and hookworms

**Overall study start date**

16/10/2014

**Completion date**

29/11/2014

**Eligibility****Key inclusion criteria**

1. Written informed consent signed by parents and/or legal guardian; and oral assent by children
2. Able and willing to be examined by a study physician at the beginning of the study
3. Able and willing to provide two stool samples at the beginning (baseline) and approximately three weeks after treatment (follow-up)
4. Positive for *T. trichiura* eggs in the stool
5. Absence of major systemic illnesses (e.g. cancer, diabetes, clinical malaria or hepato-splenic schistosomiasis) as assessed by a medical doctor, upon initial clinical assessment
6. No known or reported history of chronic illness as cancer, diabetes, chronic heart, liver or renal disease
7. No recent anthelmintic treatment (within past 4 weeks)
8. No known allergy to study medications

**Participant type(s)**

Patient

**Age group**

Child

**Sex**

Both

**Target number of participants**

350

**Key exclusion criteria**

1. No written informed consent by parents and/or legal guardian
2. Presence of any abnormal medical condition, judged by the study physician
3. History of acute or severe chronic disease such as cancer, diabetes, chronic heart, liver or renal disease
4. Recent use of anthelmintic drug (within past 4 weeks)
5. Attending other clinical trials during the study
6. Negative diagnostic result for *T. trichiura* (absence of helminth eggs in stool)

**Date of first enrolment**

16/10/2014

**Date of final enrolment**

29/11/2014

## **Locations**

**Countries of recruitment**

Switzerland

Tanzania

**Study participating centre**

Swiss Tropical and Public Health Institute

Basel

Switzerland

4051

## **Sponsor information**

**Organisation**

Swiss National Science Foundation (SNSF) (Switzerland)

**Sponsor details**

Wildhainweg 3

Bern

Switzerland

3001

**Sponsor type**

Research organisation

**Website**

<http://www.snf.ch>

**ROR**

<https://ror.org/00yjd3n13>

## **Funder(s)**

**Funder type**

Research organisation

**Funder Name**

Swiss National Science Foundation (Switzerland)

**Alternative Name(s)**

Schweizerischer Nationalfonds, Swiss National Science Foundation, Fonds National Suisse de la Recherche Scientifique, Fondo Nazionale Svizzero per la Ricerca Scientifica, Fonds National Suisse, Fondo Nazionale Svizzero, Schweizerische Nationalfonds, SNF, SNSF, FNS

**Funding Body Type**

Private sector organisation

**Funding Body Subtype**

Trusts, charities, foundations (both public and private)

**Location**

Switzerland

## Results and Publications

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date****Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/01/2016		Yes	No