Effectiveness of cervical osteopathic manipulation techniques of high velocity and low amplitude in patients with whiplash-associated disorders

Submission date	Recruitment status No longer recruiting	Prospectively registered		
14/05/2014		☐ Protocol		
Registration date 21/07/2014	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited	Condition category	Individual participant data		
18/12/2017	Musculoskeletal Diseases			

Plain English summary of protocol

Background and study aims

Whiplash is a neck injury caused by a sudden movement forward, backward or sideways movement of the head. It often happens after a sudden impact from, for example, a road traffic accident and results in damage to the ligaments and tendons in the neck. Road traffic accidents and their effects on social-health costs are an important problem for society. Whiplash and whiplash associated disorders (WAD) for example pain, reduced range of neck movement, neurological damage and even fracture) are the number one reason for people visiting the emergency room after being involved in a road traffic accident. Osteopathic high velocity, low amplitude (HVLA) techniques, and mobilization techniques of the cervical spine, either alone or combined with other treatments, are methods used to treat neck problems. Both treatments result in similar short term changes, but there is not enough data to assess their long-term effects, making it difficult to develop the best technique and when and how long the treatments should be applied. The aim of this study is to precisely and objectively contribute to the knowledge available on how good HVLA techniques are at treating whiplash and WAD. HVLA techniques are compared with other physiotherapy techniques, looking at how long the treatments are needed, the degree of pain relief they offer, how much they improve the function and mobility of the spine and analysing radiological changes in cervical curvature.

Who can participate?

Patients aged 18 to 65 with WAD admitted to the Osteopathic Centre in Terrassa.

What does the study involve?

Participants are randomly allocated into one of two groups. Those in the study group receive HVLA treatment and the control group receive conventional manual osteopathic therapy and therapeutic exercises. The results from the two groups are then compared to assess how well they treat whiplash and WAD.

What are the possible benefits and risks of participating?

Participants are unlikely to directly benefit from the study, but the results could help future sufferers of whiplash, resulting in better management of their symptoms and avoiding excessive procedures. The risks associated with this study are rare and generally mild, but include some neurodegenerative reactions (sweating, low blood pressure, feeling sick and/or rapid heart rate) during, or immediately after, the therapy.

Where is the study run from? Health Consortium of Terrassa (Spain)

When is the study starting and how long is it expected to run for? May 2014 to December 2015

Who is funding the study?

The Osteopathic Centre, Terrassa and the Health Consortium of Terrassa (Consorci Sanitari de Terrassa) (Spain)

Who is the main contact? Mr Joan Parera Joanpt1@gmail.com

Contact information

Type(s)

Scientific

Contact name

Mr Joan Parera

ORCID ID

https://orcid.org/0000-0002-3969-2433

Contact details

Sant Marian 99 5-2 Terrassa Spain 08221

joanpt1@gmail.com

Additional identifiers

Protocol serial number N/A

Study information

Scientific Title

Research project on the effectiveness of cervical OSTEOpathic manipulation techniques of high velocity and low amplitude in patients with whiplash-associated disorders: a Randomized Controlled single-blind Trial

Acronym

OSTEO-RCT

Study objectives

Patients who have suffered cervical whiplash, specific osteopathic techniques to adjust the treatment, of high velocity and low amplitude (SAT-HVLA) of the cervical spine, obtain better results than those treated with combined treatments found in cervical therapeutic exercise manuals.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee at the Health Consortium of Terrassa - Terrassa Hospital (Consorci Sanitari de Terrassa Hospital de Terrassa), 20/12/2013

Study design

Randomized controlled single-blind trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Whiplash associated disorders (WAD)

Interventions

The participants are randomised to two groups:

1. Treatment: The technique of choice, will be the Specific Adjusting Technique (SAT). This is a High Velocity Low Amplitude Technique, whose objective is to reverse the injury caused by the trauma.

Throughout the trial we will make three interventions with SAT (one every two weeks)

In the first intervention, we will manipulate the upper cervical segment (C2/C3)

In the second intervention, we will manipulate the low cervical segment (C5/C6)

In the third intervention, we will manipulate the upper thoracic segment (T1/T2)

2. Control: The participants will receive conventional treatment, consisting in manual therapy techniques and therapeutic exercises, over four weeks.

Participants are followed up at 11 days, 30 days, and 110 days after treatment.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Current primary outcome measures as of 01/02/2017:

- 1. Functionality (disability), measured using the Northwick Park Neck Pain Questionnaire (NPQ) at baseline, 11 days, 30 days, and 110 days after treatment
- 2. Cervical mobility, measured using the Cervical Range Of Motion Instrument (CROM) at baseline, 11 days, 30 days, and 110 days after treatment
- 3. Radiological changes in the cervical curvature, measured using the Cobb method (COOB) at baseline and 30 days after treatment

Previous primary outcome measures:

- 1. Functionality, measured using the Northwick Park Neck Pain Questionnaire (NPQ)
- 2. Cervical mobility, measured using the Cervical Range Of Motion Instrument (CROM)
- 3. Radiological changes in the cervical curvature (CA)

Key secondary outcome(s))

Current secondary outcome measures as of 01/02/2017:

- 1. Pain, measured using the Visual Analogue Scale (VAS) at baseline, 11 days, 30 days, and 110 days after treatment
- 2. Anxiety, measured using the Hospital Anxiety and Depression Scale (HAD) at baseline, 30 days, and 110 days after treatment
- 3. Depression, measured using the Hospital Anxiety and Depression Scale (HAD) at baseline, 30 days, and 110 days after treatment

Previous secondary outcome measures:

- 1. Quality of life
- 2. Mobility of the cervical spine
- 3. Lordosis of the cervical spine

Completion date

31/12/2015

Eligibility

Key inclusion criteria

- 1. Patients ages 18 to 65 years presenting whiplash syndrome (WAD)
- 2. They must have radiological studies (A-P and lateral projections) with previous signed informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

Sex

All

Key exclusion criteria

- 1. Chronic cervical pain
- 2. Bone pathology
- 3. Systemic pathologies (acute infections, cancerous or inflammatory processes)

Date of first enrolment

22/05/2014

Date of final enrolment

01/07/2015

Locations

Countries of recruitment

Spain

Study participating centre

The Osteopathic Centre, Terrassa (Centre d'osteopatia Terrassa)

Spain

08224

Sponsor information

Organisation

The Osteopathic Centre, Terrassa (Centre d'osteopatia Terrassa) (Spain)

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

The Osteopathic Centre, Terrassa (Centre d'osteopatia Terrassa) (Spain)

Funder Name

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results		31/01/2017	15/02/2017	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes