

A mouthwash medicine for the reduction or prevention of mouth ulceration caused by radiotherapy for head and neck cancer

Submission date 25/10/2022	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 13/02/2023	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 11/03/2025	Condition category Oral Health	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

This study is looking at a new medication called NG11-2 to see if it can reduce a side effect called severe Radiation-induced Oral Mucositis (RIOM) that can happen to people getting radiation treatment for head and neck cancer. Severe RIOM can make patients very sick and make it hard for them to finish their cancer treatment. Right now, there are no approved drugs that can prevent or treat severe RIOM. This study will test different doses of NG11-2 to find the best one, and then more patients will take that dose to see if it works. This study is being done at multiple hospitals.

Who can participate?

Patients aged 18 years or older, with head and neck cancer.

What does the study involve?

Daily treatment with NG11-2 will occur prior to radiotherapy for 5 days/week for up to 7 weeks. Throughout the radiotherapy regimen, no more than 60 minutes prior to the completion of radiotherapy, on each weekday the patient will be instructed to

a) rinse their mouth twice with room temperature water in the 5 minutes prior to NG11-2 treatment then

b) rinse their mouth with 12.5ml of NG11-2 for between 90 & 120 seconds duration, gargling twice for approximately three seconds each if possible during this period then

c) expectorate the solution

Treatment will continue for up to 7 weeks with a 6-week follow-up period.

What are the possible benefits and risks of participating?

Benefits:

Possibly prevention/ reduction in severe Radiation-induced Oral Mucositis during the patient's radiotherapy treatment

Risks:

The study will fit in with the usual treatment and the only additional things that patients will have to do are: give a small number of extra blood samples, have some patches put on their

chest for an electrocardiogram (ECG), and fill out a short questionnaire a few times. Patients will also have to rinse their mouths with the NG11-2 solution before each radiation treatment. The study drug is made with adrenaline, which is a well-understood drug, but it may cause some mild side effects like irritation in the mouth or a temporary change in heart rate. The study team will closely monitor patients for any side effects and will stop treatment if needed. Although the scientific rationale and earlier laboratory studies suggest that NG11-2 treatment is unlikely to have a negative impact on your tumour treatment, we cannot yet exclude the risk of a negative impact on your anticancer therapy when treated with NG11-2.

Where is the study run from?
VasoDynamics Ltd (UK)

When is the study starting and how long is it expected to run for?
October 2022 to August 2024

Who is funding the study?
VasoDynamics Ltd (UK)

Who is the main contact?
Dr Mary Lei, Mary.Lei@gstt.nhs.uk

Study website
<https://vasodynamics.co.uk/ng11/>

Contact information

Type(s)
Public

Contact name
Mr Gary Bower

Contact details
Stevenage Biocatalyst Centre
Gunnels Wood Rd
Stevenage
United Kingdom
SG1 2FX
+44 1438 300880
gary.bower@vasodynamics.co.uk

Type(s)
Principal Investigator

Contact name
Dr Mary Lei

Contact details
Dept of Clinical Oncology
14th floor, Tower Wing
Guy's Hospital

Great Maze Pond
London
United Kingdom
SE1 9RT
+44 20 71884219
Mary.Lei@gstt.nhs.uk

Type(s)

Scientific

Contact name

Dr Ningfeng Fiona Li

Contact details

Stevenage Biocatalyst Centre
Gunnels Wood Rd
Stevenage
United Kingdom
SG1 2FX

-

fiona.li@vasodynamics.co.uk

Additional identifiers

EudraCT/CTIS number

2022-002409-99

IRAS number

1004528

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

NG11-2 P1b, IRAS 1004528, CPMS 54459

Study information

Scientific Title

A phase-1b dose escalation study to assess the effect of NG11-2 on radiation induced oral mucositis in patients with head & neck cancer

Acronym

NG11-2-P1b

Study objectives

The main objective of the trial is to see if the use of vasoconstrictors (such as Adrenaline), when administered as a mouthwash in head and neck cancer patients prior to their radiotherapy, is

tolerable and can reduce or prevent the development of severe radiation induced oral mucositis. More specifically, the trial aims to identify an appropriate dose of NG11-2 IMP based on safety and activity/preliminary efficacy for subsequent assessment in follow-up clinical trials.

Although the primary objective of the trial is to evaluate the safety & tolerability of NG11-2 mouthwash, the secondary objective is to obtain preliminary data on the activity/efficacy of NG11-2 mouthwash in the reduction of severe RIOM.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 23/01/2023, (West Midlands - Coventry & Warwickshire Research Ethics Committee, The Old Chapel, Royal Standard Place , Nottingham, NG1 6FS, UK; +44 207 104 8184; coventryandwarwick.rec@hra.nhs.uk), ref: 22/WM/0262

Study design

Interventional non-randomized dose escalation study

Primary study design

Interventional

Secondary study design

Non randomised study

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

See study outputs table

Health condition(s) or problem(s) studied

Reduction of radiation-induced oral mucositis in head & neck cancer patients undergoing radiotherapy with or without concomitant chemotherapy

Interventions

This is an unblinded, single-arm study. Patients will undergo daily treatment with NG11-2 immediately prior to radiotherapy for 5 days/week (Mon-Fri) for up to 7 weeks. The dose level will vary from 0.92 mg/mL to 5.5 mg/mL during the dose escalation process. The NG11-2 is given as an oral rinse.

Throughout the radiotherapy regimen, no more than 60 minutes prior to the completion of radiotherapy, on each weekday the patient will be instructed to a) rinse their mouth twice with room temperature water in the 5 minutes prior to NG11-2 treatment then b) rinse their mouth with 12.5 ml of NG11-2 for between 90 & 120 seconds duration, gargling twice for approximately three seconds each if possible during this period then c) expectorate the solution. Treatment will continue for up to 7 weeks with a 6-week follow-up period.

Intervention Type

Drug

Phase

Phase I

Drug/device/biological/vaccine name(s)

NG11-2 Mouthwash, 0.92 – 5.50 mg/mL [L-adrenaline base]

Primary outcome measure

Occurrence of Dose Limiting Toxicities (DLT) & Serious Adverse Events (SAE). Serious Adverse Events will be evaluated throughout the study from time of patient consent until 42 days after last IMP dose. Dose limiting toxicities will be evaluated from time of first dose until 42 days after last IMP dose. Additionally all available safety data (AEs, SAEs and DLTs) will be reviewed and evaluated by a Safety Committee (Sponsor CMO, Study Medical Monitor & Chief Investigator) before each dose escalation.

Secondary outcome measures

1. Duration of severe RIOM (WHO, RTOG and NCI-CTCAE v5 Gr 3 or above) evaluated by Oral Mucositis assessments at screening and twice weekly in each week of study treatment in weeks 1-7, and then at 7, 14, 28 and 42 days post end of Radiotherapy, in the follow-up period.
2. Incidence & Time to onset of severe RIOM (WHO, RTOG and NCI-CTCAE v5 Gr 3 or above)
3. Patient Reported Outcome Measure - Oral Mucositis (PROM-OM) evaluated at the same timepoints as the clinical Oral Mucositis assessments, namely screening and twice weekly in each week of study treatment in weeks 1-7, and then at 7, 14, 28 and 42 days post end of Radiotherapy, in the follow-up period.

Overall study start date

21/10/2022

Completion date

20/08/2024

Eligibility

Key inclusion criteria

1. Male or female head and neck cancer patients scheduled for radiotherapy, aged 18 or over on day of signing informed consent;
2. Histologically confirmed head and neck cancer with or without previous resective surgery, having not been previously treated with radiation therapy
3. Scheduled to receive a mean radiation dose of no less than 30 Gy on either (a) Extended Oral Cavity, or (b) Buccal Mucosa, or (c) Lips, according to CT-based delineation of Organ-At-Risk (OAR) Guideline, with or without concurrent chemotherapy (Section 13.8 for guidance);
4. Rendered dentally fit for radiotherapy;
5. Have a Performance Status >60 on the Karnofsky scale
6. No evidence of active systemic infections at the time of screening;
7. No oral fungal infection by visual examination or swab test;
8. Body Mass Index ≥ 18.5 kg/m²
9. Normalcy of diet evaluated with the Performance Status Scale for Head and Neck cancer patients (PSSHN) ≥ 30 ;
10. Ability to retain and swill liquid inside the oral cavity for at least 90 seconds;
11. Female patients of childbearing potential must have a negative urine or serum pregnancy

test during the 2 week Screening period

12. Provide written informed consent for the trial.

13. Consent to utilize medically acceptable methods of contraception throughout the study period if of child-bearing potential;

14. Be able to comply with protocol procedures (such as Oral Mucositis assessment and swish /spit) and study schedule

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

Up to 32 maximum - estimated 20-24

Total final enrolment

15

Key exclusion criteria

1. Tumour of the larynx
2. Mentally or legally incapacitated, in the opinion of the PI, which could interfere with the ability of the patient to understand or adhere to the requirements of the study;
3. Open or unhealed non-cancerous wounds or ulcers in the oral cavity;
4. WHO Oral Mucositis grade of 2 or more
5. Uncontrolled Hypertension, defined as blood pressure in adults >150/100mm Hg at screening;
6. A known clinically significant abnormal ECG, such as arrhythmia and active ischemia, within 6 months prior to treatment;
7. Xerostomia or hyposalivation;
8. Known allergy or intolerance to sympathomimetic drugs (e.g., pseudoephedrine, epinephrine), alcohol, or any NG11-2 excipient;
9. Receiving monoamine oxidase (MAO) inhibitors or antidepressants (tricyclic or imipramine types); Use of MAO inhibitor in the 14 days prior to treatment, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine;
10. History or current evidence of clinically significant laboratory abnormality or any disease condition that might confound the results of the study, or interfere with the patient's participation for the full duration of the study
11. Current participation in any other oral mucositis studies or completion/ withdrawal from any other oral mucositis studies within the last 3 months
12. Patient is, at the time of signing informed consent, a regular user of any illicit drugs, which would interfere with cooperation with the requirements of the trial;
13. Patient is pregnant or breastfeeding, or expecting to conceive or father children within the projected duration of the study;
14. Any other condition which, in the opinion of the Investigator, means that the patient is not a good candidate for study enrolment

Date of first enrolment

01/05/2023

Date of final enrolment

30/06/2024

Locations

Countries of recruitment

England

Northern Ireland

Scotland

United Kingdom

Study participating centre**Guys Hospital**

Guys Hospital

Great Maze Pond

London

United Kingdom

SE1 9RT

Study participating centre**Western General Hospital**

Crewe Road South

Edinburgh

Lothian

United Kingdom

EH4 2XU

Study participating centre**Aberdeen Royal Infirmary**

Foresterhill Road

Aberdeen

United Kingdom

AB25 2ZN

Study participating centre

Queen Elizabeth Hospital
Queen Elizabeth Medical Centre
Edgbaston
Birmingham
United Kingdom
B15 2TH

Study participating centre
Belfast City Hospital
51 Lisburn Rd
Belfast
United Kingdom
BT9 7AB

Sponsor information

Organisation
VasoDynamics Ltd

Sponsor details
Stevenage Biocatalyst Centre
Gunnels Wood Rd
Stevenage
England
United Kingdom
SG1 2FX
+44 1438 300880
gary.bower@vasodynamics.co.uk

Sponsor type
Industry

Funder(s)

Funder type
Industry

Funder Name
Vasodynamics

Results and Publications

Publication and dissemination plan

Peer reviewed scientific journals
Internal report
Conference presentation
Publication on website
Other publication
Submission to regulatory authorities
The study database will not be shared outside of the Sponsor group.
The overall study results will be disseminated on publicly available websites

Intention to publish date

28/02/2025

Individual participant data (IPD) sharing plan

Datasets are not expected to be made available due to Sponsor commercial confidentiality and the early phase nature of the study.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	version 1.2		12/05/2023	No	Yes
HRA research summary			28/06/2023	No	No
Participant information sheet	version 2.0		20/12/2023	No	Yes
Results article			11/03/2025	Yes	No