

A UK trial for the investigation of stable chest pain: can we improve patient experience, outcomes and NHS cost-efficiency compared to the current NICE guidelines?

| | | |
|--|---|---|
| Submission date 19/01/2022 | Recruitment status No longer recruiting | <input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol |
| Registration date 24/01/2022 | Overall study status Ongoing | <input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results |
| Last Edited 28/07/2025 | Condition category Circulatory System | <input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year |

Plain English summary of protocol

Background and study aims

Coronary artery disease (narrowing of the heart arteries) is the typical cause of angina (cardiac chest pain) and is becoming more common as people live longer with multiple heart disease risk factors (e.g. diabetes, smoking, high blood pressure, and high cholesterol). In the UK, about 2 million people have angina and about 200,000 are referred to hospital each year for further investigation. There are lots of different tests to investigate suspected angina, but doctors have different opinions as to the best approach, and international guidelines make very different recommendations. This leads to wide practice variation, inefficiency of healthcare resources and it may adversely impact on patient experience and outcomes. What most doctors agree on is that we do too many invasive X-ray angiograms. About 60% of angiograms performed do not find significant narrowing of the heart arteries, and as such, do not lead to a direct change in management for that patient.

Recent UK NICE guidelines recommend a non-invasive CT angiogram as the first-line test for all patients with suspected angina. This one-size-fits-all approach does not recognise individual patient characteristics, risk factors or likelihood of them having disease. Whilst cardiac CT is an excellent test to exclude narrowing of the heart arteries when they are normal, it can lead to over-estimation of severity when moderate disease is present. It has been suggested that this approach may lead to further increases in the rates of invasive angiography, more additional downstream investigations and increased NHS costs, with no difference in patient outcomes. In addition, the recent UK NICE guidelines if fully adopted would require major service reorganisation in every hospital, and large capital and revenue investment to accommodate the 700% predicted increase in cardiac CT capacity. Thus many doctors believe that this major NHS service reorganisation should first be robustly evaluated in the setting of a clinical trial to demonstrate that it is both appropriate and cost-efficient.

The aim of this study is to try to improve the investigation and management of patients with suspected cardiac chest pain (angina), both in terms of NHS resource use and also patient experience and outcomes; producing results that would be highly generalisable across the NHS.

Who can participate?

Males aged 45 years and over and females aged 50 years and over with atypical or typical angina and at least one major cardiac risk factor (diabetes, peripheral arterial disease, cerebrovascular disease, current or past tobacco use, high blood pressure, dyslipidaemia or a family history of premature coronary artery disease) referred to NHS cardiology outpatient services, requiring further investigation according to NICE CG95 guidelines, and who are deemed suitable for coronary revascularisation if required.

What does the study involve?

Patients will have either a computerised tomography cardioangiogram (a CT scan of the heart), a cardiac MRI scan, a stress echocardiogram (an ultrasound scan of the heart) or a nuclear perfusion scan of the heart with single-photon emission computerised tomography (MPS-SPECT), depending on which group they are randomly allocated to and their calculated risk of having heart disease. Patients will also complete three questionnaires at the start of the study and after 6 and 12 months.

What are the possible benefits and risks of participating?

All of the study procedures are part of routine clinical care and no additional procedures will be undertaken. Participants may be less likely to have unnecessary tests. Patients will also have more contact with the clinical care team and have more opportunities to ask questions and be informed about their health, which some patients find helpful.

Where is the study run from?

University of Leeds and Leeds Clinical Trials Research Unit (UK)

When is the study starting and how long is it expected to run for?

May 2019 to April 2027

Who is funding the study?

1. Heart Research UK
2. British Heart Foundation (UK) (quality of life sub-study)

Who is the main contact?

Dr Laura Jones
L.M.Jones@Leeds.ac.uk

Contact information

Type(s)

Principal investigator

Contact name

Prof John Greenwood

ORCID ID

<https://orcid.org/0000-0002-2861-0914>

Contact details

Division of Biomedical Imaging
LIGHT Building
Clarendon Way

Leeds
United Kingdom
LS2 9JT
+44 (0)113 343 8469
J.Greenwood@Leeds.ac.uk

Type(s)

Principal investigator

Contact name

Dr Peter Swoboda

ORCID ID

<https://orcid.org/0000-0001-7162-7079>

Contact details

Division of Biomedical Imaging
LIGHT Building
Clarendon Way
Leeds
United Kingdom
LS2 9JT
+44 (0)113 39 28206
P.Swoboda@Leeds.ac.uk

Type(s)

Public

Contact name

Dr Laura Jones

Contact details

Leeds Institute of Cardiovascular and Metabolic Medicine
University of Leeds
Leeds
United Kingdom
LS2 9JT
+44 (0)7921121815
L.M.jones@Leeds.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

302218

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

CPMS 50598, IRAS 302218

Study information

Scientific Title

A pragmatic approach to the investigation of stable chest pain: a UK, multi-centre, randomised trial to improve patient experience, outcomes and NHS cost efficiency

Acronym

CE-MARC 3

Study objectives

The primary question is: 'can we improve the management of patients with new-onset chest pain, both from a patient perspective and an NHS (payer) perspective?'

The proposed trial will address this by answering the following: compared to the current UK NICE guidelines, can a more pragmatic diagnostic pathway to guide the management of patients with suspected cardiac chest pain, result in a) lower rates of invasive angiography without evidence of obstructive disease, b) less downstream resource utilisation and c) improved patient experience and outcomes.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 01/10/2021, London Bridge Research Ethics Committee (postal address: not available; +44 (0)207 1048202; +44 (0)207 1048124; londonbridge.rec@hra.nhs.uk), REC ref: 21/LO/0666

Study design

Randomized; Both; Design type: Diagnosis, Imaging, Validation of investigation /therapeutic procedures

Primary study design

Interventional

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Stable chest pain

Interventions

People who come to the hospital outpatients department with stable new-onset chest pain or those that have been admitted to cardiology wards will be approached to take part in the trial. If they take part they will be in one of the two arms of the trial and this will be randomly decided (by a computer). The two strategies for the first-line test are:

1. Usual care following the NICE guidelines (this essentially means having a non-invasive CT coronary angiography (CTCA), anatomical test)
2. Pragmatic care following the European Society of Cardiology (ESC) 2019 guidelines whereby a

patient could get either a) an MRI scan of the heart (functional test), b) a stress echocardiogram (ultrasound scan) of the heart (functional test), c) a nuclear perfusion scan of the heart (functional test), d) a non-invasive CTCA with added computer calculations (fractional flow reserve [FFR]-CT, anatomical test) which is no different for the patient than having a CTCA. The choice of the first test in this arm would be by shared decision making between the patient and the doctor (cardiologist) guided by contemporary risk stratification determined by the doctor. There will be no extra visits but patients may be asked to complete three short (optional) questionnaires that ask about their health and well-being at baseline, 6 and 12 months. The questionnaires are validated and are expected to take no longer than 20 minutes to complete in total (at each time point). They can be completed during hospital visits or submitted electronically via the Clinical Trials Research portal (the link will be sent out via email or SMS).

In either arm, if the first test is abnormal patients may be asked to go for an invasive angiogram (this would also be the case if a patient was not in the trial). If the first test is normal then NICE guideline-directed medical therapy would be recommended by the doctor. If the first test is inconclusive then the patient could have second-line non-invasive testing or invasive angiography, based upon shared decision-making and in line with current UK NICE guidelines.

The researchers will collect all the test results to see how many of the invasive angiograms were abnormal in each of the trial arms (and how many were normal - i.e. how many showed no evidence of significant disease). They will analyse questionnaire results completed at baseline, 6 and 12 months, and will review medical records at the same intervals.

Patients will also be followed up for up to 10 years to see if they have any further downstream tests, and if they have any health problems like a heart attack (thankfully only very few people have).

Setting: Secondary care cardiology departments that have well-established clinical NHS services for both anatomical and functional cardiac imaging. Initially nine high volume, experienced research centres that are geographically spread and with ethnically diverse populations have been selected (Leeds, Glasgow, Leicester, Bristol, Oxford, Barts (London), Liverpool, Wakefield and Southampton); other centres could join based on their clinical services and trial experience.

Intervention Type

Other

Primary outcome(s)

Time to composite endpoint of cardiovascular death, myocardial infarction, unobstructive coronary artery disease (CAD) at invasive angiography, defined as the invasive reference standard FFR measurement of >0.80 (or instantaneous wave-free ratio [iFR] ≥ 0.90), i.e. no functional ischaemia, at the time of coronary angiography (or no coronary stenosis $>70\%$ on quantitative coronary angiography should FFR/iFR be deemed clinically inappropriate/unsafe to perform)

Key secondary outcome(s)

1. Patient-reported quality of life measured using the Seattle Angina Questionnaire (SAQ), Euroqol EQ-5D-5L, SF12v2 at 0, 6 and 12 months
2. Guideline-directed medical therapy (GDMT) usage measured using patient questionnaire results and/or GP records at 0, 6 and 12 months
3. Major Adverse Cardiovascular Events (all cause death, myocardial infarction [MI], unplanned percutaneous coronary intervention [PCI]/coronary artery bypass graft surgery [CABG])

measured using patient questionnaire results and/or GP records at 3 years

4. Resource use and cost efficiency measured using patient questionnaire results and/or GP records at 12 months

Completion date

26/04/2027

Eligibility

Key inclusion criteria

1. Males ≥ 45 years or females ≥ 50 years
2. Atypical or typical angina
3. At least one major cardiac risk factor (diabetes, peripheral arterial disease, cerebrovascular disease, current or past tobacco use, hypertension, dyslipidaemia or family history of premature coronary artery disease)
4. Referred to NHS cardiology outpatient services
5. Requiring further investigation according to NICE CG95 guidelines
6. Deemed suitable for coronary revascularisation if required

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

45 years

Sex

All

Key exclusion criteria

In order to keep the trial generalisable to the UK population at large, only issues related to patient safety/appropriateness will form the exclusion criteria; these will include:

1. Prior normal CT coronary angiography (CTCA) within the last 2 years or prior CTCA with extensive calcification
2. Clinically unstable cardiac symptoms
3. Known coronary artery disease (including previous myocardial infarction [MI], acute coronary syndrome [ACS] or coronary revascularization)
4. Contraindication to CTCA or functional cardiac imaging
5. Pregnancy and/or breastfeeding
6. Known chronic renal failure (eGFR $< 30\text{ml}/\text{min}/1.73\text{m}^2$)
7. Inability to give written informed consent

Date of first enrolment

27/04/2022

Date of final enrolment

08/08/2025

Locations**Countries of recruitment**

United Kingdom

England

Scotland

Study participating centre**United Leeds Teaching Hospital NHS Trust**

Trust Hq

Leeds General Infirmary

Great George St

Leeds

United Kingdom

LS1 3EX

Study participating centre**Oxford University Hospitals**

John Radcliffe Hospital

Headley Way

Headington

Oxford

United Kingdom

OX3 9DU

Study participating centre**Pinderfields Hospitals NHS Trust**

Trust Hq, Rowan House

Pinderfields General Hospital

Aberford Road

Wakefield

United Kingdom

WF1 4EE

Study participating centre**Northumbria Healthcare NHS Foundation Trust**

North Tyneside General Hospital

Rake Lane

North Shields
United Kingdom
NE29 8NH

Study participating centre
Harefield Hospital NHS Trust
Harefield
United Kingdom
UB9 6JH

Study participating centre
Royal Devon University Healthcare NHS Foundation Trust
Royal Devon University NHS Ft
Barrack Road
Exeter
United Kingdom
EX2 5DW

Study participating centre
South Tees Hospitals NHS Trust
Middlesbrough General Hospital
Ayresome Green Lane
Middlesbrough
United Kingdom
TS5 5AZ

Study participating centre
University Hospital Southampton NHS Foundation Trust
Southampton General Hospital
Tremona Road
Southampton
United Kingdom
SO16 6YD

Study participating centre
Central Manchester University Hospitals NHS Foundation Trust
Trust Headquarters, Cobbett House
Manchester Royal Infirmary
Oxford Road

Manchester
United Kingdom
M13 9WL

Study participating centre
Bradford Teaching Hospitals NHS Foundation Trust
Bradford Royal Infirmary
Duckworth Lane
Bradford
United Kingdom
BD9 6RJ

Study participating centre
Nenedoc (kettering)
Kettering General Hospital
Rothwell Road
Kettering
United Kingdom
NN16 8UZ

Study participating centre
The Royal Wolverhampton NHS Trust
New Cross Hospital
Wolverhampton Road
Heath Town
Wolverhampton
United Kingdom
WV10 0QP

Study participating centre
Norfolk & Norwich University Hospital
Colney Lane
Colney
Norwich
United Kingdom
NR4 7UY

Study participating centre
Nottingham University Hospitals NHS Trust - City Campus
Nottingham City Hospital
Hucknall Road

Nottingham
United Kingdom
NG5 1PB

Study participating centre
NIHR Barts Clinical Research Facility
11th Floor
Royal London Hospital
Stepney Way
London
United Kingdom
E1 1BB

Study participating centre
Kings College Hospital
Mapother House
De Crespigny Park
Denmark Hill
London
United Kingdom
SE5 8AB

Study participating centre
Glenfield General Hospital
Grobby Road
Leicester
United Kingdom
LE3 9QP

Study participating centre
The James Cook University Hospital
Marton Road
Middlesbrough
United Kingdom
TS4 3BW

Study participating centre
NIHR Barts Cardiovascular Biomedical Research Unit
The Royal London Hospital
Whitechapel
London

United Kingdom
E1 1BB

Study participating centre

North Bristol NHS Trust

Southmead Hospital
Southmead Road
Westbury-on-trym
Bristol
United Kingdom
BS10 5NB

Study participating centre

Basildon University Hospital

Nethermayne
Basildon
United Kingdom
SS16 5NL

Study participating centre

Queen Elizabeth University Hospital

1345 Govan Road
Glasgow
United Kingdom
G51 4TF

Sponsor information

Organisation

University of Leeds

ROR

<https://ror.org/024mrx33>

Funder(s)

Funder type

Charity

Funder Name

Heart Research UK; Grant Codes: TR2442/19/24

Alternative Name(s)

HRUK

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United Kingdom

Funder Name

British Heart Foundation

Alternative Name(s)

The British Heart Foundation, the_bhf, BHF

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request (Prof. John Greenwood; P.Greenwood@Leeds.ac.uk) following the publication of the pre-specified primary and secondary endpoints. Data will be fully anonymised.

IPD sharing plan summary

Available on request

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|---|-------------|--------------|------------|----------------|-----------------|
| Protocol article | | 13/06/2025 | 18/07/2025 | Yes | No |
| HRA research summary | | | 26/07/2023 | No | No |
| Participant information sheet | version 1.0 | 16/08/2021 | 20/01/2022 | No | Yes |

