# Is the cardiac model of rehabilitation is more effective than standard care in reducing cerebrovascular risk factors post-transient ischaemic attack?

Submission date	Recruitment status No longer recruiting	<ul><li>Prospectively registered</li></ul>		
08/07/2009		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
11/09/2009		[X] Results		
<b>Last Edited</b> 01/08/2013	Condition category Circulatory System	[] Individual participant data		

## Plain English summary of protocol

Not provided at time of registration

## Contact information

## Type(s)

Scientific

#### Contact name

Mr Hayden Kirk

## Contact details

Rehabilitation Research, MP 886 Southampton General Hospital Tremona Road Southampton United Kingdom SO16 6YD hjsk1k06@soton.ac.uk

# Additional identifiers

## Protocol serial number

RHM MED0832; PRF/08/04

# Study information

## Scientific Title

A single centre phase II feasibility study evaluating if the cardiac model of rehabilitation is more effective than standard care in reducing cerebrovascular risk factors post-transient ischaemic attack

## Acronym

**Ex4TIA Study** 

## Study objectives

Providing a cardiac model of rehabilitation (exercise and lifestyle strategies) for people within one month of a transient ischaemic attack (TIA)/minor stroke will improve their physiological status and reduce their vascular risk as defined by the cardiovascular and cerebrovascular score. We will ask the following questions:

- 1. Does the intervention produce significant physiological changes associated with atherosclerotic disease?
- 2. Does the intervention enhance lifestyle changes more effectively than standard care?
- 3. Do the patients perceive any benefit from the intervention in relation to standard care?

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Southampton and South West Hampshire Research Ethics Committee (B) approved on the 16th April 2009 (ref: 09/H0501/46)

## Study design

Single centre exploratory single-blind two group randomised controlled phase II feasibility study

## Primary study design

Interventional

## Study type(s)

Quality of life

## Health condition(s) or problem(s) studied

Cerebrovascular disease - transient ischaemic attack/minor strokes

## **Interventions**

Although a single centre study, this trial is registered for ethical purposes as a two site study as recruitment occurs in a different site (Acute Hospital Trust) to the intervention (Primary Care Trust).

## Intervention group:

Cardiac Rehabilitation Programme (Exercise and Lifestyle Education). Participants will have baseline data collected within 1 month of their cerebrovascular event following which they will then be randomised to standard care or standard care and cardiac rehabilitation with endpoint data collection 5 months subsequently. The intervention is an existing standard NHS cardiac rehabilitation programme consisting of Phases 2 - 4:

Phase 2 = risk stratification, education, family involvement and support

Phase 3 = structured exercise & education sessions

Phase 4 = patient directed continuation of the exercise component and the long term maintenance of individual goals

Control group:

Standard care.

All participants primary and secondary outcomes (with the exception of the qualitative interviews) will be recorded at the same time. Participants will have initial baseline measures recorded within one month of their cerebrovascular event (TIA/minor stroke). All participants will then continue for a five month period with standard care (routine TIA investigations and pharmaceutical treatments along with a short piece of lifestyle advice) or standard care and the intervention. All participants will return for endpoint data collection in the fifth month after initial baseline data collection. A number of patients will then be randomly selected (estimation 20) to be interviewed in order to gain a subjective analysis of their views regarding standard care and the cardiac model of rehabilitation.

## Intervention Type

Other

### **Phase**

Phase II

## Primary outcome(s)

Cardiac Risk Score: an algorithmic score that assesses the future risk of cardiac events based on age, sex, smoking status, resting blood pressure, diabetic status, total cholesterol and high-density/low-density lipoprotein cholesterol. Measured at baseline and 5 months.

## Key secondary outcome(s))

Measured at baseline and 5 months:

- 1. Biomarkers (C-reactive protein, fibrinogen)
- 2. Cardiovascular Disease Score: similar to the Cardiac Risk Score but allows for the identification of cerebrovascular risk
- 3. Exercise frequency: self-reported
- 4. Exercise capacity: maximal oxygen uptake (VO2) estimation derived from the Astrand-Rhyming cycle ergometer test (for participants not taking Beta Blockers)
- 5. Obesity: body mass index (BMI) and Waist-to-Hip ratio
- 6. Diet: quantity of fruit and vegetables consumed daily
- 7. Mood: Hospital Anxiety and Depression (HAD) score
- 8. Quality of Life: 36-item short form health survey (SF-36)
- 9. Qualitative analyses purposive sampling of up to 10 participants from each arm with thematic analyses of semi-structured interviews

## Completion date

20/10/2010

# **Eligibility**

# Key inclusion criteria

- 1. Diagnosis (within 1 month of incident) of:
- 1.1. TIA resolution of symptoms less than 24 hours of onset (not suspected patent foramen

ovale [PFO]), or

- 1.2. Minor stroke National Institutes of Health Stroke Scale (NIHSS) score less than 3
- 2. Lives within geographic locality (GP postcodes)
- 3. Independently mobile (can use stick but no falls within 2 months)
- 4. No significant visual/speech impairment
- 5. Cognitive capacity to undertake group exercises (no apparent dementia)
- 6. Able to give verbal and written consent
- 7. Aged greater than 18 years, either sex
- 8. Considered medically fit for exercise (Canadian Angina score and SIGN 2002 guidance)
- 9. No previous experience of cardiac rehabilitation
- 10. No current or recent participation in research

## Participant type(s)

Patient

## Healthy volunteers allowed

No

## Age group

Adult

## Lower age limit

18 years

### Sex

All

## Key exclusion criteria

- 1. TIA of suspected patent foramen ovale (hole in heart) origin with no other significant vascular risk factors
- 2. Live outside of the area covered by Southampton Cardiac Rehabilitation Team
- 3. Aphasia or other communication problems affecting ability to consent or to understand information given
- 4. Apparent dementia or significant cognitive impairment (mini-mental test [MMT] score less than 7)
- 5. Previously undertaken cardiac rehabilitation
- 6. Involved in current research or have recently been involved in any research

## Date of first enrolment

20/07/2009

## Date of final enrolment

20/10/2010

# Locations

### Countries of recruitment

United Kingdom

England

Study participating centre Rehabilitation Research, MP 886 Southampton United Kingdom SO16 6YD

# Sponsor information

## Organisation

Southampton University Hospitals NHS Trust (UK)

### ROR

https://ror.org/0485axj58

# Funder(s)

## Funder type

Research organisation

## **Funder Name**

Physiotherapy Research Foundation (UK) (ref: PRF/08/04)

## **Funder Name**

Private Physiotherapy Education Foundation (UK)

# **Results and Publications**

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration

## **Study outputs**

Output type	Details			Peer reviewed?	Patient-facing?
Results article	results (patients' experiences)	01/09/2013		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes