Efficacy and safety of lymphdiaral basistropfen in the treatment of chronic low-back pain

Submission date [] Prospectively registered Recruitment status 18/05/2009 No longer recruiting [] Protocol [] Statistical analysis plan Registration date Overall study status 17/06/2009 Completed [X] Results [] Individual participant data **Last Edited** Condition category 16/04/2019 Musculoskeletal Diseases

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Andre-Michael Beer

Contact details

Klinikum Blankenstein Modellabteilung für Naturheilkunde Im Vogelsang 5-11 Hattingen Germany D 45527

andre.beer@klinik-blankenstein.de

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number NCT01049373

Secondary identifying numbers

PSC 144/03

Study information

Scientific Title

Efficacy and safety of lymphdiaral basistropfen (a homoeopathic remedy [HDC]) in the treatment of chronic low-back pain considering consitution and diathesism: a double blind, randomised, placebo controlled, single-centre study

Study objectives

To evaluate superiority of HDC in comparison to placebo in the treatment of chronic low-back pain in relation to pain, functional impairment, quality of life, and state of health during a 15-week treatment period.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Bergmannsheil Bochum Ethikkommission, 23/07/2003

Study design

Interventional double-blind randomised placebo-controlled single-centre study

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Low-back pain

Interventions

Intervention: lymphdiaral basistropfen, a homoeopathic remedy of 12 drugs, oral solution Control: placebo oral solution

Each is to be taken 3 times daily, 10 drops in some liquid over 15 weeks.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Lymphdiaral basistropfen

Primary outcome measure

Change in FFbH-R following 15 weeks treatment

Secondary outcome measures

- 1. Change in FFbH-R following 2 weeks treatment
- 2. Change in pain score (SES) following 2 weeks treatment
- 3. Change in pain score (ES) following 15 weeks treatment
- 4. Change in strength of pain (Visual analog scale VAS) following 2 weeks treatment
- 5. Change in strength of pain (Visual analog scale VAS following 15 weeks treatment
- 6. Change in state of health (BF-S) following 2 weeks treatment
- 7. Change in state of health (BF-S) following 15 weeks treatment
- 8. Change in Oswestry Score following 2 weeks treatment
- 9. Change in Oswestry Score following 15 weeks treatment
- 10. Change in short form health survey 12 items (SF-12) following 2 weeks treatment
- 11. Change in short form health survey 12 items (SF-12) following 15 weeks treatment
- 12. Correlation of efficacy with the constitutional type of the patient, measured by the Hattinger Konstitutional Manual (HKM) and the Hattinger Konstitutional Questionnaire (HKF)
- 13. Amount of analgesics used
- 14. Number of days with incapability to work under 15 weeks treatment
- 15. Tolerability within 15 weeks treatment

Overall study start date

05/12/2003

Completion date

05/05/2007

Eligibility

Key inclusion criteria

- 1. Male and female patients
- 2. Aged 18 75 years
- 3. Chronic low-back pain lasting at least for 6 months
- 4. Hanover functional ability questionnaire (FFbH-R) score less than 70%
- 5. At least one of the following diagnoses:
- 5.1. Chronic lumbar ischialgia with or withour radicular radiation
- 5.2. Chronic degenerative lumbar syndrome
- 5.3. Spondylarthrosis
- 5.4. Chronic facette syndrome
- 5.5. Lumbage with protrusion of the intervertebral disc
- 5.6. Lumbar radiculopathia
- 5.7. Lumbar and other intervertebral disc impairments with radiculopathia
- 5.8. Back pain at different locations of the spine
- 6. Written, informed consent

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

140 (70 verum, 70 placebo)

Total final enrolment

137

Key exclusion criteria

- 1. Participation in another clinical trial/GCP-trial within 30 days prior to screening
- 2. Participation in this trial in an earlier time
- 3. Treatment with lymphdiaral basistropfen within 3 month prior to enrolment
- 4. Pregnancy and lactation
- 5. Non-compliance
- 6. Incapability to understand the sense of the study
- 7. Abuse of analgesics, opiates or other drugs
- 8. Chronic pain that are as strong as or even stronger than the pain caused by the low-back and that need to be treated with analgesics
- 9. Malign diseases
- 10. Pathological neurological states
- 11. Epilepsy
- 12. Operation of the spine within 3 month prior to enrolment
- 13. Fractures of the spine
- 14. Bechterew's disease
- 15. Alcohol abuse
- 16. Consuming diseases
- 17. Cachexia
- 18. Palsy of the legs or anal sphincter due to acute impairment of the intervertebral disc
- 19. Catheterisation or CT-controlled intra-articular injection in the lumbar region
- 20. Hypersensitivity against one of the ingredients or excipients of the study drugs or against composite plants in general
- 21. Systemic, progressive diseases like tuberculosis, leucosis, collagenosis, multiple sclerosis, acquired immune deficiency syndrome (AIDS), human immunodeficiency virus (HIV) infection, or other auto-immune diseases

Date of first enrolment

05/12/2003

Date of final enrolment

05/05/2007

Locations

Countries of recruitment

Germany

Study participating centre Klinikum Blankenstein Hattingen Germany

D 45527

Sponsor information

Organisation

PASCOE pharmazeutische Praeparate GmbH (Germany)

Sponsor details

Schiffenberger Weg 55 Giessen Germany D-35394

-

anja.braschoss@pascoe.de

Sponsor type

Industry

Website

http://www.pascoe-global.com

ROR

https://ror.org/02xbx2821

Funder(s)

Funder type

Industry

Funder Name

PASCOE pharmazeutische Praeparate GmbH (Germany)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results				No	No
Results article	results	28/06/2012	16/04/2019	Yes	No