# Repetitive transcranial magnetic stimulation (rTMS) for the treatment of chronic tinnitus

Submission date Recruitment status Prospectively registered 12/09/2007 No longer recruiting [X] Protocol [ ] Statistical analysis plan Registration date Overall study status 18/01/2008 Completed [X] Results [ ] Individual participant data **Last Edited** Condition category 17/08/2017 Nervous System Diseases

#### Plain English summary of protocol

Not provided at time of registration

# **Contact information**

#### Type(s)

Scientific

#### Contact name

Prof Goeran Hajak

#### Contact details

Universitaetsstr. 84 Regensburg Germany 93053 +49 (0)941 941 2011 goeran.hajak@medbo.de

#### Additional identifiers

Protocol serial number 05/06\_1

# Study information

#### Scientific Title

Repetitive transcranial magnetic stimulation (rTMS) for the treatment of chronic tinnitus

#### **Study objectives**

Chronic tinnitus is a severe and disabling disease with so far no efficient treatment. Accumulating data point to the involvement of dysfunctional neuronal activity in the central nervous system as one possible underlying cause of chronic tinnitus. rTMS has been shown to be able to non-invasively modulate cortical activity and holds therapeutic potential in other treatment-resistant diseases such as major depression. Pilot studies revealed promising therapeutic potential of rTMS in the treatment of chronic tinnitus.

The primary objective of this trial is to evaluate the efficacy of real rTMS versus sham rTMS in the treatment of chronic tinnitus by means of change of tinnitus severity according to the tinnitus questionnaire of Goebel and Hiller (baseline versus day 12).

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Ethics committee of the University of Regensburg, 24/10/2006

#### Study design

Randomized double-blind placebo-controlled multi-center trial

#### Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Chronic tinnitus

#### **Interventions**

rTMS will be administered according to current safety guidelines. Figure-of-eight-coils will be used for real stimulation. Sham stimulation will be carried out by tilting the coil 45° away from the skull with one wing touching the skull. The stimulation parameters have been chosen according to successful pilot studies. Patients will be randomized to 2 parallel treatment groups:

Group A will receive real stimulation: 2 x 5 sessions, 1 Hz rTMS, stimulation intensity 110% related to the individual motor threshold, 2000 stimuli per session, coil position 10-20 guided over left primary auditory cortex.

Group B will receive sham stimulation by angulation of the magnetic coil 45° away from the skull with one wing touching the skull. Coil positioning and stimulation parameters as for group A.

Treatment will be conducted over a period of 2 weeks, at a frequency of 5 sessions/week.

#### Intervention Type

Procedure/Surgery

#### Primary outcome(s)

Tinnitus severity, measured using the tinnitus questionnaire of Goebel and Hiller at baseline and day 12

#### Key secondary outcome(s))

Tinnitus severity, measured using the tinnitus questionnaire of Goebel and Hiller, Tinnitus Handicap Inventory (THI), Tinnitus Severity scale and Cinical Global Impression Scale during the follow-up period (screening, baseline, days 5, 67 and 181)

#### Further outcome measures:

- 1. Quality of life, measured by the 12-item Short Form health survey (SF-12) at baseline, days 5, 12, 18, 67 and 181
- 2. Depressive symptoms, measured by the Beck Depression Inventory (BDI) at baseline, days 5, 12, 18, 67 and 181
- 3. Psychometric parameters of tinnitus, assessed by audiological evaluation at screening and day 18
- 4. Structural neuroplastic adaptation processes, detected by voxel-based morphometry at baseline and day 12
- 5. Cortical excitability, assessed by paired-pulse TMS at baseline and day 12

#### Completion date

01/11/2010

# **Eligibility**

#### Key inclusion criteria

- 1. Male or female in- and out-patients, age 18-70
- 2. Diagnosis of chronic tinnitus
- 3. Patient has a score of greater than or equal to 38 on the Tinnitus Handicap Inventory
- 4. Tinnitus duration of more than 6 months
- 5. Age-adjusted normal sensorineuronal hearing determined by an audiogram within the last 4 weeks, i.e. no more than 5 dB below the 10% percentile (DIN EN ISO 7029) of the appropriate age and gender group in all measured standard frequencies. Furthermore, no conductive hearing loss of more than 15 dB in neither of the measured standard frequencies
- 6. Patient naïve to rTMS-treatment

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Upper age limit

70 years

#### Sex

All

#### Key exclusion criteria

- 1. Objective tinnitus
- 2. Other forms of tinnitus treatment at the same time
- 3. Clinically relevant psychiatric comorbidity as judged by an experienced psychiatrist
- 4. Concomitant treatment with psychotropic drugs
- 5. History of or evidence of significant brain malformation or neoplasm, head injury, cerebral vascular events, neurodegenerative disorder affecting the brain or prior brain surgery
- 6. Severe unstable somatic comorbidity
- 7. Cardiac pace makers, other electronic implants, intracranial metallic particles
- 8. History of seizures or epileptiform activity
- 9. Pregnancy and lactation
- 10. Women in child bearing age without contraception
- 11. Patients who cannot communicate reliably with the investigator or who are not likely to cope with the requirements of the trial
- 12. Patient unwilling or unable to give written informed consent
- 13. Participation in a clinical trial within the last 30 days before start of this clinical trial or similar participation in another clinical trial

# Date of first enrolment 01/11/2007

Date of final enrolment 01/11/2010

#### Locations

#### **Countries of recruitment** Germany

Study participating centre University of Regensburg Regensburg Germany 93053

# Sponsor information

#### Organisation

University of Regensburg (Germany)

#### **ROR**

https://ror.org/01eezs655

# Funder(s)

#### Funder type

Government

#### Funder Name

Deutsche Forschungsgemeinschaft within a clinical studies programme (ref: HA 3547/4-1)

#### Alternative Name(s)

German Research Association, German Research Foundation, Deutsche Forschungsgemeinschaft (DFG), DFG

#### **Funding Body Type**

Government organisation

#### Funding Body Subtype

National government

#### Location

Germany

## **Results and Publications**

Individual participant data (IPD) sharing plan

### IPD sharing plan summary

#### **Study outputs**

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	results	01/11/2017	Yes	No
Protocol article	protocol	15/04/2008	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes