Treatment of Diabetic Foot Wounds by Vacuum-Assisted Closure (VAC®)

Submission date	Recruitment status	[X] Prospectively registered
13/07/2010	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
29/07/2010	Completed	Results
Last Edited	Condition category	Individual participant data
29/07/2010	Nutritional, Metabolic, Endocrine	Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Ms Dörthe Seidel

Contact details

Institut für Forschung in der Operativen Medizin Private Universität Witten / Herdecke gGmbH Ostmerheimerstraße 200 Haus 38 Köln Germany 51109 +49 (0) 221 9895719 doerthe.seidel@uni-wh.de

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

DRKS00000059

Study information

Scientific Title

Treatment of Diabetic Foot Wounds by Vacuum-Assisted Closure (VAC®): A multicentre, randomised controlled trial

Acronym

DiaFoVAC

Study objectives

This randomised controlled trial (RCT) comparing vacuum-assisted closure and conventional moist wound therapy (CMWT) for diabetic foot wounds was designed in order to:

- 1. Prove that VAC® therapy leads to complete healing of diabetic foot wounds more of-ten and earlier as compared to conventional moist wound therapy
- 2. Show that VAC® therapy is safe and effective also when provided in an ambulatory care setting
- 3. Generate more data on optimal patient selection and optimal combination of VAC® therapy with surgical closure

Ethics approval required

Old ethics approval format

Ethics approval(s)

The local ethics committee (Ethik-Kommission der Universität Witten/Herdecke) approved on the 8th of May 2009 (ref: 12/2009)

Study design

Multicentre randomised controlled clinical superiority trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Diabetic foot syndrome / chronic or amputational diabetic foot wound / wound treatment

Interventions

Participants will be randomised to receive either:

- 1. Intervention: Vacuum-assisted-closure® based on negative pressure wound therapy (V.A.C® NPWT)
- 2. Control / Comparator: Conventional Moist Wound Therapy (CMWT) according to current available evidence based guidelines

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

Time until complete (100%) wound closure, defined by the following criteria:

- 1. Full (100%) re-tepithelialization of the wound (i.e. no granulation tissue visible)
- 2. No drainage
- 3. No need for wound dressing

Secondary outcome measures

- 1. Clinical endpoints
- 1.1. Wound size (area of wound opening, maximal length and width)
- 1.2. Composition of wound tissue (percentage of epithelia layer, granulation tissue and fibrin)
- 1.3. Pain in wound surrounding (until wound closure and at 12 months)
- 1.4. Incidence and time of wound bed preparation
- 1.5. Secondary resections and amputations
- 1.6. Recurrence of wound or ulcer at same site
- 1.7. Complication rates (local and systemic)
- 1.8. Total complication rate
- 1.9. Mortality of any cause (within 12 month)
- 2. Health economic endpoints
- 2.1. Number of dressing changes
- 2.2. Length of treatment
- 2.3. Length of hospital stay
- 2.4. Time until orthopedic footwear supply possible
- 2.5. Time until return to work
- 2.6. Quality-of-life during treatment
- 2.7. Quality of life after Healing (at 6 and 12 months)
- 2.8. Patient satisfaction
- 2.9. Costs (in and out hospital)

Overall study start date

01/06/2011

Completion date

30/09/2013

Eligibility

Key inclusion criteria

- 1. Evidence of diabetes mellitus type I, II or III as defined by the necessity for antidiabetic drugs for ≥ 12 months
- 2. All wounds corresponded to University of Texas grade 2 or 3 in depth
- 2.1. either a chronic wound which is planned to receive radical surgical debridement
- 2.2. or an amputational wound which results from a planned below-ankle amputation
- 3. Participant has given written informed consent

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

300

Key exclusion criteria

- 1. Pregnancy
- 2. Age <18 years
- 3. Missing Compliance
- 4. Untreated or refractory phlegmona or untreated osteomyelitis of the wound
- 5. Exposed blood vessels in or around the wound that can't be protected or have a heightened risk for a bleeding with relevant circulation effects
- 6. Any revascularization procedure or amputation of the same extremity within the last 5 days
- 7. Indication for above-ankle amputation of the same extremity
- 8. Any extreme defective position of toes, foot or the whole extremity which could lead to ischemia of the wound area and impair wound healing
- 9. Evidence of inadequate perfusion
- 9.1. via MR angiography within the last 3 month or
- 9.2. sonographic confirmation of absence of peripheral pulse at A. tibialis posterior or A. dorsalis pedis) or
- 9.3. transcutaneous oxygen measurements of the dorsum of the foot ≤30 mmHq or
- 9.4. ankle brachial indices ≥ 0.7 and ≤ 1.2 and toe pressure at ≤ 30 mmHg
- 10. Renal insufficiency with need for dialysis
- 11. Anemia which is not caused infection
- 12. Long term use of oral (p.o.) steroids (> 30days)
- 13. Chronic anticoagulation except acetylsalicylate, clopidogrel (e.g. by vitamin K an-tagonists) at screening that can't be bridged to heparins during participation in the trial
- 14. Nerve lesions or active Charcot disease (Diabetic neuro-osteoarthropathy) of either lower extremity that will interfere with wound treatment
- 15. Malignancy in the wound or the periwound area or any other malignancy requiring immunosuppressant therapy or chemotherapy
- 16. Wounds resulting from burns, untreated cellulitis or collagen vascular disease
- 17. Presence of necrotic tissue that cannot be debrided
- 18. Concomitant severe skin disease that may impair wound healing
- 19. Connective tissue disease that might impair wound healing
- 20. Hematological disorders or condition that might impair wound healing

- 21. History of significant chronic anemia as evidenced by hemoglobin concentration of < 10 g/dl within 30 days of screening
- 22. Any kind of chronic venous insufficiency (CVI) or CVI that leads to varicose eczema, lo-cal inflammation and an increased risk of ulcers
- 23. Use of VAC® negative pressure wound therapy (NPWT) or any other suction device system on the study wound \leq 8 days prior to screening
- 24. Previous treatment with hyperbaric oxygen therapy (HBO) or normothermic therapy in the past 8 days
- 25. Application of recombinant or autologous growth factors on the study wound \leq 8 days prior to screening
- 26. Application of skin or dermal substitutes and dressings with living cells capable of producing growth factors (e.g. Oasis®, Apligraf®, Dermagraft™, or Integra®) on the study extremity < 15 days prior to screening
- 27. Allergy to any disposal component of the VAC® or MWT used materials
- 28. Simultaneous participation in other interventional trials interfering with one of the approaches in this trial
- 29. Previous participation in this trial

Date of first enrolment

01/06/2011

Date of final enrolment

30/09/2013

Locations

Countries of recruitment

Germany

Study participating centre Institut für Forschung in der Operativen Medizin Köln Germany 51109

Sponsor information

Organisation

Private University of Witten/Herdecke gGmbH (Germany)

Sponsor details

Medical Faculty Alfred-Herrhausen-Str. 50 Witten Germany 58448 sekretariat-neugebauer@uni-wh.de

Sponsor type

University/education

Website

http://www.uni-wh.de

ROR

https://ror.org/00yq55g44

Funder(s)

Funder type

Industry

Funder Name

Kinetic Concepts, Inc. (USA)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration