# Blood flow of the knee cap in patients with knee pain

Submission date	Recruitment status  No longer recruiting	[X] Prospectively registered		
10/12/2021		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
15/12/2021		[X] Results		
Last Edited	Condition category	Individual participant data		
29/11/2023	Musculoskeletal Diseases			

#### Plain English summary of protocol

Background and study aims

Patellofemoral pain (pain at the front of your knee, around your kneecap) is a frequent and often long-standing painful condition of the knee. Although the available scientific evidence suggests that exercise therapy is an effective treatment, there are still a lot of patients who continue to suffer from knee pain. This study elaborates on a theory of disturbed bone blood flow in the bone of the knee cap. Reduced oxygen saturation of that specific tissue potentially explains the pain while prolonged sitting and descending stairs, but this has never been studied before. Therefore the aim of the study is to observe bone blood perfusion while the participant is seated (protocol 1) and simulates descending stairs (protocol 2). The oxygen saturation will be measured with a spectroscope, which is a non-invasive device and will be placed on the skin of the knee cap.

#### Who can participate?

Patients with patellofemoral pain and healthy persons can participate. Patellofemoral pain is defined as pain around or behind the knee cap and usually gets worse while descending stairs or during activities like running, jumping, and hopping. Before participating in this study the principal investigator will assess the presence of patellofemoral pain and exclude other conditions of the knee that could explain symptoms. Healthy persons are required to be completely pain-free and without other specific conditions like surgery in the past. Fulfillment of these requirements for healthy persons will also be assessed by the principal investigator prior to participation.

#### What does the study involve?

Persons who meet the criteria described above will be asked to stay seated for 15 minutes. Thereafter the spectroscope measurements will be started and the patient bends the knee to 90 degrees and will be asked to keep in that position for 30 minutes while the measurement of bone blood perfusion will take place (protocol 1). Thereafter, the participant will be asked to stand up and simulate descending stairs (protocol 2) while again the spectroscope measurements will be performed.

What are the possible benefits and risks of participating?
By participating in this study one can contribute to developing new insights into the bone blood

perfusion and maybe change the way of thinking of best treatment strategies for this frequent and often long-standing painful condition. The Medical Ethics Committee concluded that the risks for participants are low to absent. This is due to the non-invasive character of the bone blood perfusion measurements.

Where is the study run from?

The study runs from the orthopedic surgery department of the Amsterdam Medical Centers in the Netherlands.

When is the study starting and how long is it expected to run for? February 2021 to December 2022

Who is funding the study? Investigator initiated and funded

Who is the main contact?
Martin Ophey, m.j.ophey@amsterdamumc.nl

#### Contact information

#### Type(s)

Scientific

#### Contact name

Mr Martin Ophey

#### **ORCID ID**

https://orcid.org/0000-0002-8674-0477

#### Contact details

Meibergdreef 9 Amsterdam Netherlands 1105AZ +31-650616509 m.j.ophey@amsterdamumc.nl

#### Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

NL77408.018.21

## Study information

#### Scientific Title

Bone blood perfusion of the patella measured with non-invasive Near Infrared Spectroscopy in patients with patellofemoral pain

#### **Acronym**

NIRSpatella

#### **Study objectives**

In patients with patellofemoral pain bone blood perfusion of the patella bone is reduced when compared to healthy controls.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Approved 15/11/2021, Medical Ethics Committee of the Amsterdam Medical Center (AMC, Meibergdreef 9, 1105AZ, Amsterdam, the Netherlands; +31(0)205669111; indienenmetc@amsterdamumc.nl), ref: 2021\_088#B2021652

#### Study design

Single-center single-blinded observational case-control study

#### Primary study design

Observational

#### Study type(s)

Other

#### Health condition(s) or problem(s) studied

Patellofemoral pain in young adults

#### **Interventions**

Bone blood perfusion will be measured with non-invasive Near InfraRed Spectroscopy (NIRS). The measurements will take place in two different postures, which usually provoke symptoms of knee pain in patients with patellfemoral pain. The first posture is prolonged sitting with the knee bend for 30 minutes and the second posture mimics descending stairs on a decline stepdown exercise.

#### Intervention Type

Other

#### Primary outcome(s)

Bone blood perfusion expressed as concentrations of oxygenated hemoglobin (O2Hb) and deoxygenated hemoglobin (HHb) measured using NIRS during the two conditions.

#### Key secondary outcome(s))

Measured at a single time point (baseline) unless otherwise stated:

- 1. Anterior Knee Pain Scale Dutch Version (AKPS-DV), 13 item questionnaire
- 2. Tegner Score, 10 item level of physical activity
- 3. Blood pressure (mmHq), sphygmomanometer

- 4. Prepatellar skinfold thickness measured with Harpenden skinfold caliper
- 5. Patella width measured with caliper
- 6. Decline step-down test (DSDT) measuring pain-free flexion angle
- 7. Lower limb range of motion (LLROM) measuring soft tissue flexibility of the kinetic chain
- 8. Visual Analogue Scale (VAS) measuring pain intensity while performing protocol 1 and 2

#### Completion date

31/12/2022

## Eligibility

#### Key inclusion criteria

- 1. Age 18-40 years
- 2. Diagnosis of patellofemoral pain for more than 3 months
- 3. No other specific knee conditions (ACL, meniscus, OA, jumpers knee)
- 4. Non-traumatic onset of PFP

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Upper age limit

40 years

#### Sex

ΔII

#### Total final enrolment

60

#### Key exclusion criteria

- 1. PFP due to trauma or post-surgery
- 2. Other specific knee conditions (ACL, meniscus, OA, jumpers knee)

#### Date of first enrolment

01/02/2022

#### Date of final enrolment

10/05/2022

#### Locations

#### Countries of recruitment

Netherlands

# Study participating centre YsveldFysio

Nieuwstraat 9 Nijmegen Netherlands 6511PT

# Sponsor information

#### Organisation

Amsterdam University Medical Centers

#### **ROR**

https://ror.org/05grdyy37

# Funder(s)

#### Funder type

Other

#### **Funder Name**

Investigator initiated and funded

#### **Results and Publications**

#### Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study will be published as a supplement to the subsequent results publication.

#### IPD sharing plan summary

Published as a supplement to the results publication

#### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		29/11/2023	29/11/2023	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes