The protective effect of a nasal corticosteroid (Avamys[®]) on exercise induced airway obstruction in cold air in children

Submission date 09/02/2009	Recruitment status No longer recruiting	Prospectively registered
		[_] Protocol
Registration date	Overall study status	[] Statistical analysis plan
03/07/2009	Completed	[_] Results
Last Edited	Condition category	Individual participant data
03/07/2009	Respiratory	[] Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s) Scientific

Contact name Mrs Janneke van Leeuwen

Contact details

Ariensplein 1 Enschede Netherlands 75 11 JX +31 (0) 53 487 2310 J.C.van.Leeuwen@student.rug.nl

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers FF1, NL26953.044.09

Study information

Scientific Title

The protective effect of a nasal corticosteroid (Avamys®) on exercise induced airway obstruction in cold air in children: a randomised double-blind placebo-controlled single-centre trial

Acronym

YSCO

Study objectives

Three weeks of treatment with fluticasone furoate (Avamys®) in children will reduce exercise induced fall in forced expiratory volume in one second (FEV1) and maximum inspiratory flow rate at 50% of vital capacity (MIF50).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Medical Ethical Review Committee (METC), Medical Centre Twente (Medisch Spectrum Twente), Enschede, approval pending as of 03/07/2009.

Study design

Randomised double-blind placebo-controlled single-centre trial

Primary study design

Interventional

Secondary study design Randomised controlled trial

Study setting(s) Other

Study type(s) Prevention

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Allergic rhinitis and exercise induced asthma in children

Interventions

Three weeks +/- 5 days of treatment with fluticasone furoate (Avamys®) or placebo. Avamys® will be administered through a nasal spray, once daily, 27.5 µg in each nostril. In the first week of the study there will be a double dosing.

Intervention Type

Drug

Phase Not Applicable

Drug/device/biological/vaccine name(s)

Fluticasone furoate (Avamys®)

Primary outcome measure

1. Analyse the reduction in exercise induced fall of FEV1 after three weeks of treatment with fluticasone furoate

2. Analyse the reduction in exercise induced fall of MIF50 after three weeks of treatment with fluticasone furoate

Secondary outcome measures

1. To analyse the reduction in exercise induced increase of airway resistance, measured with the forced oscillation technique (FOT), after three weeks of treatment with fluticasone furoate 2. To analyse the reduction in exercise induced decrease of airway reactance, measured with the forced oscillation technique (FOT), after three weeks of treatment with fluticasone furoate 3. Analyze the reduction in fractional exhaled nitric oxide (FeNO), measured with miniNIOX® after three weeks of treatment with fluticasone furoate

4. To analyse the increase in quality of life, measured with the Paediatric Asthma Quality of Life Questionnaire (PAQLQ), after three weeks of treatment with fluticasone furoate 5. To analyse the increase in control of asthma, measured with the Asthma Control Questionnaire (ACQ), after three weeks of treatment with fluticasone furoate

Overall study start date

05/03/2009

Completion date

05/04/2009

Eligibility

Key inclusion criteria

1. Both males and females, aged between 12 and 17 years

2. Clinical history of allergic rhinitis and/or allergic asthma

3. Ability to perform reproducible lung function tests, i.e. coefficient of the predicted value variation in three of five consecutive measurements less than 5%

4. Maximal FEV1 greater than 70% of predicted value

5. Clinically stable period at least three weeks before the study period

Participant type(s)

Patient

Age group Child

Lower age limit 12 Years Upper age limit

17 Years

Sex Both

Target number of participants

91

Key exclusion criteria

1. Use of intranasal or systemic corticosteroids in the last four weeks prior to the study

2. Use of antihistamines, cromoglycates, anticholinergics in two weeks prior to the study

3. Use of long acting bronchodilators 24 hours before testing

4. Use of short acting bronchodilators eight hours before testing

5. Use of systemic corticosteroids, antihistamines, cromoglycates, anticholinergics, during the study

6. Other pulmonary or cardiac disorder

7. Deviation of the FEV1 of more than 12% from baseline spirometry and the FEV1 before subsequent exercise provocation challenges

8. Signs of gastro-oesophageal reflux

Date of first enrolment 05/03/2009

Date of final enrolment 05/04/2009

Locations

Countries of recruitment Netherlands

Study participating centre Ariensplein 1 Enschede Netherlands 75 11 JX

Sponsor information

Organisation Paediatric Research Foundation Enschede, Medical Centre Twente (Netherlands)

Sponsor details

Ariensplein 1 Enschede Netherlands 7511 JX +31 (0) 53 487 2310 kindergeneeskunde@ziekenhuis-mst.nl

Sponsor type Hospital/treatment centre

Website http://www.mstwente.nl

ROR https://ror.org/033xvax87

Funder(s)

Funder type Hospital/treatment centre

Funder Name

Paediatric Research Foundation Enschede, Medical Centre Twente (Stichting Pediatrisch Onderzoek Enschede, Medisch Spectrum Twente) (Netherlands)

Results and Publications

Publication and dissemination plan Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary Not provided at time of registration