S-ketamine to reduce negative pressure wound therapy dressing change associated pain

Submission date 01/12/2013	Recruitment status Stopped	[X] Prospectively registered [_] Protocol
Registration date	Overall study status	Statistical analysis plan 1 Results 1 Results
Last Edited	Condition category	 Individual participant data Decord updated in last year
11/07/2022	Signs and Symptoms	

Plain English summary of protocol

Background and study aims

We are carrying out a study on patients undergoing negative pressure wound therapy dressing changes. Common analgesics (painkillers) such as local anesthesia with lidocaine alone are often not effective for this procedure. Opioids are the most effective analgesics but their use is limited by numerous side effects, such as nausea, vomiting, constipation, urinary retention, dysphoria (an emotional state characterized by anxiety, depression or unease), euphoria (a feeling or state of intense excitement and happiness), altered cognitive (mental) function and hypoventilation (slow breathing). Other sedatives used for procedural sedation such as propofol or benzodiazepines offer no analgesia (pain relief) and may cause hypoventilation, alterations in hemodynamics and drowsiness. Ketamine is an old anaesthetic agent that also has analgesic properties. Ketamine also reduces the need for opioids and thus helps to reduce side effects caused by opioid medication.

Who can participate?

Adult patients aged over 18 undergoing elective wound dressing changes in the plastic surgery unit. 23 patients will be enrolled in the study.

What does the study involve?

The wound dressing changes will take place every 3-4 days for each patient. Each patient will be randomly allocated to receive either S-ketamine or saline on their first procedure and then on the second procedure the patient will be given the remaining medicine. Each patient will be given midazolam as a premedication before the procedure and a solution containing lidocaine with saline. If the pain is too great during the procedure, the patient will be given fentanyl as a rescue medication.

What are the possible benefits and risks of participating?

The possible benefits of participating in the study are improved analgesia during the procedure, reduced need for rescue medication (fentanyl) and fewer side effects caused by opioid medication. Possible risks are the side effects caused by ketamine. The most common side effect of ketamine is altered mental state. However, that can be prevented by giving midazolam as a premedication.

Where is the study run from?

The study is run from the Helsinki University Central Hospital Töölö Hospital, Finland.

When is the study starting and how long is it expected to run for? It is anticipated that recruitment for the study will start in April 2014. Participants will be enrolled on the study for a period of one year, recruiting the last patient on April 2015. However, the recruitment period will be extended if necessary until 23 patients are enrolled.

Who is funding the study? Helsinki University Central Hospital, Finland.

Who is the main contact? Dr Vesa Kontinen Dr Elina Brinck

Contact information

Type(s) Scientific

Contact name Dr Elina Brinck

Contact details Helsinki University Central Hospital Töölö Hospital Topeliuksenkatu 5 PB 266 Helsinki Finland 00029

Additional identifiers

EudraCT/CTIS number 2013-005114-35

IRAS number

ClinicalTrials.gov number Nil known

Secondary identifying numbers Nil known

Study information

Scientific Title

S-ketamine to reduce negative pressure wound therapy dressing change associated pain: a prospective, randomized, blinded cross-over study

Acronym KETAPLAST

Study objectives

S-ketamine effectively alleviates procedural pain and reduces the need for rescue medication (fentanyl) during negative pressure wound therapy dressing change compared to local anaesthesia.

Ethics approval required Old ethics approval format

Ethics approval(s) Operative ethics committee of the Helsinki University Central Hospital, 12t/02/2014, ref: 38/13 /03/02/2014

Study design Prospective randomized blinded cross-over study

Primary study design Interventional

Secondary study design Randomised cross over trial

Study setting(s) Hospital

Study type(s) Treatment

Participant information sheet See additional files

Health condition(s) or problem(s) studied

Procedural pain related to negative pressure wound therapy dressing changes

Interventions

 Intervention group: S-ketamine dose 0.25 mg/kg will be administered during negative pressure wound therapy dressing change in addition to local anaesthesia
 Control group: placebo (saline)

The wound dressing changes will take place every 3-4 days for each patient. Each patient will be randomized to receive either S-ketamine or saline on their first procedure and then on the second procedure, she or he will be given the remaining medicine. Each patient will be given midazolam as a premedication before the procedure and a solution containing Lidocaine 1% with saline will be absorbed into the tissue treated. If the pain score (NRS scale) exceeds 5 during the procedure, she or he will be given fentanyl 0.5-1 ug/kg as a rescue medication.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

S-ketamine

Primary outcome measure

1. Pain measured using the NRS scale during the procedure, at the end of the procedure, and at three months after the procedure

2. Possible need for rescue medicine (fentanyl) during the procedure and, if needed, how much

Secondary outcome measures

To evaluate and compare the possible side effects such as altered mental state, desaturation, changes in blood pressure and heart rate associated with the treatment of S-ketamine or rescue medication (fentanyl):

1. Confusion measured with Richmond Agitation Sedation Scale (RASS) at 30 minutes and 60 minutes after the drug investigated (S-ketamine or placebo) has been given

2. Saturation with pulse oximetry (SpO2 %) at 15 minutes, 30 minutes and 60 minutes after the drug investigated (S-ketamine or placebo) has been given

3. Hemodynamic parameters (blood pressure mmHg and pulse beats/minute) at 15 minutes, 30 minutes and 60 minutes after the drug investigated (S-ketamine or placebo) has been given

Overall study start date

01/04/2014

Completion date

30/06/2021

Reason abandoned (if study stopped)

Lack of staff/facilities/resources

Eligibility

Key inclusion criteria

Adult patients over 18 years undergoing elective negative pressure wound therapy wound dressing changes in the plastic surgery unit

Participant type(s) Patient

Age group Adult

Lower age limit 18 Years

Sex Both

Target number of participants

23

Key exclusion criteria

- 1. Body mass index over 35
- 2. Unstable ischaemic cardiac disease
- 3. Increased intracranial pressure
- 4. Increased intraocular pressure
- 5. Gravidity
- 6. Lactation
- 7. Hypersensitivity or allergy to ketamine, fentanyl or midazolam
- 8. Severe psychiatric disease
- 9. Inability to use the Numerical Rating Scale (NRS) pain scale

Date of first enrolment 01/04/2014

Date of final enrolment

31/12/2020

Locations

Countries of recruitment Finland

Study participating centre Helsinki University Central Hospital Helsinki Finland 00029

Sponsor information

Organisation

Helsinki University Central Hospital (Finland)

Sponsor details

c/o Vesa Kontinen Jorvi Hospital Turuntie 150 PB 800 Helsinki Finland 00029 **Sponsor type** Hospital/treatment centre

Website http://www.hus.fi/en/Pages/default.aspx

ROR https://ror.org/02e8hzf44

Funder(s)

Funder type Hospital/treatment centre

Funder Name

Helsinki University Central Hospital (Finland)

Results and Publications

Publication and dissemination plan

This clinical trial will be a part of a doctoral thesis (In Finland, 3-5 trials are required for PhD in Medicine). Plans to publish the results of in a high-impact peer reviewed journal during 2018-2019.

Intention to publish date

30/06/2022

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from elina.brinck@hus.fi

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet		19/01/2016	19/01/2017	No	Yes