

# Vascular function in patients with chronic vessel inflammation

<b>Submission date</b> 05/01/2016	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 09/01/2016	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 07/01/2016	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Chronic periaortitis is a rare disease of unknown cause that leads to inflammatory tissue around the body's main artery, the aorta. It is known that increased arterial stiffness is associated with an increased risk of heart disease and death. We aim to study arterial stiffness in patients with chronic periaortitis.

### Who can participate?

Patients with chronic periaortitis, patients with high blood pressure, and 16 healthy volunteers, matched for sex and age (age range: 34 to 74).

### What does the study involve?

Participants have their arterial stiffness measured with two methods: pulse wave velocity (a probe placed over the neck artery and the main leg artery) and digital volume pulse (a probe placed on the index finger). We study the change of digital volume pulse before and after giving the patients the drug nitroglycerin sublingually (under the tongue).

### What are the possible benefits and risks of participating?

Benefits include knowing whether you have a marker of heart disease risk and contributing to knowledge about a very rare disease. Sublingual nitroglycerin can sometimes cause a headache, sickness and dizziness. These side effects usually last less than 10 minutes and are not dangerous.

### Where is the study run from?

University Hospital of Bern (Switzerland).

### When is the study starting and how long is it expected to run for?

January to May 2008.

### Who is funding the study?

Investigator initiated and funded

Who is the main contact?

Dr Claudia Scheuter

## Contact information

### Type(s)

Scientific

### Contact name

Dr Claudia Scheuter

### ORCID ID

<https://orcid.org/0000-0001-7088-3652>

### Contact details

Department of Nephrology and Hypertension

Inselspital University Hospital

Bern

Switzerland

3010

## Additional identifiers

### Protocol serial number

1491

## Study information

### Scientific Title

Arterial stiffness and vascular function in patients with chronic periaortitis – a case-control study

### Study objectives

Patients with periaortitis show an elevated arterial stiffness and a reduced vascular function

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Kantonale Ethikkommission Bern (Ethics Committee of the State of Bern, Switzerland), 07/04/2008, ref: 025/08

### Study design

Observational cross-sectional case-control Study

### Primary study design

Observational

### Study type(s)

Diagnostic

### **Health condition(s) or problem(s) studied**

Chronic periaortitis, also known as Ormond's disease or retroperitoneal fibrosis is a rare disease of unknown etiology with pathological features of vascular inflammation and fibrosis.

### **Interventions**

We compared arterial stiffness and endothelial function in 16 patients with chronic periaortitis to age- and sex-matched hypertensive and healthy controls in a 1:1 distribution. Aortic stiffness was assessed by carotid-to-femoral pulse wave velocity with Doppler measurements. Endothelium-independent vasodilation was assessed before and after sublingual administration of glyceryl trinitrate by analyzing changes in arterial stiffness using digital volume pulse (DVP) and the DVP derived stiffness index (SIDVP).

### **Intervention Type**

Device

### **Primary outcome(s)**

Arterial stiffness, measured by pulse wave velocity (PWV). PWV was recorded in a single session, using a Pulse Trace PWV© (Micro Medical Ltd, Rochester, Kent, UK) with a 4-MHz continuous wave directional Doppler pencil probe. Pulse Trace PWV© calculates the time lag between the R wave of the electrocardiogram and the arrival of the arterial pulse wave at 2 different sites, thus inferring pulse wave velocity. First, electrodes for an electrocardiogram are placed, followed by placing the Doppler probe over (1) the right carotid artery and (2) the right or left femoral artery. The position of the Doppler probe is optimized at each site to achieve a signal of sufficient quality (i.e. stable pulse waveform). About 10 systolic peaks are recorded at both sites to obtain a mean time lag at the carotid artery and at the femoral artery. The distance between the 2 detection sites is measured.

### **Key secondary outcome(s)**

Digital volume pulse (DVP) as a measure for endothelial function, blood pressure, heart rate and pulse pressure. DVP was measured photoplethysmographically in the same single session as PWV, using the Pulse Trace© Pulse Contour Analysis device (Micro Medical Ltd, Rochester, Kent, UK). The probe were placed on the index finger of the dominant hand. Pulse contour waves recorded over a period of 10 seconds were averaged by the device to a single waveform. Blood pressure was measured while participants were recumbent for at least 10 minutes.

### **Completion date**

30/05/2008

## **Eligibility**

### **Key inclusion criteria**

Patients: patients with a diagnosis of chronic periaortitis that were followed at our outpatient clinic for Nephrology and Hypertension and patients with known arterial hypertension that were followed at our outpatient clinic for Nephrology and Hypertension, matched for age and sex in a 1:1 fashion.

Healthy volunteers: healthy volunteers from the community and clinic staff, matched 1:1 for age and sex.

**Participant type(s)**

Mixed

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Key exclusion criteria**

Patients with atrial fibrillation and other cardiac arrhythmias that interfered with study measurements

**Date of first enrolment**

07/04/2008

**Date of final enrolment**

30/05/2008

**Locations****Countries of recruitment**

Switzerland

**Study participating centre**

**Inselspital University Hospital Bern**

Department of Nephrology and Hypertension

Bern

Switzerland

3010

**Sponsor information****Organisation**

Inselspital University Hospital Bern (Switzerland)

**ROR**

<https://ror.org/01q9sj412>

**Funder(s)**

**Funder type**

Other

**Funder Name**

Investigator initiated and funded

**Results and Publications**

Individual participant data (IPD) sharing plan

**IPD sharing plan summary**

Stored in repository