A new integrative weaning index of discontinuation from mechanical ventilation

Submission date	Recruitment status No longer recruiting	Prospectively registered	
02/02/2009		Protocol	
Registration date	Overall study status	Statistical analysis plan	
10/02/2009	Completed	[X] Results	
Last Edited	Condition category	Individual participant data	
29/03/2010	Respiratory		

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Sergio Nemer

Contact details

Hospital de Clínicas de Niterói Rua La Sale, 12 Centro, Niterói Rio De Janeiro Brazil 24020-090 snnemer@urbi.com.br

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

A new integrative weaning index of discontinuation from mechanical ventilation: an observational study

Study objectives

A new integrative weaning index can predict the weaning outcome better than traditional indexes.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committees of Hospital de Clínicas de Niterói approved in August 2004 (ref: USP/FM/SBD-033/07)

Study design

Observational cohort blinded (physician) trial

Primary study design

Observational

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Mechanical ventilation weaning

Interventions

Patients were classified as weaned or not weaned according to the ability to sustain spontaneous breathing for more than 48 hours post-extubation. Continuous positive airway pressure (CPAP) of 5 cm H2O was provided to avoid reconnection to the ventilator where necessary. Patients that remained in non invasive ventilation (NIV) were considered as 'weaning in progress' and could not be classified as weaned or not weaned. Patients that presented with post-extubation respiratory failure received all care to avoid reintubation but were ventilated by NIV where necessary.

The indexes were measured by the respiratory physiotherapists before the spontaneous breathing trials (SBTs). The decision to return to mechanical ventilation was made by the physician in charge (who was completely blind about the results of the indexes evaluated), based on the signs of poor tolerance.

The following measurements were taken for each weaning index:

- 1. Rapid shallow breathing index (f/Vt ratio)
- 2. Quasi-static compliance of the respiratory system (Cqst,rs)
- 3. Tidal volume
- 4. Respiratory rate
- 5. PaO2/FiO2 ratio
- 6. Volume minute
- 7. Airway occlusion pressure (P 0.1)
- 8. Integrative weaning index (Cqst,rs x arterial oxygen saturation / f/Vt ratio). This is a newly proposed index, used in some hospitals in Brazil.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Better accuracy in predicting weaning outcome

Key secondary outcome(s))

Adequate prognosis

Completion date

01/01/2008

Eligibility

Key inclusion criteria

Patients (both males and females) more than 24 hours in mechanical ventilation in weaning process.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

- 1. Patients younger than 18 years old
- 2. Neurological and neuromuscular diseases
- 3. Patients who have been tracheostomised

Date of first enrolment

01/09/2004

Date of final enrolment

01/01/2008

Locations

Countries of recruitment

Brazil

Study participating centre

Hospital de Clínicas de Niterói

Rio De Janeiro Brazil 24020-090

Sponsor information

Organisation

Niterói Hospital (Hospital de Clínicas de Niterói) (Brazil)

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Niterói Hospital (Hospital de Clínicas de Niterói) (Brazil)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	results	01/07/2009	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes