Progesterone in recurrent miscarriages (PROMISE) study

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
10/03/2009		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
17/03/2009		[X] Results		
Last Edited	Condition category	[] Individual participant data		
31/05/2016	Pregnancy and Childbirth			

Plain English summary of protocol

Background and study aims

This study was undertaken to test whether giving the hormone progesterone to pregnant women with a history of repeated unexplained early pregnancy losses could increase the number of pregnancies leading to live births after at least 24 weeks of gestation.

Who can participate?

Women aged 18-39 with unexplained recurrent miscarriages (three or more consecutive or non-consecutive first trimester miscarriages). A pregnancy loss is considered to be unexplained if conditions known to increase the risk of miscarriage are absent.

What does the study involve?

Participants were randomly allocated to one of two groups: one group received progesterone twice daily as vaginal pessaries and the other group received a placebo (dummy) with an identical appearance, from soon after a positive urinary pregnancy test, and no later than 6 weeks of pregnancy, until 12 completed weeks of pregnancy (or earlier if the pregnancy ended before 12 weeks).

What are the possible benefits and risks of participating?

There is substantial evidence from IVF practice that progesterone supplementation is safe to the mother and fetus at the proposed dose. Moreover, recent studies of vaginal progesterone in the context of prevention of preterm birth have not shown any evidence of short-term safety concerns in the participants.

Where is the study run from? 45 hospitals in the UK and the Netherlands

When is the study starting and how long is it expected to run for? From May 2009 to September 2014

Who is funding the study? NIHR Health Technology Assessment Programme - HTA (UK) Who is the main contact?

- 1. Dr Arri Coomarasamy (a.coomarasamy@bham.ac.uk)
- 2. Dr Rajendra Rai (r.rai@imperial.ac.uk)

Contact information

Type(s)

Scientific

Contact name

Dr Arri Coomarasamy

ORCID ID

https://orcid.org/0000-0002-3261-9807

Contact details

3rd Floor, Academic Department Birmingham Women's Hospital Metchley Park Road Edgbaston Birmingham United Kingdom B15 2TG +44 (0)121 623 6835 a.coomarasamy@bham.ac.uk

Type(s)

Scientific

Contact name

Dr Rajendra Rai

Contact details

Department of Obstetrics & Gynaecology St Mary's Hospital Mint Wing South Wharf Road London United Kingdom W2 1PG +44 (0)207 886 2475 r.rai@imperial.ac.uk

Additional identifiers

Protocol serial number HTA 08/38/01

Study information

Scientific Title

First trimester progesterone therapy in women with a history of unexplained recurrent miscarriages: a randomised double-blind placebo-controlled multi-centre trial (The PROMISE [PROgesterone in recurrent MIScarriagE] Trial)

Acronym

PROMISE

Study objectives

- 1. In women with unexplained recurrent miscarriages, progesterone (400 mg pessaries, twice daily), started soon as possible after a positive pregnancy test (and no later than 6 weeks gestation) and continued to 12 weeks of gestation, compared to placebo, increases live births beyond 24 completed weeks by at least 10% (principal objective)
- 2. Progesterone improves secondary outcomes such as gestation at delivery, on-going pregnancy at 12 weeks, survival at 28 days of neonatal life
- 3. Progesterone, compared to placebo, does not incur substantial adverse effects to the mother or the neonate
- 4 Explore differential or subgroup effects of progesterone in prognostic subgroups
- 5. Perform an economic evaluation for cost-effectiveness

More details can be found at http://www.nets.nihr.ac.uk/projects/hta/083801 Protocol can be found at http://www.nets.nihr.ac.uk/__data/assets/pdf_file/0018/52911/PRO-08-38-01.pdf

Ethics approval required

Old ethics approval format

Ethics approval(s)

REC: West Midlands Research Ethics Committee, 19/10/2009, ref: 09/H1208/44

Study design

Randomised double-blind placebo-controlled multi-centre trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Unexplained recurrent miscarriages

Interventions

Intervention group: Progesterone pessaries (400 mg twice daily) started soon as possible after a positive pregnancy test (and no later than 6 weeks gestation) and continued to 12 weeks of gestation

Control group: Placebo

Total duration of follow-up per participant: 42 weeks

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Progesterone

Primary outcome(s)

Live births beyond 24 weeks

Key secondary outcome(s))

- 1. Gestation at delivery
- 2. Clinical pregnancy at 6-8 weeks
- 3. Ongoing pregnancy at 12 weeks (range 11-13 weeks)
- 4. Miscarriage rate
- 5. Survival at 28 days of neonatal life
- 6. Congenital abnormalities with specific examination for genital anomalies
- 7. Adverse events

Completion date

01/09/2014

Eligibility

Key inclusion criteria

- 1. Women with unexplained recurrent miscarriages (3 or more consecutive first trimester miscarriages)
- 2. Age 18-39 years at randomisation (likelihood of miscarriages due to chromosomal aberrations is higher in older women; such miscarriages are unlikely to be prevented by progesterone therapy)
- 3. Spontaneous conception (as confirmed by urinary pregnancy tests)
- 4. Willing and able to give informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

39 years

Sex

Female

Key exclusion criteria

- 1. Inability to conceive spontaneously within 1 year of recruitment
- 2. Antiphospholipid syndrome (lupus anticoagulant and/or anticardiolipin antibodies [IgG or IgM]); other recognised thrombophilic conditions (testing according to usual clinic practice)
- 3. Intrauterine abnormalities (as assessed by ultrasound, hysterosonography, hysterosalpingogram, or hysteroscopy)
- 4. Fibroids distorting uterine cavity
- 5. Abnormal parental karyotype
- 6. Other identifiable causes of recurrent miscarriages (tests initiated only if clinically indicated) e.
- g., diabetes, thyroid disease and systemic lupus erythematosus (SLE)

Date of first enrolment

01/06/2010

Date of final enrolment

01/10/2013

Locations

Countries of recruitment

United Kingdom

Netherlands

Study participating centre
45 hospitals in the UK and the Netherlands

_

Sponsor information

Organisation

Imperial College London (UK)

ROR

https://ror.org/041kmwe10

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme

Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	26/11/2015		Yes	No
Results article	results	01/05/2016		Yes	No
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes