Microparticle orientated risk evaluation and prediction of pre-eclampsia among risk pregnancies

Submission date	Recruitment status	Prospectively registered
02/04/2009	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
28/07/2009	Completed	Results
Last Edited	Condition category	Individual participant data
28/07/2009	Pregnancy and Childbirth	Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Justine Sayuri Fitzgerald

Contact details

Abteilung für Geburtshilfe der Universitätsfrauenklinik Jena Bachstr. 18 Jena Germany 07743 +49 (0)3641 514 320 fitzgerald@med.uni-jena.de

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

FI 1635/1-1

Study information

Scientific Title

Microparticle Orientated Risk Evaluation and Prediction of Pre-eclampsia Among Risk gravidas: a multicentre prospective phase I prognostic marker study

Acronym

MORE PrePARd

Study objectives

The objective is to elucidate whether serum microparticle concentrations of pregnant women with abnormal uterine perfusion (a high risk group), both measured at 20th week of gestation, can discriminate between women who will develop preeclampsia (case group) and those who will not (control group).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Local ethics committee (Ethik-Kommission der Universitätsklinikum Jena) approved on the 26th August 2008 (ref: 2370-08/08)

Study design

Multicentre prospective blinded phase I prognostic marker pilot study

Primary study design

Interventional

Secondary study design

Non randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please contact fitzgerald@med.uni-jena.de or ekkehard. schleussner@med.uni-jena.de to request a patient information sheet

Health condition(s) or problem(s) studied

Pre-eclampsia/pregnancy

Interventions

Total of 80 - 100 pregnant women with abnormal uterine perfusion, including 20 women who develop pre-eclampsia during pregnancy (cases) and 60 - 80 woman who do not develop pre-eclampsia (controls). Patients with risk pregnancies for preeclampsia will be recruited for marker measurement between 19th and 21st week of gestation.

Experimental test:

Measurement of syncytiotophoblast microparticle (STBM) concentration in maternal serum at time of study inclusion.

Reference test:

Assessment of objective clinical symptoms of preeclampsia according to the German Society of Obstetrics and Gynecology (DGGG)-diagnostic criteria at 8-week intervals:

- 1. Hypertension defined as blood pressure (RR) systolic greater than or equal to 140 mmHg and /or diastolic greater than or equal to 90 mmHg, and
- 2. Proteinuria: Dipstick-Test greater than or equal to 1+ or greater than or equal to 300 mg total protein in 24-hour urine collection

There are two optional visits between the 19th and 21st week of gestation and birth: first visit at 27th to 29th week of gestation and second visit at 35th to 37th week of gestation to monitor these risk pregnancies. Final outcome will be assessed at 6 weeks post-partum.

Intervention Type

Other

Phase

Phase I

Primary outcome measure

Development of pre-eclampsia during current pregnancy. Final outcome will be assessed 6 weeks post-partum either by study doctor or attending doctor.

Secondary outcome measures

Pre-eclampsia associated complications:

- 1. Differentiation early onset/severe and late onset/mild pre-eclampsia
- 2. Intra-uterine growth restriction (IUGR) or small for gestational age (SGA)
- 3. Induction of preterm labour/delivery
- 4. Intra-uterine foetal demise (IUFD)
- 5. Placental abruption

Final outcome will be assessed 6 weeks post-partum either by study doctor or attending doctor.

Overall study start date

20/04/2009

Completion date

20/05/2009

Eligibility

Key inclusion criteria

- 1. Abnormal uterine artery Doppler ultrasound
- 2. Singleton pregnancy
- 3. Appropriate-for-gestational-age foetus
- 4. 19th 21st week of gestation
- 5. Healthy, normotensive gravidas
- 6. Women, all ages considered normal-risk pregnancy: 18 40 years

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Female

Target number of participants

80 -100 pregnant women

Key exclusion criteria

- 1. Concurrent participation in interventional clinical studies
- 2. Multiple pregnancy
- 3. Premature rupture of membrane (PROM)
- 4. Suspected/diagnosed infected amnion syndrome (IAS)
- 5. Pre-term labour/cervical incompetence
- 6. Foetal genopathia
- 7. Suspected foetal defects
- 8. Pre-existing maternal disease, e.g. diabetes mellitus, cardiovascular or renal disease

Date of first enrolment

20/04/2009

Date of final enrolment

20/05/2009

Locations

Countries of recruitment

Germany

Study participating centre Abteilung für Geburtshilfe der Universitätsfrauenklinik Jena

Jena Germany 07743

Sponsor information

Organisation

University Hospital of Jena (Universitätsklinikum Jena) (Germany)

Sponsor details

Prof. Dr. med. Ekkehard Schleussner Abteilung Geburtshilfe Klinik für Frauenheilkunde und Geburtshilfe Bachstr. 18 Jena Germany 07743 +49 (0)3641 9 33230 ekkehard.schleussner@med.uni-jena.de

Sponsor type

Hospital/treatment centre

Website

http://www.geburtshilfe.uniklinikum-jena.de

ROR

https://ror.org/035rzkx15

Funder(s)

Funder type

Research council

Funder Name

German Research Council (Deutsche Forschungsgemeinschaft [DFG]) (Germany) (ref: FI 1635/1-1)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration