

A pilot study to assess flutamide monotherapy compared With maximal androgen blockade in metastatic prostate cancer

Submission date 19/08/2002	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 19/08/2002	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 29/10/2019	Condition category Cancer	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr - -

Contact details
UKCCCR Register Co-ordinator
MRC Clinical Trials Unit
222 Euston Road
London
United Kingdom
NW1 2DA

Additional identifiers

Protocol serial number
SCTO46

Study information

Scientific Title
A pilot study to assess flutamide monotherapy compared With maximal androgen blockade in metastatic prostate cancer

Study objectives

Not provided at time of registration

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Prostate cancer

Interventions

1. Group A: Oral flutamide 250 mg three times daily
2. Group B: Oral flutamide 250 mg three times daily plus either orchidectomy or the lutenizing hormone releasing hormone (LHRH) analogue Zoladex (goserelin acetate). Zoladex is given by monthly injection at the dosage recommended by the manufacturer

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Flutamide, goserelin acetate

Primary outcome(s)

Not provided at time of registration

Key secondary outcome(s))

Not provided at time of registration

Completion date

30/09/1996

Eligibility**Key inclusion criteria**

1. Histologically confirmed prostate carcinoma
2. Skeletal metastases on bone scan, with radiological conformation if appropriate, or typical

sclerotic metastases on X-ray

3. Life expectancy >6 months

4. Suitable for treatment by any of the study therapies

5. No previous hormonal therapy

6. Not currently receiving corticosteroids, spironolactone or aminoglutethamide

7. Adequate renal and hepatic function

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Male

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/01/1994

Date of final enrolment

30/09/1996

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

MRC Clinical Trials Unit

London

United Kingdom

NW1 2DA

Sponsor information

Organisation

Schering-Plough Ltd (UK)

ROR

<https://ror.org/00148fb49>

Funder(s)

Funder type

Industry

Funder Name

Schering-Plough Ltd (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes