

A national patient registry for radiofrequency ablation (RFA) for Barrett's oesophagus

Submission date 12/06/2013	Recruitment status Recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 14/06/2013	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 08/02/2021	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

<http://www.cancerresearchuk.org/cancer-help/trials/a-trial-looking-new-treatment-2-pre-cancerous-conditions-affecting-food-pipe-halo>

Contact information

Type(s)

Scientific

Contact name

Dr Laurence Lovat

Contact details

University College London Hospitals NHS Trust

235 Euston Square

London

United Kingdom

NW1 2BU

-

l.lovat@uclh.nhs.uk

Additional identifiers

Protocol serial number

Protocol Version 3.1

Study information

Scientific Title

Radiofrequency ablation of Barrett's columnar lined oesophagus and squamous dysplasia: multicentre trial with long term follow up using a central database

Study objectives

The aim of this prospective, multi-centre patient registry study is to provide a tool for participating physician investigators to collect outcome data related to the use of the HALO Ablation Systems. We hypothesise that RFA is safe and effective in treating patients with dysplasia arising in Barrett's oesophagus and will maintain a durable response.

Ethics approval required

Old ethics approval format

Ethics approval(s)

National Research Ethics Service (NRES), ref: 08/H0714/27

Study design

Prospective multi-centre patient registry

Primary study design

Observational

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Oesophageal Cancer/ Barrett's Oesophagus

Interventions

HALO radiofrequency ablation (HALO RFA)

The study collects the outcomes of patients undergoing the intervention which is RFA. The study collects not just demographic data on patients eligible for RFA treatment but also the effectiveness of the treatment in curing them of early oesophageal neoplasia. The study collates the number of procedures required to get to this point but also examines the reasons why we fail in a minority of patients so we can try understand these patients to help improve our practice and streamline our approach to them.

Intervention Type

Procedure/Surgery

Primary outcome(s)

1. Endoscopic clearance rate for Barrett's esophagus. Percentage of patients with no endoscopically visible Barrett's oesophagus at 1 year follow-up
2. Histological clearance rate for intestinal metaplasia. Percentage of patients with no histological evidence of intestinal metaplasia at 1 year follow-up
3. Histological clearance rate for columnar dysplasia. Percentage of patients with baseline dysplasia who have no histological evidence of dysplasia at 1 year follow-up
4. Sub-squamous intestinal metaplasia. Percentage of patients at 1 year follow-up with sub-squamous intestinal metaplasia, with or without dysplasia, that is covered completely by an intact layer of squamous epithelium with no communication with the surface

Key secondary outcome(s)

Ability of aneuploidy and other molecular changes to predict success or failure of this therapy.

Completion date

31/03/2028

Eligibility

Key inclusion criteria

1. Patients will be recruited from those referred for ablative management of Barrett's oesophagus or squamous dysplasia with HALO RFA
2. Patients must have no contraindications to endoscopy
3. Males and non-pregnant females over the age of 21 years. Female patients who are premenopausal must practice a medically acceptable form of contraception.
4. Patients must sign an informed consent form

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

21 years

Sex

All

Key exclusion criteria

1. Presence of invasive carcinoma of the oesophagus
2. Patients in whom endoscopy is contraindicated
3. Patients with oesophageal varices
4. Previous radiotherapy
5. Patients who have undergone Hellers myotomy
6. Pregnant females
7. People under the age of 21 years

Date of first enrolment

31/03/2008

Date of final enrolment

30/09/2027

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
University College London Hospitals NHS Trust
London
United Kingdom
NW1 2BU

Sponsor information

Organisation

Joint UCLH/UCL Biomedical Research Unit (UK)

ROR

<https://ror.org/02jx3x895>

Funder(s)

Funder type

Industry

Funder Name

BAARX Medical Inc. (USA)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/08/2015		Yes	No
Results article	results	05/02/2021	08/02/2021	Yes	No