

Airogym exercise on solute removal

| | | |
|--|--|--|
| Submission date 11/08/2016 | Recruitment status No longer recruiting | <input type="checkbox"/> Prospectively registered |
| Registration date 19/09/2016 | Overall study status Completed | <input type="checkbox"/> Protocol |
| Last Edited 15/09/2016 | Condition category Urological and Genital Diseases | <input type="checkbox"/> Statistical analysis plan |
| | | <input type="checkbox"/> Results |
| | | <input type="checkbox"/> Individual participant data |
| | | <input type="checkbox"/> Record updated in last year |

Plain English summary of protocol

Background & Study aims

Kidney dialysis is a medical procedure that removes waste products and excess fluid from the blood in people whose kidneys have stopped working properly, namely patients with chronic kidney disease (CKD) or end stage kidney failure. The procedure is expensive and there is limited availability of these sorts of expensive therapies in the majority of emerging countries and more so in African nations. The aim of this study is to find out whether exercise during haemodialysis (the most common type of kidney dialysis) has an effect on solute (i.e. waste product) removal from the blood and to identify whether there are any other possible benefits such as preventing deep vein thrombosis, improvement in blood circulation and reduction in swelling and edema (build-up of fluid in the tissues).

Who can participate?

End stage kidney disease patients aged between 30-50 years.

What does the study involve?

This study is called a Quasi-experimental investigation, since patients participating in study know whether they are in the intervention (test) group or the control group. They are randomly allocated to one of the groups. Patients in the intervention group exercise for fifteen minutes every hour using a Airogym exercise cushion. They exercise for a total of sixty minutes during each four-hour hemodialysis session. Patients in the control group do not have to exercise during their 4 hour hemodialysis sessions. Blood is taken from all patients in both groups once a month before and after the session. They are also all encouraged to exercise regularly, eat a healthy diet, get involved in social activities and attend psychological counselling (if necessary) to maintain their quality of life during the study.

What are the possible benefits and risks of participating?

No benefits are expected for the participants, but this study will improve the future treatment of end-stage kidney disease patients. There is no additional risk involved in this study.

Where is the study run from?

1. Newcastle Private Hospital Renal unit (South Africa)
2. Mediclinic Bloemfontein (South Africa)

Who is funding the study?
Durban University of Technology, Durban (South Africa)

Who is the main contact?
Professor Jamila Khatoon Adam

Contact information

Type(s)
Scientific

Contact name
Prof Jamila Khatoon Adam

ORCID ID
<https://orcid.org/0000-0001-6266-699X>

Contact details
Department of Biomedical & Clinical Technology
41-43 M L Sultan Road
Durban University of Technology
Durban
South Africa
4000

Additional identifiers

Protocol serial number
4426

Study information

Scientific Title
Effect of exercise on solute removal and edema on end stage renal disease patients

Acronym
Airogym exercise

Study objectives

1. Patients participating in the intervention group will show marked reduction in edema compared to the control group during a nine- month period of study
2. Patients participating in the intervention group will show a marked increase in solute removal compared to patients in the control group

Ethics approval required
Old ethics approval format

Ethics approval(s)
Institutional Research Ethics Committee, Durban University of Technology, 30/08/2006

Study design

Unblinded randomised controlled trial

Primary study design

Interventional

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Renal artery obstruction

Interventions

Between September 2006 to December 2006, 34 end stage renal patients for the study were recruited from two MediClinic Hospital Renal Units (namely, Newcastle Private Hospital Renal Unit and Bloemfontein MediClinic Renal Units, South Africa) where they have been on maintenance hemodialysis for at least 3 months.

Participants are randomly allocated to one of two groups.

1. The intervention group: this group exercised on a Airogym exercise cushion for 15 minutes every hour over a 4 hour dialysis session. They exercise for a total of 60 minutes over the 4 hour period.
2. The control group: this group do not have to exercise during their 4 hour dialysis session.

During the first three months of this study patients in the intervention group did not exercise on the exercise cushion (Airogym) to establish a baseline. Thereafter, from the fourth month until the ninth month patients in the intervention group pedalled on the exercise cushion during hemodialysis. All patients exercised at a slow steady rate to target a heart rate of 100 beats per minute and then slowed down gradually. A total of sixty minutes of exercise was performed over the four hours on hemodialysis. Urea, creatinine and potassium were quantitatively measured using standard laboratory techniques from patients (pre and post hemodialysis). Edema of the lower limb was measured in centimeters around the ankles (right and left) before and after dialysis.

Randomisation Method: Random allocation will be utilised to assign the patients to their respective groups, with the patients drawing a letter (A or B) out of an envelope. A total of twenty patients will be recruited from Bloemfontein (ten in group A and ten in group B) and fourteen from Newcastle (seven in group A and seven in group B)

Follow up of patients: On completion of the study, the patients recruited continued on chronic dialysis programme as was recommended by the nephrologist.

The Airogym:

The Airogym was designed specifically to address two common problems associated with long haul flying, i.e., ankle swelling and blood clotting. Patients with end-stage renal disease on haemodialysis are generally more prone to developing deep vein thrombosis, swelling and oedema due to having chronic renal failure. They often feel fatigued and are unable to join an exercise programme at the gymnasium or do any strenuous activities. This inactivity further increases the risk of developing deep vein thrombosis, swelling and oedema due to poor blood circulation. Both can be attributed to stagnation of the circulation. Anybody sitting or standing for long periods of time will notice ankle swelling and pain in the lower legs. Fortunately in the

majority of passengers the clots dissolve without causing any apparent symptoms or signs, and leaving no long-term damage. On occasion the blood clot can grow and then a segment can break off and travel to the lungs. There are two problems associated with blood clots, one a clot in the lung, the other an effect which may take many years to manifest itself, namely the development of the post thrombotic limb. The airogym was designed specifically to promote the flow of blood through the deep veins. By pressing down on the footpad, veins in the foot are compressed, squeezing blood into the main veins in the calf. The pressure involved in squeezing the foot causes muscular contractions of the calf muscles promoting the flow of blood through the main veins. Results showed an improvement in peak velocity blood flow up to five times. The device prevented venous stasis and reduced the risk of developing blood clots and also prevented ankle swelling.

Intervention Type

Device

Primary outcome(s)

Patients experience of the exercise program, measured using questionnaires completed once a month over a nine month period.

Key secondary outcome(s)

1. Urea, creatinine and potassium levels, measured using biochemical analysis of blood samples, taken once a month for nine months from each participant before the dialysis session and afterwards
2. Degree of edema of the lower limb, measured in centimeters around the ankles (right and left) before and after dialysis
3. Statistical Analysis. Paired student's t-test was used to determine level of significance, using the statistical package SPSS 9.0 for Windows

Completion date

30/09/2007

Eligibility

Key inclusion criteria

1. On a chronic hemodialysis schedule
2. Hemodynamically stable
3. Attending haemodialysis at the Bloemfontein Mediclinic Hospital or Newcastle Private Hospital Renal Unit
4. Receiving haemodialysis twice or three times a week
5. Eighteen years old and above

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Patients not on a chronic haemodialysis schedule
2. Patients medically unfit to participate (confirmed by doctor's incharge of renal unit patients)
3. Patients not willing to comply with protocol
4. Patients that are diabetic

Date of first enrolment

20/09/2006

Date of final enrolment

30/12/2006

Locations**Countries of recruitment**

South Africa

Study participating centre**Newcastle Private Hospital Renal unit**

Hospital St, Newcastle

Durban

South Africa

2940

Study participating centre**Mediclinic Bloemfontein**

Cnr Kellner & Parfitt Street

Westdene

Bloemfontein, Free State

South Africa

9301

Sponsor information**Organisation**

Durban University of Technology

ROR

<https://ror.org/0303y7a51>

Funder(s)

Funder type

University/education

Funder Name

Durban University of Technology

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not expected to be made available