

# A randomised trial of infusional 5-fluorouracil and concomitant radiotherapy in locally advanced rectal cancer, comparing three and six months of infusional 5-fluorouracil

<b>Submission date</b> 19/08/2002	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 19/08/2002	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 21/01/2019	<b>Condition category</b> Cancer	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr - -

**Contact details**  
UKCCCR Register Co-ordinator  
MRC Clinical Trials Unit  
222 Euston Road  
London  
United Kingdom  
NW1 2DA

## Additional identifiers

**Protocol serial number**  
COLORAD

## Study information

**Scientific Title**

A randomised trial of infusional 5-fluorouracil and concomitant radiotherapy in locally advanced rectal cancer, comparing three and six months of infusional 5-fluorouracil

### **Study objectives**

Not provided at time of registration

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Not provided at time of registration

### **Study design**

Randomised controlled trial

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Cancer of rectum

### **Interventions**

All patients receive continuous infusional 5-fluorouracil for 12 weeks plus radiotherapy to the pelvis, minimum 45 Gy given in twenty-five fractions over 5 weeks to start at the beginning of the 5th week of chemotherapy.

Depending upon the initial randomisation, patients receive either:

1. Group One: Continuous infusional 5-fluorouracil for a further 12 weeks.
2. Group Two: No further chemotherapy with 5-fluorouracil.

### **Intervention Type**

Drug

### **Phase**

Not Applicable

### **Drug/device/biological/vaccine name(s)**

5-fluorouracil

### **Primary outcome(s)**

Not provided at time of registration

### **Key secondary outcome(s))**

Not provided at time of registration

### **Completion date**

11/11/1994

# Eligibility

## Key inclusion criteria

1. Histologically verified adenocarcinoma of the rectum, which is either:
  - 1.1. Inoperable primary tumour
  - 1.2. Locally recurrent tumour
  - 1.3. Residual pelvic disease after resection as assessed either surgically or histopathologically
2. No evidence of metastatic disease
3. No past history of malignancy
4. No previous radiotherapy to the pelvis
5. Normal bone marrow, liver and renal function
6. No concurrent severe or life threatening illness

## Participant type(s)

Patient

## Healthy volunteers allowed

No

## Age group

Not Specified

## Sex

Not Specified

## Key exclusion criteria

Not provided at time of registration

## Date of first enrolment

01/01/1990

## Date of final enrolment

11/11/1994

# Locations

## Countries of recruitment

United Kingdom

England

## Study participating centre

**MRC Clinical Trials Unit**

London

United Kingdom

NW1 2DA

# Sponsor information

## Organisation

The Royal Marsden NHS Foundation Trust (UK)

## ROR

<https://ror.org/0008wzh48>

# Funder(s)

## Funder type

Hospital/treatment centre

## Funder Name

Royal Marsden Hospital (UK)

# Results and Publications

## Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration