

# A clinical trial to assess the effectiveness of miniplate surgical treatment to bring the upper jaw forwards in 11-14 years old children

<b>Submission date</b> 02/09/2014	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 19/09/2014	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 29/10/2024	<b>Condition category</b> Oral Health	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Children with a lower jaw protruding forward of the upper jaw have what is called a class III skeletal pattern. In some cases, it is caused by the upper jaw being set too far back. The treatment in such cases is to bring the upper jaw forward to correct the reverse bite. Patients are not usually offered corrective treatment until their general and facial growth is complete at around 17-18 years of age, at which point they may be offered orthognathic (jaw) surgery to correct the position of the jaw. The surgery involves a controlled break of the upper and lower jaw from inside the mouth which are then screwed into place to hold the jaws into their new position. There is an early treatment option, for children under the age of 10 that may prevent them from having to undergo the surgery when they are older. This means wearing reverse headgear, which brings the top jaw and teeth forwards over time and has a 70% success rate, but this option only works for children under the age of 10. There is, however, another option for 11-14 year olds. This is called Bone Anchored Maxillary Protection (BAMP). It involves places small metal plates (miniplates) from inside the mouth into the front part of the cheekbone next to the upper molars on each side. Further miniplates are positioned near the front of the lower jaw. Elastics are then attached from the upper to the lower plates, via hooks, to bring the upper jaw and teeth forwards and correct the class III skeletal pattern. At the moment, BAMP is not used routinely used in the UK. Here, we want to see how well it performs compared with a no treatment (control group) to see whether it reduces the need for orthognathic surgery at a later date.

### Who can participate?

Children aged between 11-14, with a class III skeletal pattern and a reverse bite.

### What does the study involve?

Participants are randomly allocated into one of two groups. Those in group 1 undergo BAMP surgery. Those in group 2 are given no treatment. The success of the treatment will be assessed according to how much the upper jaw moves to correct the reverse bite and whether orthognathic surgery is needed at a later date. All participants are followed up at 1 year and 3 years after the start of the trial to compare the outcomes for each group.

What are the possible benefits and risks of participating?

Possible benefits of BAMP include avoiding orthognathic surgery at a later date. The treatment is also considerably cheaper than orthognathic surgery and is a 30 minute daycase procedure rather than a treatment requiring a 3 day stay in hospital.

Where is the study run from?

The trial is multicentre with proposed sites: Tameside Hospital NHS Foundation Trust (Lead site), University Hospital of south Manchester (Wythenshawe Hospital), Bradford Royal Infirmary, Sheffield Dental Hospital, Newcastle Dental Hospital, Peterborough General Hospital, Southend Hospital, Kent and Canterbury Hospital (UK)

When is the study starting and how long is it expected to run for?

January 2015 to October 2022

Who is funding the study?

DB Orthodontics (UK)

Who is the main contact?

Dr Nicola Mandall

(see contact details below)

## Contact information

### Type(s)

Scientific

### Contact name

Dr Nicola Mandall

### Contact details

Orthodontic Department  
Tameside Hospital NHS Foundation Trust  
Fountain Street  
Ashton under Lyne  
Lancashire  
Lancashire  
United Kingdom  
OL6 9RW

## Additional identifiers

EudraCT/CTIS number

IRAS number

160532

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

## Study information

### Scientific Title

A multicentre randomised clinical trial to investigate whether Bone Anchored Maxillary Protraction (BAMP) reduces the need for orthognathic (facial) surgery

### Acronym

BAMP

### Study objectives

Null hypothesis: There is no difference in the effectiveness of BAMP versus no treatment /control group in terms of: need for later orthognathic facial surgery, skeletal forward movement of the maxilla (upper jaw), correction of reverse overjet, facial profile improvement.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

NRES committee North West Greater Manchester Central, 19/02/2015, ref: 15/NW/0022

### Study design

Multicentre randomized controlled trial

### Primary study design

Interventional

### Secondary study design

Randomised controlled trial

### Study setting(s)

Hospital

### Study type(s)

Treatment

### Participant information sheet

Not available in web format, please use the contact details to request a patient information sheet

### Health condition(s) or problem(s) studied

Oral and maxillofacial surgery and orthodontics

### Interventions

Group 1: BAMP surgery

Group 2: Control/no treatment

BAMP consists of small metal plates that are inserted from inside the mouth into the front part of the cheekbone bone next to the upper molars on each side. Further miniplates are positioned

into the bone down in the lower canine region near the front of the lower jaw. Elastics are then attached from the upper to the lower plates, via hooks, to bring the upper jaw and teeth forwards and correct the class III skeletal pattern. The miniplates are placed and later removed as a daycase general anaesthetic procedure and the patient wears the elastics for around 6-8 months.

## **Intervention Type**

Procedure/Surgery

## **Primary outcome measure**

1. Need for orthognathic surgery later
2. Amount of forward movement of the maxilla (upper jaw) to correct the reverse bite
3. Reverse overjet (bite) correction

Need for orthognathic surgery will be measured through consultant clinical panel consensus method using patient records. Forward movement of the maxilla will be measured from lateral cephalogram radiographs. Reverse overbite correction will be measured from plaster study models of the patients teeth using a metal ruler. The outcomes will be measured at baseline, 1 and 3 years follow-up.

## **Secondary outcome measures**

1. Facial aesthetic improvement
2. Patient discomfort log following surgery
3. Operating time for the BAMP surgery
4. BAMP - post operative infection rates and plate failure rates

Facial aesthetic improvement will be measured using panel consensus of facial profiles photographs. Patient discomfort log will be recorded on a 7 point Likert scale with 1 = no pain at all to 7 severe pain. Operating time for BAMP surgery will be recorded from the theatre time log. Post operative infection rates and plate failure rates will be recorded from the patient notes. The outcomes will be measured at baseline, 1 and 3 years follow-up.

## **Overall study start date**

02/01/2015

## **Completion date**

01/10/2022

# **Eligibility**

## **Key inclusion criteria**

1. Age 11-14 years old
2. Class III skeletal pattern with reverse bite of at least 1mm
3. Dentally fit
4. Written consent

## **Participant type(s)**

Patient

## **Age group**

Child

**Lower age limit**

11 Years

**Upper age limit**

14 Years

**Sex**

Both

**Target number of participants**

60

**Total final enrolment**

57

**Key exclusion criteria**

1. Cleft lip and or palate or craniofacial syndrome
2. Reverse bite greater than 6mm
3. Lower jaw asymmetric
4. Muscular dystrophy (weakness)
5. Patient smokes

**Date of first enrolment**

06/07/2015

**Date of final enrolment**

01/10/2019

## **Locations**

**Countries of recruitment**

England

United Kingdom

**Study participating centre**

**Tameside Hospital NHS Foundation Trust**

Lancashire

United Kingdom

OL6 9RW

## **Sponsor information**

**Organisation**

Tameside Hospital NHS Foundation Trust (UK)

**Sponsor details**

c/o Mrs Margaret Cooper  
Tameside Hospital NHS Foundation Trust R+D department  
Fountain Street  
Ashton under Lyne  
Lancashire  
England  
United Kingdom  
OL6 9RW

**Sponsor type**

Hospital/treatment centre

**ROR**

<https://ror.org/04d713p41>

**Funder(s)****Funder type**

Industry

**Funder Name**

DB Orthodontics (UK)

**Results and Publications****Publication and dissemination plan**

Current publication and dissemination plan:  
Submitted to the Journal of Orthodontics in August 2023

Previous publication and dissemination plan:  
To be confirmed at a later date

**Intention to publish date**

01/10/2024

**Individual participant data (IPD) sharing plan**

The datasets generated during and/or analysed during the current study are not expected to be made available. The data will be held at Tameside and Glossop Integrated Care NHS Foundation Trust and copied to the R+D dept for electronic archiving.

**IPD sharing plan summary**

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">HRA research summary</a>			28/06/2023	No	No
<a href="#">Results article</a>		06/06/2024	29/10/2024	Yes	No