

Pre-exposure option for reducing HIV in the UK: an open-label randomisation to immediate or deferred daily Truvada for HIV-negative gay men

Submission date 28/02/2013	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 28/02/2013	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 13/12/2019	Condition category Infections and Infestations	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Human immunodeficiency virus (HIV) is a virus that attacks the immune system and weakens the body's ability to fight infections and disease. It is most commonly caught by having sex without a condom. The aim of this study is to assess what happens when a new approach, known as pre-exposure prophylaxis (PrEP), is added to the existing methods available to gay men in the UK to reduce their risk of catching HIV. Prophylaxis is taking a drug to prevent, or at least reduce the risk of, catching an infection. In this study a drug called Truvada is used, which is commonly used to treat HIV. It is a combination of tenofovir and emtricitabine in a single pill, to be taken once a day. A major concern about HIV-negative individuals taking PrEP is that they will decrease their use of other methods such as condoms, thereby increasing their risk of HIV and other sexually transmitted infections. This can only be assessed when people know they are taking an active drug. This study therefore offers people PrEP on an 'open label' basis, meaning that everyone who takes it knows they are taking the drug; no-one takes an inactive placebo pill.

Who can participate?

HIV-negative gay men aged 18 or over

What does the study involve?

In order to assess what happens when PrEP is used, one group of participants needs to be followed for a period before using PrEP, at a time when they are accessing the best available methods for reducing risk. To do this, participants are randomly allocated to be offered PrEP either immediately or after 12 months. All participants are offered other interventions to reduce their risk. They are followed up in clinic every 3 months for 2 years, and tested for HIV. A test for sexually transmitted infections is carried out every 6 months or more frequently if indicated. In between visits, participants are asked to complete a diary and monthly questionnaire reporting anal intercourse and their pill taking when they are on Truvada.

What are the possible benefits and risks of participating?

Not provided at time of registration

Where is the study run from?

Medical Research Council Clinical Trials Unit (MRC CTU) (UK)

When is the study starting and how long is it expected to run for?

November 2012 to October 2016

Who is funding the study?

1. Gilead Sciences (USA)
2. Health Protection Agency (HPA) (UK)
3. Medical Research Council (MRC) (UK)

Who is the main contact?

Dr Sheena McCormack

smc@ctu.mrc.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr Sheena McCormack

Contact details

MRC CTU

Aviation House

125 Kingsway

London

United Kingdom

WC2B 6NH

-

smc@ctu.mrc.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

2012-002373-56

ClinicalTrials.gov (NCT)

NCT02065986

Protocol serial number

13131

Study information

Scientific Title

Pre-exposure Option for reducing HIV in the UK: an open-label randomisation to immediate or Deferred daily Truvada for HIV-negative gay men

Acronym

PROUD

Study objectives

The aim of this study is to assess what happens when a new approach, known as pre-exposure prophylaxis (PrEP), is added to the existing methods available to gay men in the UK to reduce their risk of catching human immunodeficiency virus (HIV) infection.

Ethics approval required

Old ethics approval format

Ethics approval(s)

First MREC approval date 02/10/2012, ref: 12/LO/1289

Study design

Randomised interventional study

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Topic: Infection; Subtopic: Infection (all Subtopics); Disease: Infectious diseases and microbiology

Interventions

A major concern about HIV-negative individuals taking PrEP is that they will decrease their use of other methods such as condoms, thereby increasing their risk of HIV and other sexually transmitted infections. This can only be assessed when people know they are taking an active drug. This study will therefore offer people PrEP on an 'open label' basis, meaning that everyone who takes it will know they are taking the drug; no-one will take an inactive placebo pill. In order to assess what happens when PrEP is added, one group will need to be followed for a period before the PrEP is included, at a time when they are accessing the best available methods for reducing risk. To do this, the volunteers will be randomised to either be offered PrEP immediately or after 12 months. This study will use a drug called Truvada, which is commonly used to treat HIV. It is a combination of tenofovir and emtricitabine in a single pill, to be taken once a day. All participants will be offered other interventions to reduce their risk. They will be followed in clinic every 3 months for 2 years, and tested for HIV. A screen for sexually transmitted infections will be done every 6 months or more frequently if indicated. In between visits, participants will be asked to complete a diary and monthly questionnaire reporting anal intercourse and their pill taking when they are on Truvada.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Truvada

Primary outcome(s)

Time to accrual; Timepoint(s): time to accrual of 500 participants

Key secondary outcome(s)

No secondary outcome measures

Completion date

01/10/2016

Eligibility**Key inclusion criteria**

1. Born to male gender, age 18 years or more
2. Previously attended the enrolling clinic on at least one occasion
3. Completed a screen for HIV and sexually transmitted infections (STIs)
4. HIV negative by a routinely used assay within 4 weeks prior to or on the day of randomisation
5. Reported unprotected anal intercourse (UAI) on more than one occasion within the 90 days prior to randomisation
6. Likely, in the opinion of the volunteer, to have UAI in the next 90 days
7. Willing and able to comply with the visit schedule throughout the follow-up period
8. Willing and able to provide written informed consent

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Male

Key exclusion criteria

1. An acute viral illness that could be due to HIV seroconversion
2. Any contraindications to Truvada according to the current package insert
3. Treatment for hepatitis B infection indicated or ongoing
4. Unlikely, in the opinion of the clinician, to comply with the randomised allocation

Date of first enrolment

29/11/2012

Date of final enrolment

28/11/2013

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

MRC CTU

London

United Kingdom

WC2B 6NH

Sponsor information

Organisation

Medical Research Council Clinical Trials Unit (MRC CTU) (UK)

ROR

<https://ror.org/001mm6w73>

Funder(s)

Funder type

Industry

Funder Name

Gilead Sciences

Alternative Name(s)

Gilead, Gilead Sciences, Inc., Oligogen

Funding Body Type

Government organisation

Funding Body Subtype

For-profit companies (industry)

Location

United States of America

Funder Name

Health Protection Agency (HPA) (UK)

Alternative Name(s)

HPA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Funder Name

Medical Research Council (MRC) (UK)

Alternative Name(s)

Medical Research Council (United Kingdom), UK Medical Research Council, MRC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	02/01/2016		Yes	No
Results article	results	24/03/2016		Yes	No

Results article	results	10/12/2019	13/12/2019	Yes	No
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes