

Achilles Tendinopathy Management: platelet-rich plasma versus eccentric loading programme

Submission date 22/12/2009	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 18/02/2010	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 05/08/2014	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mr Matthew Costa

Contact details
Warwick Medical School
Clinical Sciences Research Institute
Clifford Bridge Road
Coventry
United Kingdom
CV2 2DX
matthew.costa@warwick.ac.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
PRF/09/2

Study information

Scientific Title

Achilles Tendinopathy Management: a randomised controlled trial comparing platelet-rich plasma with an eccentric loading programme

Acronym

ATM

Study objectives

There is no difference in Victorian Institute of Sports Assessment - Achilles (VISA-A) scores at 6 months between patients initially managed with platelet-rich plasma injections compared to patients managed with an eccentric loading programme.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Coventry Research Ethics Committee, 20/03/2009, ref: 09/H1210/18

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Achilles tendinopathy

Interventions

Intervention:

Platelet Rich Plasma Injection. The procedure will involve taking 52 ml of whole blood combined with 8 ml of an anti-coagulant which will be immediately centrifuged at 2400 for 12 minutes. After centrifugation the platelet layer (approximately 3 - 5 ml) will be extracted using a syringe and then injected into the Achilles tendinopathy.

Comparison:

Eccentric Loading Programme, involving two exercises. The first involves the patient being in a standing position with the heel over the edge of a step with the legs straight. The patient then slowly lowers their heels beyond the level of the step. The second exercise follows the same sequence but with the knee slightly bent, to maximise activation of the soleus muscles. These exercises are performed three times a day, 7 days a week for 12 weeks and are progressed as pain allows by adding weight via a back pack.

Secondary Sponsor Details:

University Hospitals of Coventry and Warwickshire NHS Trust
Research and Development Department
Clifford Bridge Road
Coventry CV4 8UW
United Kingdom

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

VISA-A at 6, 12, 24, 30, 36 and 52 weeks

Secondary outcome measures

EQ-5D and complications at 6, 12, 24, 30, 36 and 52 weeks

Overall study start date

01/02/2010

Completion date

01/02/2012

Eligibility

Key inclusion criteria

1. Midsubstance achilles tendinopathy diagnosed clinically through pain on palpation at a level of 2 - 6 cm above the tendon insertion and ultrasonography
2. The tendinopathy will be causing pain during loading activities and limit those activities
3. Duration of at least 3 months
4. Aged over 18 years, either sex

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

20

Key exclusion criteria

1. Tendinopathies secondary to systemic conditions such as rheumatoid arthritis and diabetes
2. Insertional Achilles tendinopathies
3. Pregnancy
4. Previous Achilles rupture or surgery
5. Dislocation or fracture of the lower limb within the preceeding 12 months

Date of first enrolment

01/02/2010

Date of final enrolment

01/02/2012

Locations**Countries of recruitment**

England

United Kingdom

Study participating centre

Warwick Medical School

Coventry

United Kingdom

CV2 2DX

Sponsor information**Organisation**

University of Warwick (UK)

Sponsor details

c/o Grants and Contracts Officer

Research Support Services

University House

Kirby Corner Road

Coventry

England

United Kingdom

CV4 7AL
N.K.Bains@warwick.ac.uk

Sponsor type
University/education

Website
<http://www2.warwick.ac.uk/>

ROR
<https://ror.org/01a77tt86>

Funder(s)

Funder type
Research organisation

Funder Name
Physiotherapy Research Foundation (UK) (ref: PRF/09/2)

Results and Publications

Publication and dissemination plan
Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	17/10/2013		Yes	No