

# Influence of the suturing technique on the wound healing of the palate after obtaining a connective tissue graft

<b>Submission date</b> 22/01/2018	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 15/02/2018	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 15/02/2018	<b>Condition category</b> Ear, Nose and Throat	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Gingival recession (also known as receding gums) is when the roots of the teeth are exposed when the gum tissue is lost. One of the main treatments is a gum transplant, where gum tissue (called a graft) is collected from other parts of the mouth and placed on the areas where the gums have receded. There are many different surgical techniques used described to obtain a connective tissue graft from the palate in order to decrease patient discomfort, but there is no clear answer about the influence of the suture technique and the residual flap thickness on the early healing of the donor site. The aim of this study is to examine if wound healing is related to suturing, and if palatal thickness is related to postoperative pain.

### Who can participate?

Adults aged 18 and older who have gingival recession.

### What does the study involve?

All participants undergo a procedure where a connective tissue graft is taken from the palate using a single incision technique. Participants are randomly allocated to one of two groups. Those in the first group have the donor site sutured with one method. Those in the second group receive a criss-cross suture used. The same drug therapy is given to all patients. One week after surgery, at suture removal, the healing of the donor site is evaluated.

### What are the possible benefits and risks of participating?

There are no direct benefits or risks taking part in the study.

### Where is the study run from?

Universitat internacional de Catalunya (Spain)

### When is the study starting and how long is it expected to run for?

September 2015 to May 2017

Who is funding the study?  
Universitat Internacional de Catalunya (Spain)

Who is the main contact?  
Dr Giovanni Maino (Public)

## Contact information

**Type(s)**  
Public

**Contact name**  
Dr Giovanni Maino

**Contact details**  
Josep Trueta Street, s/n  
Hospital Universitari General de Catalunya  
Vicenza  
Italy  
08195

**Type(s)**  
Scientific

**Contact name**  
Dr Cristina Valles

**Contact details**  
calle Josep tureta  
Barcelona  
Spain  
08190

## Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

**Secondary identifying numbers**  
PER-ECL-2015-03

## Study information

**Scientific Title**  
Influence of suturing technique and postoperative palatal flap thickness on wound healing and patient morbidity after connective tissue harvesting. A randomized controlled clinical trial

**Study objectives**

1. Wound healing is correlated to the suturing techniques
2. The post-operative palatal thickness is correlated with the early wound healing of the donor site
3. Palatal thickness is correlated with postoperative pain

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Comitè Etic d'Investigacio Clinica, 20/11/2015, ref: PER-ECL-2015-03

**Study design**

Single centre randomized clinical trial

**Primary study design**

Interventional

**Secondary study design**

Randomised controlled trial

**Study setting(s)**

Hospital

**Study type(s)**

Prevention

**Participant information sheet****Health condition(s) or problem(s) studied**

Palatal healing

**Interventions**

Using a computerized randomisation method, participants are allocated to one of two groups.

Group 1: Participants receive a continuous interlocking suture used in the donor site

Group 2: Participants receive a criss-cross suture used in the donor site.

The envelope indicating the group to which the patient belongs was opened at the time of the suture. Before the surgery measurement of the palate was taken with an endodontic file and flowable light curing composite. After the obtention of a standardized connective tissue graft (12mm x 8mm x 1,5mm) using the single incision technique, measurement of the residual flap thickness was taken with the same method explained before.

After one week, healing of the donor site is evaluated and pain perception recorded daily until 8 weeks after the intervention.

**Intervention Type**

Procedure/Surgery

**Primary outcome measure**

Association between suture technique and early healing index is measured one week after surgery and after suture removal, the early-wound healing index (EHI) using five different degrees:

1. Complete flap closure without fibrin line at the palate
2. Complete flap closure with fibrin line at the palate
3. Complete flap closure with small fibrin clot(s) at the palate
4. Incomplete flap closure with partial necrosis of the palatal tissue
5. Incomplete flap closure with complete necrosis of the palatal tissue (more than 50 % of the former flap is involved)

### **Secondary outcome measures**

1. Association between preoperative palatal thickness and early healing is measured one week after surgery using a early-wound healing index (EHI) (Fickl et al. 2014) and the preoperative measurement taken with the endodontic file and light curing flowable composite
2. Association between postoperative palatal flap thickness and early healing is measured one week after surgery using a early-wound healing index (EHI) (Fickl et al. 2014) and the postoperative measurement taken with the endodontic file and light curing flowable composite
3. Pain Perception of the patients was determined with VAS pain values daily until 8 weeks after surgery

### **Overall study start date**

20/09/2015

### **Completion date**

20/05/2017

## **Eligibility**

### **Key inclusion criteria**

1. Men and women over 18 years old
2. Showing Miller Class I,II or III (Miller 1985) single gingival recession

### **Participant type(s)**

Patient

### **Age group**

Adult

### **Lower age limit**

18 Years

### **Sex**

Both

### **Target number of participants**

40 patients total, 20 each group

### **Key exclusion criteria**

1. Any systemic conditions that might impact directly on the systemic inflammatory status (e.g. autoimmune diseases)

2. Severe cognitive or psychiatric disorders
3. Chronic immunosuppressant therapy
4. Lactating females and current pregnancy
5. Patients with coagulation disorders excluded from the trial
6. Smokers who consume more than 10 cigarettes per day
7. Patients using partial or full denture, which has any contact with the palate

**Date of first enrolment**

21/10/2015

**Date of final enrolment**

20/02/2017

## **Locations**

**Countries of recruitment**

Spain

**Study participating centre**

Universitat Internacional de Catalunya

Calle Josep Tureta s/n

Barcelona

Spain

08190

## **Sponsor information**

**Organisation**

Universitat Internacional de Catalunya

**Sponsor details**

Calle Josep Tureta

San Cugat del Valles

Spain

08195

+34 93 504 20 30

clinica@uic.es

**Sponsor type**

University/education

**ROR**

<https://ror.org/00tse2b39>

# Funder(s)

## Funder type

University/education

## Funder Name

Universitat Internacional de Catalunya

# Results and Publications

## Publication and dissemination plan

Planned publication in a high-impact reviewed journal

## Intention to publish date

15/07/2018

## Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

## IPD sharing plan summary

Data sharing statement to be made available at a later date