

Low level laser treatment for masticatory muscle pain

Submission date 28/08/2015	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 10/09/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 04/11/2015	Condition category Signs and Symptoms	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Myofascial pain is a long-term pain disorder, in which pain, tenderness and muscle spasms are felt in facial muscles. The masticatory muscles include all of the muscles that are responsible for movement of the jaw. These muscles are often the most affected in cases of myofascial pain, as their involvement in chewing (mastication) means that they are used a lot. Two of the most common points where this pain is felt are the masseter muscle, which is the main muscle involved in chewing, and the temporalis muscle, which is involved in closing the mouth and moving the mouth from side to side when chewing. Low-level laser therapy (LLLT) is a treatment which uses specific wavelengths of light to stimulate the cells in the body that repair tissue, reduce inflammation, and reduce pain. This non-invasive treatment has proven to be very effective for treating back and neck pain, as well as other long term pain conditions. The aim of this study is to investigate whether applying LLLT to the point of greatest pain would help to reduce masticatory muscle pain.

Who can participate?

Adults with a diagnosis of myofascial pain and natural posterior occlusion (the teeth at the back of the mouth are aligned correctly).

What does the study involve?

Participants are randomly allocated into one of three groups. For participants in the first group, the low level laser is applied precisely and continuously to the places where pain is greatest in the related muscle. For participants in the second group, the laser is applied to three previously chosen points on the masseter muscle and three points on the temporalis muscle. For participants in the third group, the laser device is switched on, but not programmed to target any particular points. Levels of pain and movement range are assessed before and after the laser treatment.

What are the possible benefits and risks of participating?

Not provided at time of registration

Where is the study run from?

Istanbul University Faculty of Dentistry Department of Prosthodontics (Turkey)

When is the study starting and how long is it expected to run for?
September 2010 to October 2011

Who is funding the study?
Istanbul University Faculty of Dentistry Department of Prosthodontics (Turkey)

Who is the main contact?
Dr Bilge Gokcen-Rohlig

Contact information

Type(s)
Scientific

Contact name
Dr Bilge Gokcen-Rohlig

ORCID ID
<http://orcid.org/0000-0003-3143-9668>

Contact details
Istanbul University Faculty of Dentistry Department of Prosthodontics
Capa
Istanbul
Türkiye
34390

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N/A

Study information

Scientific Title
Early results of low-level laser application for masticatory muscle pain: a double-blind randomized clinical study

Study objectives
Direct application of low-level laser therapy (LLLT) at the point of greatest pain would reduce muscle pain to a greater degree than would application at pre-established points and placebo.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee of the Istanbul University Faculty of Medicine , 10/05/2012, ref: 2011/1272-639

Study design

Single-centre randomised controlled trial.

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Masticatory pain

Interventions

Participants were divided randomly into three age- and sex-matched groups:

1. Laser group (LGI) in which low level laser is applied precisely and continuously to the greatest points of pain in the related muscle (masseter and/or temporalis).
2. Laser group II (LGII) in which LLLT is applied in the same manner to three predetermined points on the masseter muscle (superior [MS], middle [MM], and inferior [MI] points) and three points on the temporalis muscle (anterior [TA], middle, and posterior points)
3. Placebo group (PG) in which the laser device was switched on, but not programmed.

Intervention Type

Device

Primary outcome measure

Mandibular movement range (right, left and protrusion) before and after the assigned treatments, pain on palpation by masseter muscles on three points (superior, middle and inferior points), temporals muscles on three points (anterior, middle and posterior points) and pressure pain thresholds for the same points by algometer.

Secondary outcome measures

Pressure pain thresholds of the examined muscles are investigated before and after each treatment via visual analogue score (VAS), before any after the treatments

Overall study start date

01/09/2010

Completion date

01/10/2011

Eligibility

Key inclusion criteria

1. Diagnosis of myofascial pain according to the Research Diagnostic Criteria for Temporomandibular Disorder (RDC/TMD)
2. Aged between 18–60 years
3. Natural posterior occlusion

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

30

Key exclusion criteria

1. Disc displacement with reduction or without reduction with or without limited opening, arthralgia, arthritis, or arthrosis
2. General inflammatory connective tissue disease (e.g. rheumatoid arthritis)
3. Psychiatric disorder
4. Tumour
5. Heart disease or pacemaker
7. Pregnancy
8. Symptoms that could be referred to other orofacial region diseases (e.g. toothache, neuralgia, migraine)
9. Treatment or medication use for headache or bruxism in the last 2 years
10. Local skin infection over the masseter muscle

Date of first enrolment

01/09/2010

Date of final enrolment

01/10/2011

Locations

Countries of recruitment

Türkiye

Study participating centre

Istanbul University Faculty of Dentistry Department of Prosthodontics

Istanbul universitesi dis hek. fak. cene yuz protezi 6.kat

Capa

Istanbul

Türkiye

34390

Sponsor information

Organisation

Istanbul University Faculty of Dentistry Department of Prosthodontics

Sponsor details

Istanbul University Faculty of Dentistry Department of Prosthodontics

Capa

Istanbul

Türkiye

34390

Sponsor type

Hospital/treatment centre

ROR

<https://ror.org/03a5qrr21>

Funder(s)

Funder type

University/education

Funder Name

Istanbul University Faculty of Dentistry Department of Prosthodontics

Results and Publications

Publication and dissemination plan

Manuscript has been submitted to BMC Oral Health.

Intention to publish date**Individual participant data (IPD) sharing plan**

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	23/10/2015		Yes	No