

# A study on side effects such as nausea and vomitus after medication to prevent excessive bleeding after a caesarean section

<b>Submission date</b> 03/02/2013	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
<b>Registration date</b> 28/02/2013	<b>Overall study status</b> Completed	<input type="checkbox"/> Protocol
<b>Last Edited</b> 07/03/2018	<b>Condition category</b> Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

At the end of a cesarean section in Belgium, in general some medication is given to the mother to prevent excessive bleeding. The two products used most frequently are called carbetocin and oxytocin and work well. No difference has been demonstrated as far as the effect on prevention of bleeding is concerned. A major side effect of this kind of medication is nausea and vomitus and a lowering of blood pressure resulting in dizziness. The aim of the present study is to compare nausea, vomiting and changes in blood pressure between carbetocin and oxytocin.

### Who can participate?

Any pregnant woman for whom a caesarean section is planned in a term ( full grown) pregnancy without any other complication can participate.

### What does the study involve?

Participants are randomly allocated to one of two groups. All women receive an intravenous medication to prevent bleeding and no one (participant, operating or treating team) will not know whether the medication used is carbetocin or oxytocin.

### What are the possible benefits and risks of participating?

There is no difference with routine management during which both medications are used. The results of this study will help future patients to receive the product with the fewest side effects.

### Where is the study run from?

The study will be run in the Obstetrics Ward at Antwerp University Hospital UZA, Belgium.

### When is the study starting and how long is it expected to run for?

The study will be running for one year and will start in March 2013.

### Who is funding the study?

This study is not funded by any pharmaceutical company, all costs are taken care of by the local Fund for perinatal Medicine of the Department of Obstetrics at Antwerp University Hospital

Who is the main contact?  
Prof Dr Yves Jacquemyn  
Yves.jacquemyn@uza.be

## Contact information

### Type(s)

Scientific

### Contact name

Prof Yves Jacquemyn

### Contact details

Wilrijkstraat 10

Edegem

Belgium

2650

-

yves.jacquemyn@uza.be

## Additional identifiers

### Protocol serial number

OBGYN-1

## Study information

### Scientific Title

OBGYN-1: A randomized controlled trial on nausea, vomitus and blood pressure changes comparing carbetocine versus oxytocine in the prevention of hemorrhage after caesarean section

### Acronym

NAVOCASY

### Study objectives

Single shot intravenous carbetocin is accompanied with more nausea and vomitus as compared to intravenous oxytocin.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Ethics Committee Ethisch Comité Universitair Zieknhusi Antwerpen, 21/02/2011, ref: B300201110299

### Study design

Prospective randomized controlled trial

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Pregnancy and delivery

**Interventions**

Standard intravenous carbetocin vs standard intravenous oxytocin

**Intervention Type**

Drug

**Phase**

Not Applicable

**Drug/device/biological/vaccine name(s)**

Carbetocin, oxytocin

**Primary outcome(s)**

From the moment the participant enters the operating theatre until one hour later, blood pressure and heart frequency are carefully registered every 5 minutes or more as during routine monitoring of surgery. Nausea and vomitus are evaluated before surgery (just before incision), during surgery: before giving the intravenous medication and after 5, 10 and 15 minutes based on the standardized scale:

0: no nausea, no vomitus

1: mild nausea, no vomiting

2: nausea but with retching, no vomiting

3: nausea and vomiting

**Key secondary outcome(s)**

Any other complaint

**Completion date**

15/02/2014

**Eligibility****Key inclusion criteria**

Term uncomplicated pregnant women who undergo a planned cesarean section

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

Female

**Key exclusion criteria**

1. Unplanned caesarean section
2. Complications including: hypertension, preterm labour, any gastrointestinal or cardiovascular disease necessitating treatment

**Date of first enrolment**

15/02/2013

**Date of final enrolment**

15/02/2014

## Locations

**Countries of recruitment**

Belgium

**Study participating centre**

Antwerp University Hospital

Edegem

Belgium

2650

## Sponsor information

**Organisation**

Antwerp University Hospital (Belgium)

**ROR**

<https://ror.org/01hwamj44>

## Funder(s)

**Funder type**

Hospital/treatment centre

**Funder Name**

Department of Obstetrics, Antwerp University Hospital (Belgium)

# Results and Publications

## Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	02/01/2018		Yes	No