# Early Diabetes Intervention Trial

Submission date	<b>Recruitment status</b> No longer recruiting	Prospectively registered		
19/06/2002		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
19/06/2002	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
06/01/2011	Nutritional. Metabolic. Endocrine			

## Plain English summary of protocol

Not provided at time of registration

## Contact information

## Type(s)

Scientific

### Contact name

Dr Rury R Holman

#### Contact details

Diabetes Trials Unit OCDEM, Churchill Hospital Old Road, Headington Oxford United Kingdom OX3 7LJ

## Additional identifiers

### Protocol serial number

Study Numbers:Bay g 5421/0740 and Bay g 5421/200031

## Study information

Scientific Title

### **Acronym**

**EDIT Study** 

**Study objectives** 

The Early Diabetes Intervention Trial (EDIT) was a six-year, prospective, randomised, double-blind, placebo-controlled study in subjects thought to be at increased risk of developing diabetes, and who had two consecutive fasting plasma glucose levels in the range 5.5 to 7.7 mmol/L. The primary aim of the trial was to determine whether deterioration in glycaemic tolerance towards diabetes could be delayed or prevented using an alpha-glucosidase inhibitor (acarbose) or a biguanide (metformin). The trial results were presented at the 2003 Diabetes UK meeting.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Not provided at time of registration

### Study design

Prospective, randomised, double-blind, placebo-controlled study

### Primary study design

Interventional

### Study type(s)

**Not Specified** 

### Health condition(s) or problem(s) studied

Impaired glucose tolerance

#### Interventions

Subjects valid for inclusion were randomly assigned to acarbose 50 mg tds, metformin 500 mg tds, a combination of the two, or matching placebo.

### Intervention Type

Drug

#### Phase

**Not Specified** 

## Drug/device/biological/vaccine name(s)

acarbose, metformin

### Primary outcome(s)

Not provided at time of registration

### Key secondary outcome(s))

Not provided at time of registration

### Completion date

31/12/2003

## **Eligibility**

### Key inclusion criteria

Subjects aged 30 to 70 years inclusive, with fasting hyperglycaemia; defined as a fasting plasma glucose (FPG) level  $\geq$ 5.5 mmol/l and <7.8 mmol/l on two occasions at least 1 week apart

### Participant type(s)

**Patient** 

## Healthy volunteers allowed

No

### Age group

Adult

### Sex

All

### Key exclusion criteria

Not provided at time of registration

### Date of first enrolment

01/01/1997

### Date of final enrolment

31/12/2003

## Locations

### Countries of recruitment

**United Kingdom** 

England

# Study participating centre Diabetes Trials Unit

Oxford United Kingdom OX3 7LJ

## Sponsor information

### Organisation

Bayer plc and Merck-Lipha (UK)

## Funder(s)

### Funder type

Industry

### Funder Name

Educational grants were provided by Bayer plc and Merck-Lipha (UK)

## **Results and Publications**

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration

### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/06/2003		Yes	No
Results article	Hyper Glycaemia results	01/01/2009		Yes	No
Abstract results				No	No
Abstract results				No	No
Abstract results	abstract 450-P	01/06/2000		No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes